


# **Does VADER sentiment analysis produce useful data for suicide prevention outreach on Twitter?**

A Case Study

# The Data We Know


Distribution of Twitter users in the United States as of December 2016, by age group



Source  
comScore  
© Statista 2018

Additional Information:  
United States; comScore; December 2016; 18 years and older


Suicide Rates by Age from 2000 to 2016



Highcharts.com

# Which Data Could Help National and Local Orgs?

Best social media platforms for influence marketing according to influencers in the United States as of July 2016



Sources  
eMarketer; SheSpeaks  
© Statista 2018

Additional Information:  
United States; SheSpeaks; July 2016; 347 respondents; among influencers with an active blog and who were act



## American Foundation for Suicide Prevention

@afspnational

Preventing suicide through research, advocacy, and education. Account not monitored 24/7. Thoughts of suicide? Call: 800-273-8255 / Text: 741741

📍 USA

🌐 [afsp.org](https://afsp.org)

📅 Joined January 2009



## SFSuicidePrevention

@SFSuicide

We're a 24-hour suicide prevention hotline available to anyone in emotional crisis. If you need help or someone you know is in crisis, please call 415.781.0500

📍 San Francisco

🌐 [sfsuicide.org](https://sfsuicide.org)

📅 Joined October 2011



# Project Overview & Goals

**Goal:** To create actionable data (maps, keywords, trends) for suicide and crisis prevention organizations to target ads to Twitter users experiencing emotional distress.

---

**Data source:** English language tweets (as selected by Twitter) from several US States

**Cleaning & parsing data:** Created a list of words and phrases that indicate emotional stress and distress, cleaned raw csv pulled from Twitter filtering by distress list, human filter; created dataframes, re-encoded csvs to UTF-8.

**Sentiment Analysis Analyzer:** VADER

**Visualization:** WordCloud and statistical graphics



# Assumptions & Caveats

Sample size and time series are not large enough to make definitive judgments from data.

VADER does not pick up on sarcasm.

(Ex: 'Warren Beatty has never been so entertaining.' compound: 0.5777, neg: 0.0, neu: 0.616, pos: 0.384)

VADER is not great with simultaneously reading polarity and strength and slang.

(Ex: 'VADER sentiment analysis is the s\*\*\*.' compound: 0.6124, neg: 0.0, neu: 0.556, pos: 0.444)

We don't know how time and day are related to Twitter usage or sentiment. There may be a day/time that is more reflective of suicidal sentiments and better suited for targeted ads.



# Data Gathering and Clean-up

- Twitter is a poor source of large data sets
  - Limited volume per call
  - No historical data sets for free
- Twitter data requires a lot of cleaning
  - Not uniform data
  - Significant filtering required
- Public data is already presented in summary format
  - Scraping not feasible
  - Purpose-built summaries



# Exploration of Data


- Researched Trigger Words
- Filtered Tweets
- Measured Sentiment
- Discovered our hypothesis was incorrect
  - Led us to explore why...

“The word ibuprofen was 16 times more likely to predict the person texting would need emergency services than the word suicide.”


- <https://mashable.com/2017/03/09/artificial-intelligence-suicide-risk/#HikouaxJEmqi>

# Exploration and Analysis

Sentiment Analysis on Suicidal Tweets - 2009 1.6m Tweets USA



Sentiment Analysis on Suicidal Tweets - California









**Let's Go to JUPYTER NOTEBOOK!**



# Conclusions and Implications

Sentiment analysis of tweets with “suicide trigger words” may not produce useful information for local and national suicide prevention twitter outreach . *Derived from 4 pie and 1 bar chart.*

- Twitter may not be the preferred social medium for expressing suicidal inclinations.
- The data is likely inconclusive due to the small time frame.
- There may be a better sentiment analyzer (textblob, SlangSD, etc.)
- There is evidence that specific words, and not sentiments, is a better indicator (ex. ibuprofen) of potential self-harm



# References

Gilbert, CJ Hutto Eric. "Vader: A parsimonious rule-based model for sentiment analysis of social media text." *Eighth International Conference on Weblogs and Social Media (ICWSM-14)*. Available at (2018/04/16) [http://comp. social. gatech. edu/papers/icwsm14.vader. hutto. pdf](http://comp.social.gatech.edu/papers/icwsm14.vader.hutto.pdf). 2014.

American Foundation for Suicide Prevention. Available at (2018/04/16) <https://afsp.org/about-suicide/suicide-statistics/>

Statista. "Distribution of Twitter users in the United States as of December 2016, by age group" Available at (2018/04/16) <https://www.statista.com/statistics/192703/age-distribution-of-users-on-twitter-in-the-united-states/>

Statista. "Best social media platforms for influence marketing according to influencers in the United States as of July 2016." Available at (2018/04/16) <https://www.statista.com/statistics/617525/best-social-media-for-influence-marketing-per-influencers/>

Crisis Trends. " Co-occurring issues. Texters in all states experiencing anxiety/stress." [www.crisistrends.org](http://www.crisistrends.org)



**Thank you**