

# CONVERGENCE OF A MODIFIED NEWTON METHOD FOR A MATRIX POLYNOMIAL EQUATION ARISING IN STOCHASTIC PROBLEM

SANG-HYUP SEO, JONG-HYEON SEO, AND HYUN-MIN KIM<sup>†</sup>

**ABSTRACT.** We consider the Newton iteration for a matrix polynomial equation which arises in stochastic problem. In this paper, we show that the elementwise minimal nonnegative solution of the matrix polynomial equation can be obtained using Newton's method if the equation satisfies the sufficient condition, and the convergence rate of the iteration is quadratic if the solution is simple. Moreover, we show that the convergence rate is at least linear if the solution is non-simple, but we can apply a modified Newton method whose iteration number is less than the pure Newton iteration number. Finally, we give a numerical experiment which is related with our issue.

## 1. INTRODUCTION

We consider a matrix polynomial equation(MPE) with  $n$ -degree defined by

$$(1.1) \quad P(X) = \sum_{k=0}^n A_k X^k = A_n X^n + A_{n-1} X^{n-1} + \cdots + A_1 X + A_0 = 0,$$

where the coefficient matrices  $A_k$ 's are  $m \times m$  matrices. Then, the unknown matrix  $X$  must be an  $m \times m$  matrix.

The MPE (1.1) often occurs in the theory of differential equations, system theory, network theory, stochastic theory, quasi-birth-and-death and other areas [1–4, 7, 14, 21–23].

Davis [5, 6] and Higham, Kim [15, 16] studied the Newton method for a quadratic matrix equation. Guo and Laub [11] considered a nonsymmetric algebraic Riccati equation, and they proposed iteration algorithms which converge to the minimal positive solution. In [8], Guo provided a sufficient condition for the existence of non-negative solutions of nonsymmetric algebraic Riccati equations. Kim [20] showed that the minimal positive solutions also can be found by the Newton method with the zero initial matrices in some different types of quadratic equations. Hautphenne, Latouche, and Remiche [12] studied the Newton method for the Markovian binary tree.

Seo and Kim [26, 27] studied the Newton iteration for a quadratic matrix equation and a matrix polynomial equation. Specially, in [26], they provided a relaxed Newton method whose convergence is faster than the pure one. Guo and Lancaster [10] analyzed and provided a modification about Newton's method for algebraic

---

2010 *Mathematics Subject Classification.* 65H10.

*Key words and phrases.* matrix polynomial equation, elementwise positive solution, elementwise nonnegative solution,  $M$ -matrix, Newton's method, convergence rate, acceleration of a method.

<sup>†</sup>Corresponding Author.

This research was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science and Technology(2017R1A5A1015722, 2017R1D1A3B04033516).

Riccati equations. They showed that the modification of Newton's method is better than the pure one if the minimal nonnegative solution is non-simple.

**Assumption 1.1.** For the MPE (1.1),

- 1) The coefficient matrices  $A_k$ 's are nonnegative except  $A_1$ .
- 2)  $-A_1$  is a nonsingular  $M$ -matrix.
- 3)  $A_0$ ,  $A_1$ , and  $\sum_{k=2}^n A_k$  are irreducible.

The goal of this paper is to propose a modified Newton method of the MPE (1.1) which satisfies Assumption 1.1. This MPE is useful for stochastic theory, quasi-birth-and-death area, and so on. The modified Newton method is better than the pure Newton's method if the elementwise minimal positive solution is non-simple. The idea of the modified Newton method is from the modification of Newton's method for algebraic Riccati equations of [10]. In [10], Guo and Lancaster showed that  $\|Y_{i+1} - S\| < c\varepsilon$  for the modified iteration  $Y_{i+1}$ , the solution  $S$ , a constant  $c > 0$ , and small  $\varepsilon > 0$ . On the other hand, we show that the modified Newton iteration  $Y_{i+1}$  for the MPE is closer to the solution  $S$  than the pure Newton iteration  $X_{i+1}$ .

We start with some basic definitions.

**Definition 1.2.** Let a matrix  $A \in \mathbb{R}^{m \times m}$ .  $A$  is an  $Z$ -matrix if all its off-diagonal elements are nonpositive.

It is clear that any  $Z$ -matrix  $A$  can be written as  $sI - B$  with  $B \geq 0$  and  $s \in \mathbb{R}$ . Then  $M$ -matrix can be defined as follows.

**Definition 1.3.** A matrix  $A \in \mathbb{R}^{m \times m}$  is an  $M$ -matrix if  $A = rI - B$  for some nonnegative matrix  $B$  with  $r \geq \rho(B)$  where  $\rho$  is the spectral radius; it is a singular  $M$ -matrix if  $r = \rho(B)$  and a nonsingular  $M$ -matrix if  $r > \rho(B)$ .

The following result is well known and can be found in [9] and [25] for example.

**Theorem 1.4.** For a  $Z$ -matrix  $A$ , the following are equivalent:

- (1)  $A$  is a nonsingular  $M$ -matrix.
- (2)  $A^{-1}$  is nonnegative.
- (3)  $Av > 0$  for some vector  $v > 0$ .
- (4) All eigenvalues of  $A$  have positive real parts.

**Definition 1.5.** A positive solution  $S_1$  of the matrix equation  $P(X) = 0$  is the *elementwise minimal positive solution* and a positive solution  $S_2$  of  $P(X) = 0$  is the *elementwise maximal positive solution* if, for any positive solution  $S$  of  $P(X)$ ,

$$(1.2) \quad S_1 \leq S \leq S_2.$$

Similarly, if nonnegative solutions  $S_1$  and  $S_2$  satisfy (1.2) for any nonnegative solution  $S$ ,  $S_1$  is called the *elementwise minimal nonnegative solution* and  $S_2$  is called the *elementwise maximal nonnegative solution*.

**Definition 1.6.** [18, Definition 4.2.1, Definition 4.2.9] The *Kronecker product* of  $A = [a_{ij}] \in \mathbb{C}^{m \times n}$  and  $B = [b_{ij}] \in \mathbb{C}^{p \times q}$  is denoted by  $A \otimes B$  and is defined to be the block matrix

$$A \otimes B = \begin{bmatrix} a_{11}B & \cdots & a_{1n}B \\ \vdots & \ddots & \vdots \\ a_{m1}B & \cdots & a_{mn}B \end{bmatrix} \in \mathbb{C}^{mp \times nq}.$$

The vec operator  $\text{vec} : \mathbb{C}^{m \times n} \rightarrow \mathbb{C}^{mn}$  is defined by

$$\text{vec}(A) = [\mathbf{a}_1^T \quad \mathbf{a}_2^T \quad \cdots \quad \mathbf{a}_n^T]^T,$$

where  $\mathbf{a}_i^T = [a_{1i} \quad a_{2i} \quad \cdots \quad a_{ni}]$ .

**Lemma 1.7.** [18, Lemma 4.3.1] *Let  $A \in \mathbb{C}^{m \times n}$ ,  $B \in \mathbb{C}^{p \times q}$ , and  $C \in \mathbb{C}^{m \times q}$  be given and let  $X \in \mathbb{C}^{n \times p}$  be unknown. The matrix equation*

$$(1.3) \quad AXB = C$$

*is equivalent to the system of  $qm$  equations in  $np$  unknowns given by*

$$(1.4) \quad (B^T \otimes A)\text{vec}(X) = \text{vec}(C),$$

*that is,  $\text{vec}(AXB) = (B^T \otimes A)\text{vec}(X)$ .*

**Definition 1.8.** Let a matrix function  $F : \mathbb{C}^{m \times n} \rightarrow \mathbb{C}^{m \times n}$  be given, and let a matrix equation

$$(1.5) \quad F(X) = 0$$

be given. Then, a solution  $S \in \mathbb{C}^{m \times n}$  of (1.5) is called *simple* if the Fréchet derivative at  $S$  is nonsingular.

To reach our goal, in Section 2, we study the minimal nonnegative solution  $S$  of (1.1) and the Fréchet derivative of  $P$  in (1.1) at  $S$ , and we show the convergence of the Newton iteration for (1.1). We give an analysis about Newton's method for the non-simple minimal nonnegative solution  $S$ , in Section 3. In Section 4, we propose a modified Newton method which is better for finding the minimal nonnegative solution  $S$ . Finally, we give some numerical experiments, in Section 5.

For convenience, the notation  $\|\cdot\|$  is used instead of the Frobenius norm  $\|\cdot\|_F$  and  $\mathbb{N}_0$  is used as  $\mathbb{N} \cup \{0\}$  because the Frobenius norm and  $\mathbb{N}_0$  are used very frequently in this paper.

## 2. CONVERGENCE OF NEWTON'S METHOD FOR AN MPE

In this section, we introduce a sufficient condition of the existence of the minimal nonnegative solution of the MPE (1.1) with Assumption 1.1, and give some analysis for Newton's method.

**Theorem 2.1.** [24, Theorem 2.1] *Let the MPE (1.1) with 1) and 2) in Assumption 1.1 be given. Then, there exists the minimal nonnegative solution if*

$$(2.1) \quad B = -\sum_{k=0}^n A_k \text{ is a nonsingular or singular irreducible } M\text{-matrix}.$$

The Fréchet derivative of the matrix polynomial equation (1.1) at  $X$  in the direction  $H$  is given by

$$(2.2) \quad P'_X(H) = \sum_{k=1}^n \sum_{l=0}^{k-1} A_k X^l H X^{k-l-1}.$$

The second Fréchet derivative of the quadratic matrix equation (1.1) at  $X$  is given by

$$(2.3) \quad P''_X(K, H) = \sum_{k=2}^n \sum_{l=0}^{k-2} \sum_{j=0}^l A_k (X^l H X^j K X^{n-l-j-2} + X^l K X^j H X^{n-l-j-2}).$$

For the equation (1.1), each step of the Newton iteration with given  $X_0$  can be written as

$$(2.4) \quad X_{i+1} = X_i - P'_{X_i}{}^{-1}(P(X_i))$$

if  $P'_{X_i}$  is invertible for all  $i \in \mathbb{N}_0$ .

(2.4) can be separated into two parts as

$$(2.5) \quad \begin{cases} P'_{X_i}(H_i) = -P(X_i), \\ X_{i+1} = X_i + H_i, \end{cases} \quad i = 1, 2, \dots$$

The general approach for solving (2.5) is to solve the  $m^2 \times m^2$  linear system derived by Lemma 1.7 such as

$$\mathcal{P}'_{X_i} \text{vec}(H_i) = \text{vec}(-P(X_i)),$$

where

$$(2.6) \quad \mathcal{P}'_X = \sum_{k=1}^n \sum_{l=0}^{k-1} (X^{k-l-1})^T \otimes A_k X^l.$$

**Theorem 2.2.** *Suppose that the MPE (1.1) satisfies Assumption 1.1 and (2.1). Then, the Newton sequence  $\{X_i\}$  with  $X_0 = 0$  is well defined, is monotone non-decreasing, and converges to the elementwise minimal positive solution  $S$ . Furthermore,  $-\mathcal{P}'_{X_i}$  is a nonsingular irreducible  $M$ -matrix for  $i \in \mathbb{N}$ , and  $-\mathcal{P}'_S$  is an irreducible  $M$ -matrix.*

*Proof.* According to the proof of [24, Theorem 2.1], the elementwise minimal non-negative solution  $S$  of (1.1) is the limit of the monotone nondecreasing sequence  $\{X_i^G\}_{i=0}^\infty$  which is defined by

$$\begin{cases} X_{i+1}^G = G(X_i^G), \\ X_0^G = 0, \end{cases}$$

where

$$G(X) = -A_1^{-1} \left( \sum_{k=2}^n A_k X^k + A_0 \right).$$

Since  $X_1^G = -A_1^{-1} A_0 > 0$ , the solution  $S$  is also positive. Thus,  $S \in \{Y \in \mathbb{R}^{m \times m} | Y > 0, F(Y) \leq 0\}$ . From [26, Theorem 2.9], the Newton sequence  $\{X_i\}$  with  $X_0 = 0$  is well-defined, monotone nondecreasing, and converges to the elementwise minimal positive solution  $S$ . Moreover,

$$-\mathcal{P}'_{X_i} = - \sum_{k=1}^n \sum_{l=0}^{k-1} (X_i^{k-l-1})^T \otimes A_k X_i^l$$

is a nonsingular  $M$ -matrix for each  $i \in \mathbb{N}_0$ .

Now, it is sufficient to show that  $-\mathcal{P}'_{X_i}$  is irreducible for  $i \in \mathbb{N}$ . Since  $X_1 = -A_1^{-1} A_0 > 0$  and  $A_p \geq 0$  for all  $p \geq 2$ ,

$$\sum_{k=2}^n \sum_{l=0}^{k-1} (X_i^{k-l-1})^T \otimes A_k X_i^l \geq \sum_{k=2}^n (X_i^{k-1})^T \otimes A_k \geq 0.$$

From that  $X_i > 0$ , we obtain that  $\sum_{k=2}^n (X_i^{k-1})^T \otimes A_k$  is irreducible if and only if  $\sum_{k=2}^n \mathbf{1}_{m \times m} \otimes A_k = \mathbf{1}_{m \times m} \otimes (\sum_{k=2}^n A_k)$  is irreducible. Since  $\sum_{k=2}^n A_k$  is irreducible,

$$\sum_{k=2}^n \sum_{l=0}^{k-1} (X_i^{k-l-1})^T \otimes A_k X_i^l$$

is an irreducible nonnegative matrix. Therefore,

$$-\mathcal{P}'_{X_i} = -I_m \otimes A_1 - \sum_{k=2}^n \sum_{l=0}^{k-1} (X_i^{k-l-1})^T \otimes A_k X_i^l$$

is also irreducible.  $\square$

The next theorem follows directly from Theorem 2.2 and the well known local quadratic convergence of Newton's method.

**Theorem 2.3.** *If the matrix  $-\mathcal{P}'_S$  in Theorem 2.2 is nonsingular, then for  $X_0 = 0$ , the Newton sequence  $\{X_i\}$  converges to  $S$ , quadratically.*

### 3. ANALYSIS FOR THE SINGULAR $M$ -MATRIX $-\mathcal{P}'_S$

According to Theorem 2.3, the Newton iteration (2.4) converges quadratically if  $-\mathcal{P}'_S$  is nonsingular. In this section, we will see the convergence rate of (2.4) when  $-\mathcal{P}'_S$  is a singular  $M$ -matrix. If  $P'_S$  is not invertible, then  $P'_S$  has a null space  $\mathcal{N} = \text{Ker}(P'_S)$  and closed range  $\mathcal{M} = \text{Im}(P'_S)$ . Suppose that the direct sum  $\mathcal{N} \oplus \mathcal{M} = \mathbb{R}^{m \times m}$ . Then, we can define  $\mathbf{P}_{\mathcal{N}}$  to be the projection onto  $\mathcal{N}$  parallel to  $\mathcal{M}$  and  $\mathbf{P}_{\mathcal{M}} = I - \mathbf{P}_{\mathcal{N}}$ . For a nonzero matrix  $N_0 \in \mathcal{N}$ , define the map  $\mathcal{B}_{N_0} : \mathcal{N} \rightarrow \mathcal{N}$  given by

$$(3.1) \quad \mathcal{B}_{N_0}(N) = \mathbf{P}_{\mathcal{N}} P''_S(N_0, N).$$

In fact,  $\mathcal{B}_{N_0}$  is a linear map. The main result of this section is an application of the following theorem which shows the local convergence and the convergence rate of Newton's method under some conditions.

**Theorem 3.1.** (cf. [10, Theorem 1.5], [19, Theorem 1.1]) *Let  $\mathcal{B}_{N_0}$  in (3.1) be invertible for some nonzero  $N_0 \in \mathcal{N}$ ,  $\mathcal{N} = \text{span}\{N_0\} \oplus \mathcal{N}_1$  for some subspace  $\mathcal{N}_1$ , and let*

$$(3.2) \quad W(\rho, \theta, \eta) = \left\{ X \left| \begin{array}{l} 0 < \|X - S\| < \rho, \|\mathbf{P}_{\mathcal{M}}(X - S)\| \leq \theta \|\mathbf{P}_{\mathcal{N}}(X - S)\|, \\ \|(\mathbf{P}_{\mathcal{N}} - \mathbf{P}_0)(X - S)\| \leq \eta \|\mathbf{P}_{\mathcal{N}}(X - S)\| \end{array} \right. \right\},$$

where  $\mathbf{P}_0$  is the projection onto  $\text{span}\{N_0\}$  parallel to  $\mathcal{N}_1 \oplus \mathcal{M}$ . If  $X_0 \in W(\rho_0, \theta_0, \eta_0)$  for  $\rho_0, \theta_0, \eta_0$  sufficiently small, then the Newton sequence  $\{X_i\}$  is well defined and  $\|P'^{-1}_{X_i}\| \leq c \|X_i - S\|^{-1}$  for all  $i \geq 1$  and some constant  $c > 0$ . Moreover,

$$\lim_{i \rightarrow \infty} \frac{\|X_{i+1} - S\|}{\|X_i - S\|} = \frac{1}{2}, \quad \lim_{i \rightarrow \infty} \frac{\|\mathbf{P}_{\mathcal{M}}(X_i - S)\|}{\|\mathbf{P}_{\mathcal{N}}(X_i - S)\|^2} = 0.$$

To analyze convergence of Newton's method when  $-\mathcal{P}'_S$  is singular, we will show that (1.1) satisfies the conditions of Theorem 3.1. From this point, for convenience, we let  $\tilde{X}_i = X_i - S$ .

**Lemma 3.2.** *Suppose the matrix polynomial equation (1.1) satisfies Assumption 1.1. If the matrix  $-\mathcal{P}'_S$  is a singular  $M$ -matrix, then 0 is a simple eigenvalue of  $-\mathcal{P}'_S$ ,  $\mathcal{N} \oplus \mathcal{M} = \mathbb{R}^{m \times m}$ ,  $\mathcal{N}$  is one-dimensional and the map  $\mathcal{B}_{N_0}$  is invertible for some nonzero  $N_0 \in \mathcal{N}$ .*

*Proof.* Since  $S$  is positive and  $A_k$ 's are irreducible,  $-\mathcal{P}'_S$  is irreducible. Then, by Perron-Frobenius Theorem [17, Theorem 8.4.4], 0 is a simple eigenvalue of  $\mathcal{P}'_S$  with a positive eigenvector. Thus, we can find  $n^2$  linearly independent vectors  $\chi_1, \chi_2, \dots, \chi_{n^2}$  such that  $\chi_1 > 0$  and

$$(3.3) \quad \mathcal{X}^{-1} \mathcal{P}'_S \mathcal{X} = \begin{bmatrix} 0 & 0 \\ 0 & \mathcal{D} \end{bmatrix}, \text{ where } \mathcal{X} = \left[ \begin{array}{c|c|c|c} \chi_1 & \chi_2 & \cdots & \chi_{n^2} \end{array} \right]$$

and  $\mathcal{D}$  is an  $(n^2 - 1) \times (n^2 - 1)$  nonsingular matrix. By the same way, we also have a positive vector  $\psi$  such that  $\psi^T \mathcal{P}'_S = 0$ . Now,  $P'_S(N) = 0$  if and only if  $\mathcal{P}'_S \text{vec}(N) = 0$ . From (3.3),  $\mathcal{P}'_S \text{vec}(N) = 0$  if and only if  $\text{vec}(N) \in \text{span}(\chi_1)$ , in which case

we write  $N = a \text{vec}^{-1}(\chi_1)$  for some nonzero  $a \in \mathbb{R}$ . Thus,  $\mathcal{N} = \text{span}(\text{vec}^{-1}(\chi_1))$ . Similarly,  $\mathcal{M} = \text{span}(\text{vec}^{-1}(\chi_2), \dots, \text{vec}^{-1}(\chi_{n^2}))$ . Therefore,  $\mathcal{N}$  is one-dimensional and  $\mathbb{R}^{m \times m} = \mathcal{N} \oplus \mathcal{M}$ .

To prove the map  $\mathcal{B}_{N_0}$  is invertible for a nonzero matrix  $N_0 \in \mathcal{N}$ , we only need to show that

$$\mathbf{P}_{\mathcal{N}}(P_S''(N_0, N)) \neq 0,$$

for all nonzero  $N \in \mathcal{N}$  because  $\mathcal{B}_{N_0}$  is linear and  $\mathcal{N}$  is one-dimensional. Since  $\text{vec}^{-1}(\chi_1) > 0$  and  $S > 0$ , we have

$$\begin{aligned} P_S''(N_0, N) &= \sum_{k=2}^n \sum_{l=0}^{k-2} \sum_{j=0}^l A_k (S^l N_0 S^j N S^{n-l-j-2} + S^l N S^j N_0 S^{n-l-j-2}) \\ &= 2ab \sum_{k=2}^n \sum_{l=0}^{k-2} \sum_{j=0}^l A_k (S^l \text{vec}^{-1}(\chi_1) S^j \text{vec}^{-1}(\chi_1) S^{n-l-j-2}) \neq 0 \end{aligned}$$

where  $N = a \text{vec}^{-1}(\chi_1)$  and  $N_0 = b \text{vec}^{-1}(\chi_1)$ . Moreover,  $P_S''(N_0, N)$  is either positive or negative.

On the other hand,

$$\text{vec}(P_S''(N_0, N)) = k_1 \chi_1 + k_2 \chi_2 + \dots + k_{n^2} \chi_{n^2}$$

for some real numbers  $k_1, k_2, \dots, k_{n^2}$ . By Fundamental theorem of linear algebra in [28] and Lemma 6.3.10 in [17], we have

$$(3.4) \quad \psi^T \text{vec}(P_S''(N_0, N)) = k_1 \psi^T \chi_1.$$

Since  $P_S''(N_0, N)$  is either positive or negative and  $\psi$  is positive, the left side of (3.4) is also either positive or negative. So,  $k_1$  cannot be zero. Therefore,

$$\mathbf{P}_{\mathcal{N}}(P_S''(N_0, N)) = k_1 \text{vec}^{-1}(\chi_1) \neq 0.$$

□

**Lemma 3.3.** *Let  $S$  be the minimal positive solution of (1.1) with Assumption 1.1, and let  $\{X_i\}_{i=0}^\infty$  be a Newton sequence in (2.4). Then,*

$$\|P(X_i)\| \leq a \|\tilde{X}_i\|^2 + b \|\tilde{X}_i\| \|\tilde{X}_{i-1}\| + c \|\tilde{X}_{i-1}\|^2$$

for some positive real numbers  $a, b, c$ .

*Proof.* From Taylor's Theorem and putting  $S = X_{i-1} - \tilde{X}_{i-1}$ , we have

$$(3.5) \quad P(X_i) = P(S) + P'_S(X_i - S) + O(\|X_i - S\|^2),$$

and

$$0 = P(S) = P(X_{i-1} - \tilde{X}_{i-1}) = P(X_{i-1}) - P'_{X_{i-1}}(\tilde{X}_{i-1}) + O(\|\tilde{X}_{i-1}\|^2),$$

which is equivalent to

$$(3.6) \quad -P(X_{i-1}) + P'_{X_{i-1}}(\tilde{X}_{i-1}) = O(\|\tilde{X}_{i-1}\|^2).$$

From (2.5), we have

$$(3.7) \quad 0 = P'_{X_{i-1}}(X_i - X_{i-1}) + P(X_{i-1}),$$

and clearly

$$(3.8) \quad X_i - X_{i-1} = \tilde{X}_i - \tilde{X}_{i-1}.$$

If we subtract (3.7) from (3.5) and substitute (3.6), we obtain

$$\begin{aligned} P(X_i) &= P(S) + P'_S(\tilde{X}_i) - P(X_{i-1}) - P'_{X_{i-1}}(\tilde{X}_i - \tilde{X}_{i-1}) + O(\|\tilde{X}_i\|^2) \\ &= P'_S(\tilde{X}_i) - P'_{X_{i-1}}(\tilde{X}_i) + O(\|\tilde{X}_i\|^2) + O(\|\tilde{X}_{i-1}\|^2). \end{aligned}$$

Putting  $S = X_{i-1} - \tilde{X}_{i-1}$  in the previous equality,

$$\begin{aligned}
P(X_i) &= \sum_{k=1}^n \sum_{l=0}^{k-1} A_k (X_{i-1} - \tilde{X}_{i-1})^l \tilde{X}_i (X_{i-1} - \tilde{X}_{i-1})^{k-l-1} \\
&\quad - P'_{X_{i-1}}(\tilde{X}_i) + O(\|\tilde{X}_i\|^2) + O(\|\tilde{X}_{i-1}\|^2) \\
&= \sum_{k=1}^n \sum_{l=0}^{k-1} A_k X_{i-1}^l \tilde{X}_i X_{i-1}^{k-l-1} + O(\|\tilde{X}_i\| \|\tilde{X}_{i-1}\|) \\
&\quad - P'_{X_{i-1}}(\tilde{X}_i) + O(\|\tilde{X}_i\|^2) + O(\|\tilde{X}_{i-1}\|^2) \\
&= P'_{X_{i-1}}(\tilde{X}_i) + O(\|\tilde{X}_i\| \|\tilde{X}_{i-1}\|) - P'_{X_{i-1}}(\tilde{X}_i) + O(\|\tilde{X}_i\|^2) + O(\|\tilde{X}_{i-1}\|^2) \\
&= O(\|\tilde{X}_{i-1}\|^2) + O(\|\tilde{X}_i\| \|\tilde{X}_{i-1}\|) + O(\|\tilde{X}_i\|^2).
\end{aligned}$$

Since  $\|\cdot\|$  is a multiplicative matrix norm on  $\mathbb{R}^{m \times m}$ , we have required result.  $\square$

**Lemma 3.4.** *For any fixed  $\theta > 0$ , let*

$$\mathcal{Q} = \{i \in \mathbb{N} \cup \{0\} \mid \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| > \theta \|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|\}$$

where  $\{X_i\}$  is the Newton sequence in Theorem 2.2. Then, there exist an integer  $i_0$  and a constant  $c > 0$  such that  $\|\tilde{X}_i\| \leq c \|\tilde{X}_{i-1}\|^2$  for all  $i \geq i_0$  in  $\mathcal{Q}$ .

*Proof.* Using Taylor's Theorem and the fact that  $P'_S(\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)) = 0$ ,

$$(3.9) \quad P(X_i) = P(S) + P'_S(\tilde{X}_i) + O(\|\tilde{X}_i\|^2) = P'_S(\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)) + O(\|\tilde{X}_i\|^2).$$

Since  $P'_S|_{\mathcal{M}} : \mathcal{M} \rightarrow \mathcal{M}$  is invertible,  $\|P'_S(\mathbf{P}_{\mathcal{M}}(\tilde{X}_i))\| \geq c_1 \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\|$  for some constant  $c_1 > 0$ . For  $i \in \mathcal{Q}$ , we have

$$\|\tilde{X}_i\| \leq \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| + \|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\| \leq (\theta^{-1} + 1) \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\|.$$

Thus by (3.9),

$$(3.10) \quad \|P(X_i)\| \geq c_1 \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| - c_2 \|\tilde{X}_i\|^2 \geq c_1(\theta^{-1} + 1)^{-1} \|\tilde{X}_i\| - c_2 \|\tilde{X}_i\|^2.$$

On the other hand, from Lemma 3.3, we have

$$\|P(X_i)\| \leq c_3 \|\tilde{X}_i\|^2 + c_4 \|\tilde{X}_{i-1}\| \|\tilde{X}_i\| + c_5 \|\tilde{X}_{i-1}\|^2.$$

From (3.10) and the fact that  $X_i \neq S$  for any  $i$ , we have

$$c_1(\theta^{-1} + 1)^{-1} - c_2 \|\tilde{X}_i\| \leq c_3 \|\tilde{X}_i\| + c_4 \|\tilde{X}_{i-1}\| + c_5 \frac{\|\tilde{X}_{i-1}\|^2}{\|\tilde{X}_i\|}.$$

Since  $\tilde{X}_i$  converges to 0 by Theorem 2.2, we can find an  $i_0$  such that  $\|\tilde{X}_i\| \leq c \|\tilde{X}_{i-1}\|^2$  for all  $i \geq i_0$ .  $\square$

**Corollary 3.5.** *Assume that, for given  $\theta > 0$ ,  $\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| > \theta \|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|$  for all  $i$  large enough. Then  $X_i \rightarrow S$  quadratically.*

When  $P'_S$  is singular practically the Newton sequence converges linearly, according to the corollary we conclude that the error will generally be dominated by its  $\mathcal{N}$  component [10]. From Lemmas 3.2 and 3.4 we have the following main theorem.

**Theorem 3.6.** *If  $-P'_S$  is a singular  $M$ -matrix and the convergence rate of the Newton sequence  $\{X_i\}$  in Theorem 2.2 is not quadratic, then  $\|P'_{X_i}\| \leq c \|\tilde{X}_i\|^{-1}$  for all  $i \geq 1$  and some constant  $c > 0$ . Moreover,*

$$\lim_{i \rightarrow \infty} \frac{\|\tilde{X}_{i+1}\|}{\|\tilde{X}_i\|} = \frac{1}{2}, \quad \lim_{i \rightarrow \infty} \frac{\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\|}{\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|^2} = 0.$$

## 4. A MODIFIED NEWTON METHOD

Under the conditions in Theorem 3.6, the convergence rate of the Newton sequence is  $1/2$ . Furthermore,  $\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\|/\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|^2$  converges to 0, i.e.,  $\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i_0})\| < \varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i_0})\|$  holds for sufficiently small  $\varepsilon > 0$  and large integer  $i_0$  to make  $\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i_0})\| < 1$ . Intuitively, we understand that  $\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i_0})$  is almost terminated, and  $\{\tilde{X}_i\}_{i=i_0}^\infty$  is located near a one-dimensional subspace  $\mathcal{N}$ . Then, we will give a modified Newton method which has faster convergence than the pure one.

**Lemma 4.1.** *Let  $\{X_i\}$  be the Newton sequence in Theorem 2.2, and let the derivative  $P'_S$  be singular. Suppose that there exists  $i_0 \in \mathbb{N}_0$  such that  $i \geq i_0$  implies that*

$$(4.1) \quad \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| < \varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|,$$

for  $0 < \varepsilon \leq 1$ . Then,  $\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)$  is negative for  $i \geq i_0$ .

*Proof.* From the proof of Lemma 3.2,  $V_1 := \text{vec}^{-1}(\chi_1)$  is a positive basis for  $\mathcal{N}$ . Let  $c_{i,1}$  be a scalar such that  $\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) = c_{i,1}V_1$ . Suppose that  $c_{i,1} \geq 0$ . Then,  $\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) \geq 0$ . Since  $\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) + \mathbf{P}_{\mathcal{M}}(\tilde{X}_i) = \tilde{X}_i \leq 0$ , we obtain that  $0 \leq \mathbf{P}_{\mathcal{N}}(\tilde{X}_i) \leq -\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)$ . Thus,

$$(4.2) \quad \|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\| \leq \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| < \varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|.$$

It means that  $\varepsilon > 1$ , i.e., it contradicts the hypothesis. Therefore,  $c_{i,1} < 0$  and  $\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) < 0$ .  $\square$

Consider a polynomial  $f(x) = px^3 + 2px^2 + (9p+1)x - 1$  for  $p > 0$ . Since  $f(0) = -1$  and  $f(1) = 12p$ ,  $f$  has a root in the interval  $(0, 1)$ . From that  $f'(x) = 3px^2 + 4px + 9p + 1 > 0$  for all  $x > 0$ ,  $f$  is monotone increasing in  $(0, 1)$ , i.e., the root  $t$  of  $f$  in  $(0, 1)$  is unique. Hence, for  $x \in (0, t)$ , it holds that

$$(4.3) \quad \frac{9x + 2x^2 + x^3}{1 - x} < \frac{1}{p}.$$

The previous inequality is useful to prove the following theorem.

**Theorem 4.2.** *Let  $Y_{i+1} = X_i - 2P'^{-1}_{X_i}(P(X_i))$ ,  $p = \|\mathbf{P}_{\mathcal{N}}\|$ , and let  $\varepsilon \in (0, t)$  be given where  $t$  is the real root of  $f(x) = px^3 + 2px^2 + (9p+1)x - 1$  in  $(0, 1)$ . Suppose that  $i \geq i_0$  implies that*

$$(4.4) \quad \left| \frac{\|\tilde{X}_i\|}{\|\tilde{X}_{i+1}\|} - 2 \right| < \varepsilon, \quad \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| < \varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|$$

for some  $i_0 \in \mathbb{N}_0$ . Then, for  $i \geq i_0$ ,

$$(4.5) \quad \|Y_{i+1} - S\| < \|\tilde{X}_{i+1}\|.$$

*Proof.* From the definition of  $Y_{i+1}$ , we get that

$$\begin{aligned} Y_{i+1} - S &= X_i - 2P'^{-1}_{X_i}(P(X_i)) - S \\ &= 2(X_i - P'^{-1}_{X_i}(P(X_i))) - X_i - S \\ &= 2X_{i+1} - X_i - 2S + S \\ &= 2\tilde{X}_{i+1} - \tilde{X}_i. \end{aligned}$$

So, we will show that  $\|2\tilde{X}_{i+1} - \tilde{X}_i\| < \|\tilde{X}_{i+1}\|$ . From the hypothesis, we obtain that

$$(4.6) \quad (1 - \varepsilon)\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\| < \|\tilde{X}_i\| < (1 + \varepsilon)\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|.$$



It yields two following inequalities,

$$(4.7) \quad \begin{aligned} 2 - \varepsilon &< \frac{\|\tilde{X}_i\|}{\|\tilde{X}_{i+1}\|} < \frac{(1 + \varepsilon)\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|}{(1 - \varepsilon)\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\|}, \\ \frac{(1 - \varepsilon)\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|}{(1 + \varepsilon)\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\|} &< \frac{\|\tilde{X}_i\|}{\|\tilde{X}_{i+1}\|} < 2 + \varepsilon. \end{aligned}$$

From the previous two inequalities, we get the inequality

$$(4.8) \quad \frac{(2 - \varepsilon)(1 - \varepsilon)}{1 + \varepsilon} < \frac{\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|}{\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\|} = \frac{c_{i,1}}{c_{i+1,1}} < \frac{(2 + \varepsilon)(1 + \varepsilon)}{1 - \varepsilon},$$

where  $c_{i,1}$  and  $c_{i+1,1}$  are scalars which are in the proof of Lemma 4.1, i.e., they are negative. It is obtained that

$$(4.9) \quad \frac{-5\varepsilon + \varepsilon^2}{1 + \varepsilon}(-c_{i+1,1}) < 2c_{i+1,1} - c_{i,1} < \frac{5\varepsilon + \varepsilon^2}{1 - \varepsilon}(-c_{i+1,1}).$$

Since  $\frac{5\varepsilon + \varepsilon^2}{1 - \varepsilon} - \frac{5\varepsilon - \varepsilon^2}{1 + \varepsilon}$  is positive for  $0 < \varepsilon < 1$ ,

$$(4.10) \quad |2c_{i+1,1} - c_{i,1}| < \frac{5\varepsilon + \varepsilon^2}{1 - \varepsilon}|c_{i+1,1}|.$$

Therefore, we get the following inequality,

$$(4.11) \quad \begin{aligned} \|2\tilde{X}_{i+1} - \tilde{X}_i\| &= \|2\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}) - \mathbf{P}_{\mathcal{N}}(\tilde{X}_i) + 2\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i+1}) - \mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| \\ &\leq \|2\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}) - \mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\| + \|2\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i+1})\| + \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| \\ &< \|2\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}) - \mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\| + 2\varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\| + \varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\| \\ &= |2c_{i+1,1} - c_{i,1}|\|V_1\| + 2\varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\| + \varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\| \\ &< \frac{5\varepsilon + \varepsilon^2}{1 - \varepsilon}|c_{i+1,1}|\|V_1\| + 2\varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\| + \frac{(2 + \varepsilon)(1 + \varepsilon)}{1 - \varepsilon}\varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\| \\ &= \frac{5\varepsilon + \varepsilon^2}{1 - \varepsilon}\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\| + 2\varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\| + \frac{(2 + \varepsilon)(1 + \varepsilon)}{1 - \varepsilon}\varepsilon\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\| \\ &= \frac{9\varepsilon + 2\varepsilon^2 + \varepsilon^3}{1 - \varepsilon}\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1})\| \\ &\leq \frac{9\varepsilon + 2\varepsilon^2 + \varepsilon^3}{1 - \varepsilon}p\|\tilde{X}_{i+1}\|. \end{aligned}$$

Since  $\varepsilon \in (0, t)$ ,  $\frac{9\varepsilon + 2\varepsilon^2 + \varepsilon^3}{1 - \varepsilon} < \frac{1}{p}$  holds,

$$(4.12) \quad \|Y_{i+1} - S\| = \|2\tilde{X}_{i+1} - \tilde{X}_i\| < \|\tilde{X}_{i+1}\|.$$

□

The theoretical result in Theorem 4.2 suggests a modified Newton method, as in [10] for algebraic Riccati equations. The main ideas of the algorithm are that we choose  $X_{i+1}$  as the next step of  $X_i$  if  $\|P(Y_{i+1})\| \geq \eta$  for given tolerance  $\eta$  and the iteration is terminated if  $\|P(X_{i+1})\| < \eta$  or  $\|P(Y_{i+1})\| < \eta$ .

**Algorithm 4.3.** The Modified Newton Method for the Matrix Polynomial Equations for the given tolerance  $\eta$ .

1.  $X_0 \leftarrow 0$ ;
2. Calculate  $H$  such that  $\mathcal{P}'_{X_0} \text{vec}(H) = -\text{vec}(P(X_0))$ ;
3.  $X_0 \leftarrow X_0 + 2H$ ;
4. If  $\|P(X_0)\| < \eta$ , then go to step 7;

5.  $X_0 \leftarrow X_0 - H$ ;
6. If  $\|P(X_0)\| \geq \eta$ , then go to step 2;
7.  $S \leftarrow X_0$ .

## 5. NUMERICAL EXPERIMENTS

In this section, we compare the effectiveness of the modified Newton method and the pure one for the MPE. The experiments are made with MATLAB R2016a. The tolerance of the algorithm of Newton's method is given by  $\eta = m \times 10^{-16}$  and we will stop the iteration if  $\|P(X_{i+1})\| < \eta$ .

**Example 5.1.** This example is given to check Theorem 4.2, theoretically. So, we give the example whose solution is easy to be found, and the calculations of the iterations in the experiments are computed up to 100 digits with `vpa` function. Let an MPE (1.1) with degree  $n = 6$  be given with the following coefficients

$$(5.1) \quad \begin{cases} A_k = a_k W, & \text{for } k = 0, 2, 3, 4, 5, \\ A_1 = a_1 W - I_m, \\ A_6 = W, \end{cases}$$

where

$$(5.2) \quad W = \frac{1}{6200(m-1)}(\mathbf{1}_{m \times m} - I_m), \quad \text{and} \quad \begin{cases} a_0 = 4096, & a_1 = 56, \\ a_2 = 384, & a_3 = 1312, \\ a_4 = 321, & a_5 = 30. \end{cases}$$

Then,  $A_k$ 's satisfy Assumption 1.1 and (2.1). Hence, the MPE has the minimal positive solution  $S$ .

Let  $m = 3$ . Then, the minimal nonnegative solution

$$S = \begin{bmatrix} \frac{2r+1}{3} & \frac{1-r}{3} & \frac{1-r}{3} \\ \frac{1-r}{3} & \frac{2r+1}{3} & \frac{1-r}{3} \\ \frac{1-r}{3} & \frac{1-r}{3} & \frac{2r+1}{3} \end{bmatrix},$$

where  $r \approx -0.3287191$  which is the nearest real root to 0 of the equation  $x^6 + 30x^5 + 321x^4 + 1312x^3 + 384x^2 + 12456x + 4096 = 0$ . Furthermore,  $-\mathcal{P}'_S$  is a singular  $M$ -matrix. Therefore, we can calculate  $\mathbf{P}_{\mathcal{N}}$  and  $\mathbf{P}_{\mathcal{M}}$ , easily. In this example, in fact,  $-\mathcal{P}'_S$  is symmetric, so,  $\mathbf{P}_{\mathcal{N}}$  and  $\mathbf{P}_{\mathcal{M}}$  are orthogonal projections, i.e.,  $p = \|\mathbf{P}_{\mathcal{N}}\| = 1$ . Thus,  $t \approx 0.097985683$  is the real root of  $x^3 + 2x^2 + 10x - 1$ .

If we calculate the Newton iteration  $\{X_i\}_{i=1}^{\infty}$  in (2.4) with  $X_0 = 0$  and  $\{Y_i\}_{i=1}^{\infty}$  in Theorem 4.2, then we obtain Figures ??, ??, ?? as results of  $\left| \frac{\|\tilde{X}_i\|}{\|\tilde{X}_{i+1}\|} - 2 \right|$ ,  $\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|$ ,  $\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\|$ ,  $\|X_i - S\|$ , and  $\|Y_i - S\|$ . Figure ?? shows that  $\left| \frac{\|\tilde{X}_i\|}{\|\tilde{X}_{i+1}\|} - 2 \right| < t$  if  $i \geq 2$ . From Figure ??, we get that  $\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| < t\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|$  when  $i \geq 1$ . Finally, we see that  $\|Y_i - S\| < \|X_i - S\|$  for  $i \geq 2$  through Figure ??.

This experiment shows that, for given  $i$ ,  $Y_i$  is closer to the solution than  $X_i$  if it satisfies the conditions of Theorem 4.2. Furthermore, we obtain that  $Y_i$ 's for  $i \geq 13$  are closer to the solution than any  $X_i$ 's. It means that we do not need to compute  $X_i$  for  $i \geq 13$  since  $Y_{13}$  is thought to be a "numerical" minimal nonnegative solution which is close sufficiently to the "mathematical" minimal nonnegative solution.

**Example 5.2.** (cf. [13, Example 1]) In this example, we use Algorithm 4.3 for the following MPEs and  $m = 8$ . Let MPEs

$$(5.3) \quad Q(X) = W_0 + (W_1 - I_m)X + W_2X^2 = 0,$$

$$(5.4) \quad R(X) = W_3 + (W_4 - I_m)X + W_5X^2 = 0$$

be given, where  $W_k$  is a random nonnegative matrix which has null diagonal entries and positive off-diagonal entries such that  $W_k \mathbf{1}_m = \frac{1}{3} \mathbf{1}_m$  for  $k = 0, 1, 2$ ,  $W_3 \mathbf{1}_m = \frac{1}{2} \mathbf{1}_m$ , and  $W_k \mathbf{1}_m = \frac{1}{4} \mathbf{1}_m$  for  $k = 4, 5$ . Since  $I - \sum_{k=0}^2 W_k$  and  $I - \sum_{k=3}^5 W_k$  are singular irreducible  $M$ -matrices and the coefficients satisfy Assumption 1.1, (5.3) and (5.4) have the minimal positive solutions.

We run the pure Newton algorithm(PNA) and the modified Newton algorithm(MNA) 300 times with  $X_0 = 0$ , for (5.3) and (5.4) respectively. All of the 300 experiments are run with different coefficients. Averages of the iteration numbers are given in Table 5.1.

	PNA	MNA
$Q(X) = 0$	26	10.99
$R(X) = 0$	7	7

TABLE 5.1. Averages of the iteration numbers

For (5.3), MNA finds the solution faster than PNA. But, for (5.4), the iteration numbers of MNA and PNA are same. Figure ?? shows that the solution  $S_Q$  of (5.3) is numerically non-simple and the solution  $S_R$  of (5.4) is simple. We can see from Table 5.1 that MNA is significantly more efficient than PNA for finding non-simple solutions.

## REFERENCES

- [1] Attahiru Sule Alfa. Combined elapsed time and matrix-analytic method for the discrete time  $GI/G/1$  and  $GIX/G/1$  systems. *Queueing Syst.*, 45:5–25, 2003.
- [2] Nigel G. Bean, Leslie W. Bright, Guy Latouche, Charles E. M. Pearce, Philip K. Pollett, and Peter G. Taylor. The quasi-stationary behavior of quasi-birth-and-death processes. *Ann. Appl. Probab.*, 7(1):134–155, Feb 1997.
- [3] Dario A. Bini, Guy Latouche, and Beatrice Meini. *Numerical Methods for Structured Markov Chains*. Oxford University Press Oxford, 2005.
- [4] Geoffrey. J. Butler, Charles R. Johnson, and Henry Wolkowicz. Nonnegative solutions of a quadratic matrix equation arising from comparison theorems in ordinary differential equations. *SIAM J. Algebraic Discrete Methods*, 6(1):47–53, January 1985.
- [5] George J. Davis. Numerical solution of a quadratic matrix equation. *SIAM J. Sci. Comput.*, 2:164–175, 1981.
- [6] George J. Davis. Algorithm 598: an algorithm to compute solvent of the matrix equation  $AX^2 + BX + C = 0$ . *ACM Trans. Math. Software*, 9(2):246–254, June 1983.
- [7] Israel Gohberg, Peter Lancaster, and Leiba Rodman. *Matrix Polynomials*. Academic Press, 1982.
- [8] Chun-Hua Guo. Nonsymmetric algebraic Riccati equations and Wiener-Hopf factorization for  $M$ -matrices. *SIAM J. Matrix Anal. Appl.*, 23(1):225–242, 2001.
- [9] Chun-Hua Guo and Nicholas J. Higham. Iterative solution of a nonsymmetric algebraic Riccati equation. *SIAM J. Matrix Anal. Appl.*, 29:396–412, 2007.
- [10] Chun-hua Guo and Peter Lancaster. Analysis and modification of Newton’s method for algebraic Riccati equations. *Math. Comp.*, 67(223):1089–1105, 1998.
- [11] Chun-Hua Guo and Alan J Laub. On the iterative solution of a class of nonsymmetric algebraic Riccati equations. *SIAM J. Matrix Anal. Appl.*, 22(2):376–391, 2000.
- [12] Sophie Hautphenne, Guy Latouche, and Marie-Ange Remiche. Newton’s iteration for the extinction probability of a Markovian binary tree. *Linear Algebra Appl.*, 428:2791–2804, 2008.

- [13] Charlie He, Beatrice Meini, Noah H. Rhee, and Khosrow Sohraby. A quadratically convergent Bernoulli-like algorithm for solving matrix polynomial equations in markov chains. *Electronic transactions on numerical analysis*, 17:151–167, 2004.
- [14] Qi-Ming He and Marcel F. Neuts. On the convergence and limits of certain matrix sequences arising in quasi-birth-and-death Markov chains. *J. Appl. Probab.*, 38(2):519–541, 2001.
- [15] Nicholas J. Higham and Hyun-Min Kim. Numerical analysis of a quadratic matrix equation. *IMA J. Numer. Anal.*, 20(4):499–519, 2000.
- [16] Nicholas J. Higham and Hyun-Min Kim. Solving a quadratic matrix equation by Newton’s method with exact line searches. *SIAM J. Matrix Anal. Appl.*, 23:303–316, 2001.
- [17] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, 1985.
- [18] Roger A. Horn and Charles R. Johnson. *Topics in Matrix Analysis*. Cambridge University Press, 2nd edition, 1995.
- [19] C. T. Kelley. A shamanskii-like acceleration scheme for nonlinear equations at singular roots. *Math. Comp.*, 47(176):609–623, 1986.
- [20] Hyun-Min Kim. Convergence of Newton’s method for solving a class of quadratic matrix equations. *Honam Math. J.*, 30(2):399–409, 2008.
- [21] Peter Lancaster. *Lambda-matrices and Vibrating Systems*. Pergamon Press, 1966.
- [22] Peter Lancaster and Miron Tismenetsky. *The Theory of Matrices with Applications*. Academic Press, 2 edition, 1985.
- [23] Guy Latouche and Vaidyanathan Ramaswami. *Introduction to Matrix Analytic Methods in Stochastic Modeling*. ASA-SIAM, 1999.
- [24] Jie Meng, Sang-hyup Seo, and Hyun-min Kim. Condition numbers and backward error of a matrix polynomial equation arising in stochastic models. *J. Sci. Comput.*, January 2018. First Online : <https://doi.org/10.1007/s10915-018-0641-x>.
- [25] George Poole and Thomas Boullion. A survey on  $M$ -matrices. *SIAM Rev.*, 16(4):419–427, 1974.
- [26] Jong-Hyeon Seo and Hyun-Min Kim. Convergence of pure and relaxed Newton methods for solving a matrix polynomial equation arising in stochastic models. *Linear Algebra Appl.*, 440:34–49, 2014.
- [27] Sang-Hyup Seo, Jong-Hyun Seo, and Hyun-Min Kim. Newton’s method for solving a quadratic matrix equation with special coefficient matrices. *Honam Math. J.*, 35(3):417–433, 2013.
- [28] Gilbert Strang. *Linear Algebra and Its Applications*. Thomson, Brooks/Cole, 4th edition, 2006.

SANG-HYUP SEO, DEPARTMENT OF MATHEMATICS, PUSAN NATIONAL UNIVERSITY, BUSAN, 46241, REPUBLIC OF KOREA

Email address: [saibie1677@gmail.com](mailto:saibie1677@gmail.com)

JONG-HYEON SEO, CHUBU UNIVERSITY ACADEMY OF EMERGING SCIENCE, KASUGAI, 487-0027, JAPAN

Email address: [hyeonni94@gmail.com](mailto:hyeonni94@gmail.com)

HYUN-MIN KIM, DEPARTMENT OF MATHEMATICS, PUSAN NATIONAL UNIVERSITY, BUSAN, 46241, REPUBLIC OF KOREA

Email address: [hyunmin@pusan.ac.kr](mailto:hyunmin@pusan.ac.kr)