

# SOLVING NEARLY NON-SIMPLE MATRIX POLYNOMIAL EQUATIONS BY NEWTON'S METHOD WITH ADVANCED LINE SEARCHES

SANG-HYUP SEO

ABSTRACT.

## 1. INTRODUCTION

We consider a matrix polynomial equation(MPE) with  $n$ -degree defined by

$$(1.1) \quad P(X) = \sum_{k=0}^n A_k X^k = A_0 + A_1 X + \cdots + A_{n-1} X^{n-1} + A_n X^n = 0,$$

where the coefficient matrices  $A_k$ 's are  $m \times m$  matrices. Then, the unknown matrix  $X$  must be an  $m \times m$  matrix. A matrix  $S$  is called a solution of (1.1) if  $P(S) = 0$ .

The MPE (1.1) often occurs in the theory of differential equations, system theory, network theory, stochastic theory, quasi-birth-and-death and other areas [1–4, 9, 13, 21–23]. Specially, in quasi-birth-and-death and stochastic problems, finding the minimal nonnegative solution of a matrix equation is an important issue.

There are many researches to find the minimal nonnegative solution. Guo and Laub [12] considered a nonsymmetric algebraic Riccati equation, and they proposed iteration algorithms which converge to the minimal positive solution. In [10], Guo provided a sufficient condition for the existence of nonnegative solutions of nonsymmetric algebraic Riccati equations. Kim [20] showed that the minimal positive solutions also can be found by the Newton method with the zero initial matrices in some different types of quadratic equations. Seo and Kim [27, 29] studied the Newton iteration for matrix polynomial equations.

Newton's method is one of powerful tools to find solutions of nonlinear matrix equations. By Kantorovich theorem [16], the convergence rate of the method is quadratic if the derivative on the domain is Lipschitz continuous and at the solution is nonsingular. But, if the derivative at the solution is singular, then we cannot apply Kantorovich theorem, i.e., we cannot guarantee that the rate is quadratic. The followings are researches to analyze the problems with singular derivative at the solution and improve the method.

For general functions on Banach spaces, Reddien [25], Decker and Kelley [6, 7] gave analyses about Newton's method for singular problems. In [5], Decker, Keller, and Kelley provided an acceleration of Newton's method for singular problems and analyzed for the convergence rate of the method. Kelley and Suresh [19] suggested a new accelerated Newton's method at singular points. Decker and Kelley [8] showed an analysis for Newton's method at nearly singular roots. In [18], Kelley analyzed convergence rate of the method for functions whose high order derivatives at the

---

*Date:* July 21, 2022.

*2010 Mathematics Subject Classification.* 65H10.

*Key words and phrases.* matrix polynomial equation, positive solution, nonnegative solution,  $M$ -matrix, Newton's method, line search, acceleration of a method, nearly non-simple.

solution are singular. For specific functions on  $\mathbb{R}^{m \times n}$ , Guo and Lancaster [11] analyzed and provided a modification about Newton's method for algebraic Riccati equations at singular roots. In [28], S.-H. Seo, J.-H. Seo, and H.-M. Kim suggested a modified Newton method for matrix polynomial equations with  $n$ -degree for singular problems.

In [5, 19], accelerations for Newton's method at singular roots was suggested for general functions on Banach spaces. For specific functions on  $\mathbb{R}^{m \times n}$ , modifications of Newton's method at singular roots was provided in [11, 28]. But, the main hypotheses of the papers are similar and that the solution is non-simple, i.e., the derivative at the solution is singular. It means that the accelerations for Newton's method cannot be applied and the accelerated iterations cannot be guaranteed converge to the solutions if the solution is simple. Even if the iterations converge to the solutions, they cannot be guaranteed better than the pure iterations.

Otherwise, in [8], authors analyzed and suggested an acceleration for Newton's method about the case of nearly singular roots with the general bifurcation problem [17, 30]. In [27], it was suggested that a global accelerated Newton's method, which can be applied whether the solution is simple or not, for (1.1).

The aim of this paper is also to provide an improved Newton's method for (1.1) which satisfies following Assumption 1.1 at nearly singular roots. We show that the differences between the Newton sequences and the solution lie on nearly one-dimensional space. Moreover, with the exact line search idea in [26], we suggest a proper acceleration weight  $\alpha$ .

**Assumption 1.1.** For the MPE (1.1),

- 1) The coefficient matrices  $A_k$ 's are nonnegative except  $A_1$ .
- 2)  $-A_1$  is a nonsingular  $M$ -matrix.
- 3)  $A_0$ ,  $A_1$ , and  $\sum_{k=2}^n A_k$  are irreducible.

Here, we give some basic definitions and lemmas for this paper.

Let  $A, B \in \mathbb{R}^{m \times m}$  be matrices. If all elements of  $A$  is nonnegative, then we call that  $A$  is a *nonnegative* matrix and denote  $A \geq 0$ . In similar sense, we define  $A \leq 0$ ,  $A > 0$ , and  $A < 0$ . If a matrix  $A$  can be written as  $rI_m - B$  with  $B \geq 0$  and  $r \in \mathbb{R}$ , we call that  $A$  is a *Z-matrix*. Moreover, if  $A = rI_m - B$  is a *Z-matrix* and  $r \geq \rho(B)$  then  $A$  is called an *M-matrix*. The following is a basic theorem for *M-matrices*.

**Theorem 1.2.** For a *Z-matrix*  $A$ , the following are equivalent:

- (1)  $A$  is a nonsingular *M-matrix*.
- (2)  $A^{-1}$  is nonnegative.
- (3)  $Av > 0$  for some vector  $v > 0$ .
- (4) All eigenvalues of  $A$  have positive real parts.

Let a function  $F : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^{m \times n}$  and an equation  $F(X) = 0$  be given. If nonnegative solutions  $S_1$  and  $S_2$  of  $F(X) = 0$  are satisfy

$$(1.2) \quad S_1 \leq S \leq S_2,$$

for any nonnegative solution  $S$  of  $F(X) = 0$ , then they are called the *minimal nonnegative solution* and the *maximal nonnegative solution*, respectively. The *minimal positive solution* and the *maximal positive solution* also are can be defined, similarly. If the Fréchet derivative of  $F$  at a solution  $S$  is nonsingular, then  $S$  is called *simple*. Furthermore, we call that a not simple solution is *non-simple*. In this paper, for convenience, the notation  $\|\cdot\|$  is used instead of the Frobenius norm  $\|\cdot\|_F$  and  $\tilde{X}$  denotes  $S - X$ , the difference between  $X$  and the solution  $S$ , because they are used very frequently.

## 2. CONVERGENCE OF NEWTON'S METHOD FOR THE MPE

In this paper, we denote that

$$(2.1) \quad P'[X](H) = \sum_{k=1}^n \sum_{l=0}^{k-1} A_k X^l H X^{k-l-1}$$

is the Fréchet derivative of the MPE (1.1) at  $X$  in the direction  $H$ , and

$$(2.2) \quad P''[X](K, H) = \sum_{k=2}^n \sum_{l=0}^{k-2} \sum_{j=0}^l A_k (X^l H X^j K X^{n-l-j-2} + X^l K X^j H X^{n-l-j-2})$$

is the second Fréchet derivative at  $X$ .

For given initial guess  $X_0$ , the Newton iteration of (1.1),

$$(2.3) \quad X_{i+1} = X_i - P'[X_i]^{-1}(P(X_i)), \quad i = 0, 1, 2, \dots$$

can be separated into two parts as

$$(2.4) \quad \begin{cases} P'[X_i](H_i) = -P(X_i), \\ X_{i+1} = X_i + H_i, \end{cases} \quad i = 0, 1, 2, \dots$$

If we define that

$$(2.5) \quad \mathcal{P}'[X] = \text{vec} \circ P[X]' \circ \text{vec}^{-1} = \sum_{k=1}^n \sum_{l=0}^{k-1} (X^{k-l-1})^T \otimes A_k X^l,$$

then solving the above equation of (2.4) is equivalent to solving the following  $m^2 \times m^2$  linear system

$$(2.6) \quad \mathcal{P}'[X_i] \text{vec}(H_i) = \text{vec}(-P(X_i)),$$

where  $\text{vec}$  is the vectorization by [15, Lemma 4.3.1].

We introduce a sufficient condition of the existence of the minimal nonnegative solution of the MPE (1.1) with Assumption 1.1, and give some analysis for Newton's method.

**Theorem 2.1.** [24, Theorem 2.1] *Let the MPE (1.1) with 1) and 2) in Assumption 1.1 be given. Then, there exists the minimal nonnegative solution if*

$$(2.7) \quad -\sum_{k=0}^n A_k \text{ is a nonsingular or singular irreducible } M\text{-matrix.}$$

**Theorem 2.2.** [28, Theorem 2.2] *Suppose that the MPE (1.1) satisfies Assumption 1.1 and (2.7). Then, the Newton sequence  $\{X_i\}$  with  $X_0 = 0$  is well defined, is monotone nondecreasing, and converges to the minimal positive solution  $S$ . Furthermore,  $-\mathcal{P}'[X_i]$  is a nonsingular irreducible  $M$ -matrix for  $i \in \mathbb{N}$ , and  $-\mathcal{P}'[S]$  is an irreducible  $M$ -matrix.*

3. THE NEARLY SINGULAR DERIVATIVE  $-\mathcal{P}'[S]$ 

Generally, by Kantorovich theorem [16], the Newton iteration (2.3) converges quadratically if the derivative at the solution is nonsingular. But, if the derivative becomes nearly singular, the quadratic convergence radius will be very small. In this section, we will see the converging tendency of (2.3) when the derivative is nearly singular.

Let the MPE (1.1) satisfy Assumption 1.1, and let  $S$  be the minimal positive solution of (1.1). Then,  $-\mathcal{P}'[S]$  is an irreducible  $M$ -matrix. Therefore,  $\mu$  is a non-negative and simple eigenvalue of  $-\mathcal{P}'[S]$  if

$$\mu = \min\{|\lambda| : \lambda \text{ is an eigenvalue of } -\mathcal{P}'[S]\}$$

by Perron-Frobenius Theorem in [14].

Here, we think over an other MPE,

$$(3.1) \quad P_\varepsilon(X) = (A_0 - (\mu - \varepsilon)S) + (A_1 + (\mu - \varepsilon)I_m)X + \sum_{k=2}^n A_k X^k = 0,$$

for  $\varepsilon \geq 0$ . Clearly,  $S$  is a solution of (3.1). Since the representation of the Fréchet derivative of  $P_\varepsilon$  at  $S$  is that

$$\begin{aligned} \mathcal{P}'_\varepsilon[S] &= I_m \otimes (A_1 + (\mu - \varepsilon)I_m) + \sum_{k=2}^n \sum_{l=0}^{k-1} (S^{k-l-1})^T \otimes A_k S^l \\ &= \mathcal{P}'[S] + (\mu - \varepsilon)I_{m^2}, \end{aligned}$$

$-\mathcal{P}'_\varepsilon[S]$  is an  $M$ -matrix by Theorem 1.2, and the nonnegative number  $\varepsilon$  is the smallest simple eigenvalue of  $-\mathcal{P}'_\varepsilon[S]$ . Here, we assume that

$$(3.2) \quad N = \text{vec}^{-1}(\mathbf{n}), \quad N = \text{span}(N), \quad \mathbb{R}^{m \times m} = \mathcal{N} \oplus \mathcal{M},$$

where  $\mathbf{n}$  is a unit eigenvector of  $-\mathcal{P}'[S]$  corresponding to  $\mu$ . In addition, we define  $\mathbf{P}_\mathcal{N}$  is a projection onto  $\mathcal{N}$  parallel to  $\mathcal{M}$ , and  $\mathbf{P}_\mathcal{M} = I - \mathbf{P}_\mathcal{N}$ .

**Lemma 3.1.** *Suppose the MPE (1.1) satisfies Assumption 1.1 and (3.1) is given. Then,  $\varepsilon$  is a simple eigenvalue of  $-\mathcal{P}'_\varepsilon[S]$ ,  $\mathbf{P}_\mathcal{N} P''_\varepsilon[S](N, N) \neq 0$ , and  $P'_\varepsilon[S](\mathcal{M}) = \mathcal{M}$ .*

*Proof.* Since  $S$  is positive and  $A_k$ 's are irreducible,  $-\mathcal{P}'[S]$  is irreducible. Thus,  $-\mathcal{P}'_\varepsilon[S] = -\mathcal{P}'[S] + (\varepsilon - \mu)I_m$  is also irreducible as the diagonal entries do not effect on irreducibility of matrices. Then, by Perron-Frobenius Theorem [14],  $\varepsilon$  is a simple eigenvalue of  $-\mathcal{P}'_\varepsilon[S]$  with a positive eigenvector. Thus, we can find  $m^2$  linearly independent vectors  $\chi_1, \chi_2, \dots, \chi_{m^2}$  such that  $\chi_1 > 0$  and

$$(3.3) \quad \mathfrak{X}^{-1}(-\mathcal{P}'_\varepsilon[S])\mathfrak{X} = \begin{bmatrix} \varepsilon & 0 \\ 0 & \mathfrak{D} \end{bmatrix}, \quad \text{where } \mathfrak{X} = \begin{bmatrix} \chi_1 & \chi_2 & \cdots & \chi_{m^2} \end{bmatrix}$$

and  $\mathfrak{D}$  is an  $(m^2-1) \times (m^2-1)$  nonsingular matrix, i.e.,  $\mathcal{M} = \text{span}(\text{vec}^{-1}(\chi_2), \dots, \text{vec}^{-1}(\chi_{m^2}))$  and  $P'_\varepsilon[S](\mathcal{M}) = \mathcal{M}$ . By the same way, we also have a positive vector  $\psi_1$  such that  $-\psi_1^T \mathcal{P}'_\varepsilon[S] = \varepsilon \psi_1^T$ . Since  $N \in \mathcal{N} = \text{span}(\text{vec}^{-1}(\chi_1))$ , there exists nonzero  $a \in \mathbb{R}$  such that  $N = a \text{vec}^{-1}(\chi_1)$ . From  $\text{vec}^{-1}(\chi_1) > 0$  and  $S > 0$ , we have

$$\begin{aligned} (3.4) \quad P''_\varepsilon[S](N, N) &= P''[S](N, N) \\ &= \sum_{k=2}^n \sum_{l=0}^{k-2} \sum_{j=0}^l A_k (S^l N S^j N S^{n-l-j-2} + S^l N S^j N S^{m-l-j-2}) \\ &= 2a^2 \sum_{k=2}^n \sum_{l=0}^{k-2} \sum_{j=0}^l A_k (S^l \text{vec}^{-1}(\chi_1) S^j \text{vec}^{-1}(\chi_1) S^{n-l-j-2}) > 0. \end{aligned}$$

On the other hand,

$$(3.5) \quad \text{vec}(P''[S](N, N)) = k_1 \chi_1 + k_2 \chi_2 + \cdots + k_{m^2} \chi_{m^2}$$

for some real numbers  $k_1, k_2, \dots, k_{m^2}$ . By Fundamental theorem of linear algebra in [31] and Lemma 6.3.10 in [14], we have

$$(3.6) \quad \psi_1^T \text{vec}(P''[S](N, N)) = k_1 \psi_1^T \chi_1.$$

Since all of  $P''[S](N, N)$ ,  $\chi_1$ , and  $\psi_1$  are positive, the left side of (3.6) is positive, i.e.,  $k_1$  must be positive. Therefore,

$$(3.7) \quad \mathbf{P}_\mathcal{N}(P''[S](N, N)) = k_1 \text{vec}^{-1}(\chi_1) > 0.$$

□

**Lemma 3.2.** [28, Lemma 3.3] *Let  $S$  be the minimal positive solution of (1.1) with Assumption 1.1, and let a Newton sequence  $\{X_{\varepsilon,i}\}_{i=0}^{\infty}$  for (3.1) converge to  $S$ . Then,*

$$(3.8) \quad \|P_{\varepsilon}(X_{\varepsilon,i})\| \leq a\|\tilde{X}_{\varepsilon,i}\|^2 + b\|\tilde{X}_{\varepsilon,i}\|\|\tilde{X}_{\varepsilon,i-1}\| + c\|\tilde{X}_{\varepsilon,i-1}\|^2$$

for some positive real numbers  $a, b, c$ .

The equation (3.1) is not same as (1.1) but similar. So, Lemma 3.2 can be proved like [28, Lemma 3.3] because it is proved with just Taylor's Theorem, without the conditions of coefficient matrices.

**Lemma 3.3.** *For any fixed  $\theta > 0$ , let*

$$(3.9) \quad \Omega = \{i \in \mathbb{N} \cup \{0\} \mid \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i})\| > \theta\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{\varepsilon,i})\|\}$$

where  $\{X_{\varepsilon,i}\}$  is a Newton sequence for (3.1) which converges to  $S$ , and let  $c_1 > 0$  be the smallest singular value of  $-P'_{\varepsilon}[S]|_{\mathcal{M}}$ . Then, for  $\varepsilon < c_1\theta$ , there exist an integer  $i_0$  and a constant  $c > 0$  such that  $\|\tilde{X}_{\varepsilon,i}\| \leq c\|\tilde{X}_{\varepsilon,i-1}\|^2$  for all  $i \geq i_0$  in  $\Omega$ .

*Proof.* Using Taylor's Theorem and the fact that  $-P'_{\varepsilon}[S](\mathbf{P}_{\mathcal{N}}(\tilde{X}_{\varepsilon,i})) = \varepsilon\mathbf{P}_{\mathcal{N}}(\tilde{X}_{\varepsilon,i})$ , we obtain that

$$(3.10) \quad \begin{aligned} P_{\varepsilon}(X_{\varepsilon,i}) &= P_{\varepsilon}(S) - P'_{\varepsilon}[S](\tilde{X}_{\varepsilon,i}) + O(\|\tilde{X}_{\varepsilon,i}\|^2) \\ &= -P'_{\varepsilon}[S](\mathbf{P}_{\mathcal{N}}(\tilde{X}_{\varepsilon,i}) + \mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i})) + O(\|\tilde{X}_{\varepsilon,i}\|^2) \\ &= \varepsilon\mathbf{P}_{\mathcal{N}}(\tilde{X}_{\varepsilon,i}) - P'_{\varepsilon}[S](\mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i})) + O(\|\tilde{X}_{\varepsilon,i}\|^2). \end{aligned}$$

Since  $c_1 > 0$  is the smallest singular value of  $-P'_{\varepsilon}[S]|_{\mathcal{M}}$ ,  $\| -P'_{\varepsilon}[S](\mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i})) \| \geq c_1\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i})\|$ . For  $i \in \Omega$ , we have

$$\|\tilde{X}_{\varepsilon,i}\| \leq \|\mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i})\| + \|\mathbf{P}_{\mathcal{N}}(\tilde{X}_{\varepsilon,i})\| < (\theta^{-1} + 1)\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i})\|.$$

Thus by (3.10),

$$(3.11) \quad \begin{aligned} \|P_{\varepsilon}(X_{\varepsilon,i})\| &= \|\varepsilon\mathbf{P}_{\mathcal{N}}(\tilde{X}_{\varepsilon,i}) - P'_{\varepsilon}[S](\mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i})) + O(\|\tilde{X}_{\varepsilon,i}\|^2)\| \\ &\geq \|\varepsilon\mathbf{P}_{\mathcal{N}}(\tilde{X}_{\varepsilon,i}) - P'_{\varepsilon}[S](\mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i}))\| - c_2\|\tilde{X}_{\varepsilon,i}\|^2 \\ &\geq \left\| -P'_{\varepsilon}[S](\mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i})) \right\| - \|\varepsilon\mathbf{P}_{\mathcal{N}}(\tilde{X}_{\varepsilon,i})\| - c_2\|\tilde{X}_{\varepsilon,i}\|^2 \\ &\geq \left(c_1 - \frac{\varepsilon}{\theta}\right)\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_{\varepsilon,i})\| - c_2\|\tilde{X}_{\varepsilon,i}\|^2 \\ &\geq \left(c_1 - \frac{\varepsilon}{\theta}\right)(\theta^{-1} + 1)^{-1}\|\tilde{X}_{\varepsilon,i}\| - c_2\|\tilde{X}_{\varepsilon,i}\|^2. \end{aligned}$$

On the other hand, from Lemma 3.2, we have

$$(3.12) \quad \|P_{\varepsilon}(X_{\varepsilon,i})\| \leq c_3\|\tilde{X}_{\varepsilon,i}\|^2 + c_4\|\tilde{X}_{\varepsilon,i-1}\|\|\tilde{X}_{\varepsilon,i}\| + c_5\|\tilde{X}_{\varepsilon,i-1}\|^2.$$

With the combination of (3.11) and (3.12), we have

$$\left(c_1 - \frac{\varepsilon}{\theta}\right)(\theta^{-1} + 1)^{-1} - c_2\|\tilde{X}_{\varepsilon,i}\| \leq c_3\|\tilde{X}_{\varepsilon,i}\| + c_4\|\tilde{X}_{\varepsilon,i-1}\| + c_5\frac{\|\tilde{X}_{\varepsilon,i-1}\|^2}{\|\tilde{X}_{\varepsilon,i}\|}.$$

Since  $\tilde{X}_{\varepsilon,i}$  converges to 0, we can find an  $i_0$  such that  $\|\tilde{X}_i\| \leq c\|\tilde{X}_{i-1}\|^2$  for all  $i \geq i_0$ .  $\square$

From [8], we note that  $P_\varepsilon(X) = 0$  is the bifurcation problem where  $\varepsilon$  is a chosen continuation parameter and the solution arc  $S_\varepsilon$  is a constant arc. Specially, for  $\varepsilon = 0$ , the MPE (3.1) becomes a singular problem which can be applied by the following theorem which refer to [18, Theorem 1.1].

**Theorem 3.4.** *Let  $\mathbf{P}_N(P_0''[S](N, N))$  be nonzero and let*

$$(3.13) \quad W(\rho, \theta) = \left\{ X \mid 0 < \|\tilde{X}\| < \rho, \|\mathbf{P}_M(\tilde{X})\| \leq \theta \|\mathbf{P}_N(\tilde{X})\| \right\}.$$

*If  $X_{0,0} \in W(\rho_0, \theta_0)$  for  $\rho_0, \theta_0$  sufficiently small, then the Newton sequence  $\{X_{0,i}\}$  is well defined and  $\|P_0'[X_{0,i}]^{-1}\| \leq c\|\tilde{X}_{0,i}\|^{-1}$  for all  $i \geq 1$  and some constant  $c > 0$ . Moreover,*

$$(3.14) \quad \lim_{i \rightarrow \infty} \frac{\|\tilde{X}_{0,i+1}\|}{\|\tilde{X}_{0,i}\|} = \frac{1}{2}, \quad \lim_{i \rightarrow \infty} \frac{\|\mathbf{P}_M(\tilde{X}_{0,i})\|}{\|\mathbf{P}_N(\tilde{X}_{0,i})\|^2} = 0.$$

Defining

$$(3.15) \quad \begin{aligned} \lambda N &= \mathbf{P}_N P_\varepsilon''[S](N, N), \\ \nu N &= \mathbf{P}_N \tilde{X}, \\ s &= \text{sign}(\varepsilon \lambda), \end{aligned}$$

we consider the conical region

$$(3.16) \quad W_s(\rho, \theta) = \left\{ X \mid 0 < \|\tilde{X}\| < \rho, \|\mathbf{P}_M(\tilde{X})\| \leq \theta \|\mathbf{P}_N(\tilde{X})\|, \text{sign}(\nu) = s \right\}.$$

Then, we can apply the following theorem to the problem for  $\varepsilon > 0$ .

**Theorem 3.5.** (cf. [8, Theorem 3.22]) *Assume  $\lambda \neq 0$ , and  $\varepsilon > 0$ . Then, there are continuous functions  $\rho = \rho(\varepsilon)$ ,  $\theta = \theta(\varepsilon)$ , monotonically increasing as  $\varepsilon \downarrow 0$  such that if  $X_{\varepsilon,0} \in W_s(\rho, \theta)$  then  $P_\varepsilon'[X_{\varepsilon,0}]^{-1}$  exists and all subsequent Newton iterates remain in this set and converges to  $S$ . Further,*

$$(3.17) \quad \|\mathbf{P}_N \tilde{X}_{\varepsilon,i+1}\| < \frac{3}{4} \|\mathbf{P}_N \tilde{X}_{\varepsilon,i}\|, \quad \|\mathbf{P}_M(\tilde{X}_{\varepsilon,i+1})\| \leq K \|\tilde{X}_{\varepsilon,i}\|^2,$$

*for  $i \in \mathbb{N} \cup \{0\}$ , some  $K > 0$ , and  $\rho(\varepsilon)$ ,  $\theta(\varepsilon)$  may be chosen such that  $\rho(0) = \rho_0$ ,  $\theta(0) = \theta_0$  where  $\rho_0, \theta_0$  are values for which the conclusions of Theorem 3.4 hold.*

**Corollary 3.6.** *Assume that, for given  $\theta > 0$ ,  $\|\mathbf{P}_M(\tilde{X}_i)\| > \theta \|\mathbf{P}_N(\tilde{X}_i)\|$  for all  $i$  large enough. Then  $X_i \rightarrow S$  quadratically.*

When  $P'_S$  is singular practically the Newton sequence converges linearly, according to the corollary we conclude that the error will generally be dominated by its  $N$  component [11]. From Lemmas 3.1 and 3.3, we have the following main theorem.

**Theorem 3.7.** *If  $-\mathcal{P}'_S$  is a singular  $M$ -matrix and the convergence rate of the Newton sequence  $\{X_i\}$  in Theorem 2.2 is not quadratic, then  $\|P'_{X_i}\| \leq c\|\tilde{X}_i\|^{-1}$  for all  $i \geq 1$  and some constant  $c > 0$ . Moreover,*

$$\lim_{i \rightarrow \infty} \frac{\|\tilde{X}_{i+1}\|}{\|\tilde{X}_i\|} = \frac{1}{2}, \quad \lim_{i \rightarrow \infty} \frac{\|\mathbf{P}_M(\tilde{X}_i)\|}{\|\mathbf{P}_N(\tilde{X}_i)\|^2} = 0.$$

## 4. EXACT LINE SEARCH FOR SOME ITERATIVE POINT

- 1) Analytic하게 증명할 필요가 있는가?
- 2)  $\mathcal{N}$  안에  $\tilde{X}_i$ 가 들어있을 때,  $\tilde{X}_i$ 의 움직임에 대해 따로 증명할 필요가 있지 않을까?
- 3)  $t \in [1, 2]$ 에 대해 우리가 원하는  $S$ 가 존재함을 보여야 하지 않나?  $\leftarrow$  정확히는  $S$ 가 그 라인 근처에 놓여있다? 그걸 보여야하지 않나...

**Lemma 4.1.**  $\mathcal{N}$  안에  $\tilde{X}_i$ 가 들어있을 때,  $\tilde{X}_i$ 의 움직임에 대해 따로 증명할 필요가 있지 않을까?

*Proof.* Let  $\tilde{X}_i \in \mathcal{N}$ . Then,  $-P'[S](\tilde{X}_i) = \varepsilon \tilde{X}_i$ .

$$\begin{aligned}
 X_{i+1} &= X_i - P'[X_i]^{-1}(P(X_i)) \\
 \tilde{X}_{i+1} &= \tilde{X}_i - P'[X_i]^{-1}(P(X_i)) \\
 -P'[S](\tilde{X}_{i+1}) &= -P'[S] \left( \tilde{X}_i - P'[X_i]^{-1}(P(X_i)) \right) \\
 &= -P'[S](\tilde{X}_i) + P'[S] \left( P'[X_i]^{-1}(P(X_i)) \right) \\
 &= \varepsilon \tilde{X}_i + P'[S] \left( P'[X_i]^{-1}(P(X_i)) \right)
 \end{aligned}$$

$$\begin{aligned}
 P(X) &= P(Y) + P'[Y](X - Y) + \frac{1}{2}P''[Y](X - Y)^2 + \cdots + \frac{1}{n!}P^{(n)}[Y](X - Y)^n \\
 &= \sum_{k=0}^n \frac{1}{k!}P^{(k)}[Y](X - Y)^k
 \end{aligned}$$

$$\begin{aligned}
 P'[X](X - Y) &= P'[Y](X - Y) + P''[Y](X - Y)^2 + \cdots + \frac{1}{(n-1)!}P^{(n)}(X - Y)^n \\
 &= \sum_{k=1}^n \frac{1}{(k-1)!}P^{(k)}[Y](X - Y)^k
 \end{aligned}$$

$$\begin{aligned}
 P(X_i) &= P(S) + P'[S](-\tilde{X}_i) + \sum_{k=2}^n \frac{1}{k!}P^{(k)}[S](-\tilde{X}_i)^k \\
 &= P'[S](-\tilde{X}_i) + \sum_{k=2}^n \frac{1}{k!}P^{(k)}[S](-\tilde{X}_i)^k \\
 &= \varepsilon \tilde{X}_i + \sum_{k=2}^n \frac{1}{k!}P^{(k)}[S](-\tilde{X}_i)^k
 \end{aligned}$$

$$\begin{aligned}
 P[X_i](X_i - S) &= \sum_{k=1}^n \frac{1}{(k-1)!}P^{(k)}[S](X_i - S)^k \\
 &= P'[S](-\tilde{X}_i) + P''[S](-\tilde{X}_i)^2 + \sum_{k=3}^n \frac{1}{(k-1)!}P^{(k)}[S](-\tilde{X}_i)^k \\
 &= \varepsilon \tilde{X}_i + P''[S](-\tilde{X}_i)^2 + \sum_{k=3}^n \frac{1}{(k-1)!}P^{(k)}[S](-\tilde{X}_i)^k
 \end{aligned}$$

$$\begin{aligned}
P[X_i](X_i - S) - P(X_i) &= \sum_{k=2}^n \frac{1}{(k-1)!} P^{(k)}[S](-\tilde{X}_i)^k - \sum_{k=2}^n \frac{1}{k!} P^{(k)}[S](-\tilde{X}_i)^k \\
&= \sum_{k=2}^n \left( \frac{1}{(k-1)!} - \frac{1}{k!} \right) P^{(k)}[S](-\tilde{X}_i)^k \\
&= \sum_{k=2}^n \frac{1}{k(k-2)!} P^{(k)}[S](-\tilde{X}_i)^k
\end{aligned}$$

$$\begin{aligned}
P(X_i) &= P[X_i](X_i - X_{i+1}) = P[X_i](\tilde{X}_{i+1} - \tilde{X}_i) \\
&= P[X_i](\tilde{X}_{i+1}) - P[X_i](\tilde{X}_i)
\end{aligned}$$

□

From the previous section, we see that  $\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)$  is almost terminated more than  $\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)$  for the Newton sequence  $\{X_i\}$  which converges to  $S$  and is from a nearly singular problem. It means that  $\{\tilde{X}_i\}_{i \geq i_0}$  lies on near one-dimensional subspace  $\mathcal{N}$  for some nonnegative integer  $i_0$ . So, with just a proper scalar multiplication, we can accelerate Newton's method as

$$(4.1) \quad X_{i+1} = X_i - \alpha_i P'[X_i]^{-1}(P(X_i)), \quad i = 0, 1, 2, \dots$$

As introduced in Section 1, the proper scalar  $\alpha_i$  is suggested for specific singular problems in [11, 28]. If  $\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| < \varepsilon \|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|$  holds for small  $\varepsilon > 0$ , then  $\alpha_i = 2$  is the proper scalar. But, for nearly singular problems, we cannot guarantee that  $\alpha_i = 2$  is the proper. So, we suggest the way finding the proper scalar  $\alpha_i$  based on exact line searches in [26].

For given initial guess  $X_0$ , the Newton sequence  $\{X_i\}$  with exact line searches is defined by

$$(4.2) \quad \begin{cases} X_{i+1} = X_i + \alpha_i H_i, \\ \alpha_i = \underset{t \geq 1}{\operatorname{argmin}} \|P(X_i + tH_i)\|^2, \end{cases}$$

where  $H_i = -P'[X_i](P(X_i))$ . Thus, finding  $\alpha_i$  is equivalent to finding the minimum point for the following  $2n$ -degree polynomial

$$\begin{aligned}
(4.3) \quad p_i(t) &= \|P(X_i + tH_i)\|^2 \\
&= \operatorname{tr}((P(X_i + tH_i))^T (P(X_i + tH_i))) \\
&= \operatorname{tr} \left( \left( \sum_{k=0}^n A_k (X_i + tH_i)^k \right)^T \sum_{k=0}^n A_k (X_i + tH_i)^k \right).
\end{aligned}$$

To expand (4.3), let a function  $\tau_{(X,H)} : \{0, 1\} \rightarrow \{X, H\}$  be given by  $\tau_{(X,H)}(0) = X$  and  $\tau_{(X,H)}(1) = H$ , and let  $\mathfrak{U}_{k,l} = \{\mathbf{v} \in \{0, 1\}^k \subseteq \mathbb{R}^k \mid \mathbf{1}^T \mathbf{v} = l\}$ . In addition, for integers  $k, l$  satisfying  $0 \leq l \leq k$ , we define

$$(4.4) \quad \Phi_{(X,H)}(k, l) = \begin{cases} \sum_{\mathbf{v} \in \mathfrak{U}_{k,l}} \prod_{j=1}^k \tau_{(X,H)}(v_j) & , \text{ for } k \neq 0, \\ I_m & , \text{ for } k = 0, \end{cases}$$

where  $\mathbf{v} = [v_1 \ v_2 \ \cdots \ v_k]^T$ . Then, it is easy to verify that

$$(4.5) \quad \Phi_{(X,tH)}(k, l) = t^l \Phi_{(X,H)}(k, l), \quad \text{and} \quad (\Phi_{(X,H)}(k, l))^T = \Phi_{(X^T, H^T)}(k, l).$$



Moreover,  $(X_i + tH_i)^k = \sum_{l=0}^k \Phi_{(X_i, tH_i)}(k, l)$ , i.e.,

$$\begin{aligned} P(X_i + tH_i) &= \sum_{k=0}^n A_k (X_i + tH_i)^k \\ &= \sum_{k=0}^n A_k \sum_{l=0}^k \Phi_{(X_i, tH_i)}(k, l) \\ &= \sum_{k=0}^n \sum_{l=0}^k t^l A_k \Phi_{(X_i, H_i)}(k, l). \end{aligned}$$

Thus,

$$\begin{aligned} p_i(t) &= \text{tr} \left( \left( \sum_{r=0}^n A_r (X_i + tH_i)^r \right)^T \sum_{k=0}^n A_k (X_i + tH_i)^k \right) \\ &= \text{tr} \left( \left( \sum_{r=0}^n \sum_{s=0}^r t^s A_r \Phi_{(X_i, H_i)}(r, s) \right)^T \left( \sum_{k=0}^n \sum_{l=0}^k t^l A_k \Phi_{(X_i, H_i)}(k, l) \right) \right) \\ &= \sum_{r=0}^n \sum_{s=0}^r \sum_{k=0}^n \sum_{l=0}^k t^{s+l} \text{tr} \left( \Phi_{(X_i^T, H_i^T)}(r, s) A_r^T A_k \Phi_{(X_i, H_i)}(k, l) \right). \end{aligned}$$

In the calculation of  $\Phi_{(X_i, H_i)}(k, l)$ , there are  $\binom{k}{l}(k-1)$  matrix multiplications unless  $k = 0$ . Hence, there are about

$$\begin{aligned} &\sum_{r=1}^n \sum_{s=0}^r \sum_{k=1}^n \sum_{l=0}^k \left( \binom{r}{s}(r-1) + \binom{k}{l}(k-1) + 3 \right) \\ &= (2^{n+2}(n-2) + 8) \left( \frac{(n+1)(n+2)}{2} - 1 \right) + 3 \left( \frac{(n+1)(n+2)}{2} - 1 \right)^2 \\ &= O(2^n n^3) \end{aligned}$$

matrix multiplications to obtain all coefficients of  $p_i$  for every  $i$ .

**Theorem 4.2.** *Suppose that there exists  $i_0 \in \mathbb{N}$  such that  $\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| < \varepsilon \|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|$  for small  $\varepsilon > 0$  and  $i \geq i_0$ . Then,*

$$(4.6) \quad S \in L[X_i + H_i, X_i + 2H_i],$$

where  $L[A, B] := \{C \mid A \leq C \leq B\}$ .

**Theorem 4.3.** *Suppose that there exists  $i_0 \in \mathbb{N}$  such that  $\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| < \varepsilon \|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|$  for small  $\varepsilon > 0$  and  $i \geq i_0$ . Then, for some  $\bar{c} > 0$ ,*

$$\begin{aligned} \|P(X_i + tH_i) - P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| &< ((t-1)c_i + tc_{i+1})\bar{c}\varepsilon \\ &\leq (c_i + 2c_{i+1})\bar{c}\varepsilon, \end{aligned}$$

where  $\mathbf{P}_{\mathcal{N}}(X_i) = c_i N$ ,  $\bar{p} = \|\mathbf{P}_{\mathcal{N}}\|$ , and  $1 \leq t \leq 2$ .

*Proof.* We define  $\Psi_{(A,B)} : \{0,1\}^k \longrightarrow (\mathbb{R}^{m \times m})^k$  by  $\Psi_{(A,B)}(\mathbf{v}) = (\tau_{(A,B)}(v_i))_{i=1}^k$ . By Taylor theorem and the multilinearity of  $P^{(k)}[\cdot]$ ,

$$\begin{aligned}
P(X_i + tH_i) &= \sum_{k=1}^n \frac{1}{k!} P^{(k)}[S] \left( (t-1)\tilde{X}_i - t\tilde{X}_{i+1} \right)^k \\
&= \sum_{k=1}^n \frac{1}{k!} P^{(k)}[S] \left( \mathbf{P}_{\mathcal{N}} \left( (t-1)\tilde{X}_i - t\tilde{X}_{i+1} \right) + \mathbf{P}_{\mathcal{M}} \left( (t-1)\tilde{X}_i - t\tilde{X}_{i+1} \right) \right)^k \\
&= \sum_{k=1}^n \frac{1}{k!} \sum_{\mathbf{v} \in \{0,1\}^k} P^{(k)}[S] \Psi_{(\mathbf{P}_{\mathcal{N}}((t-1)\tilde{X}_i - t\tilde{X}_{i+1}), \mathbf{P}_{\mathcal{M}}((t-1)\tilde{X}_i - t\tilde{X}_{i+1}))}(\mathbf{v}) \\
&= \sum_{k=1}^n \frac{1}{k!} P^{(k)}[S] \left( \mathbf{P}_{\mathcal{N}} \left( (t-1)\tilde{X}_i - t\tilde{X}_{i+1} \right) \right)^k + D_i \\
&= \sum_{k=1}^n \frac{1}{k!} P^{(k)}[S] (\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S) - S)^k + D_i \\
&= P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S) - S) + D_i,
\end{aligned}$$

where

$$D_i = \sum_{k=1}^n \frac{1}{k!} \sum_{\mathbf{v} \in \{0,1\}^k \setminus \mathbf{0}} P^{(k)}[S] \Psi_{((t-1)\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}), (t-1)\mathbf{P}_{\mathcal{M}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i+1}))}(\mathbf{v}),$$

and

$$P^{(k)}[S](A)^k = P^{(k)}[S](\underbrace{A, A, \dots, A}_{k\text{-times}}).$$

Therefore,

$$\begin{aligned}
&1) \|D_i\| \\
&= \|P(X_i + tH_i) - P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| \\
&= \left\| \sum_{k=1}^n \frac{1}{k!} \sum_{\mathbf{v} \in \{0,1\}^k \setminus \mathbf{0}} P^{(k)}[S] \Psi_{((t-1)\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}), (t-1)\mathbf{P}_{\mathcal{M}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i+1}))}(\mathbf{v}) \right\| \\
&\leq \bar{c} \left\| (t-1)\mathbf{P}_{\mathcal{M}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i+1}) \right\| \\
&\leq \bar{c}(t-1) \left\| \mathbf{P}_{\mathcal{M}}(\tilde{X}_i) \right\| + \bar{c}t \left\| \mathbf{P}_{\mathcal{M}}(\tilde{X}_{i+1}) \right\| \\
&< \bar{c}\varepsilon(t-1) \left\| \mathbf{P}_{\mathcal{N}}(\tilde{X}_i) \right\| + \bar{c}\varepsilon t \left\| \mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}) \right\| \\
&= ((t-1)c_i + tc_{i+1})\bar{c}\varepsilon \\
&\leq (c_i + 2c_{i+1})\bar{c}\varepsilon.
\end{aligned}$$

$$\begin{aligned}
&\|P(X_i + tH_i)\| - \|P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| \\
&\leq \|P(X_i + tH_i) - P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| \\
&< (c_i + 2c_{i+1})\bar{c}\varepsilon
\end{aligned}$$

$$\begin{aligned}
&\|P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| - \|P(X_i + tH_i)\| \\
&\leq \|P(X_i + tH_i) - P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| \\
&< (c_i + 2c_{i+1})\bar{c}\varepsilon
\end{aligned}$$

$$\|P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| - \|P(X_i + tH_i)\| < (c_i + 2c_{i+1})\bar{c}\varepsilon$$

□

**Theorem 4.4.** *Suppose that there exists  $i_0 \in \mathbb{N}$  such that  $\|\mathbf{P}_{\mathcal{M}}(\tilde{X}_i)\| < \varepsilon \|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\|$  for small  $\varepsilon > 0$  and  $i \geq i_0$ . Then,*

$$(4.7) \quad \|P(X_i + \alpha_i H_i)\| < \|P(X_{i+1})\| \quad \text{and} \quad \|P(X_i + \alpha_i H_i)\| < \eta,$$

or

$$(4.8) \quad \|X_i + \alpha_i H_i - S\| \leq \|\tilde{X}_{i+1}\|,$$

where  $\alpha_i = \underset{1 \leq t \leq 2}{\operatorname{argmin}} \|P(X_i + tH_i)\|^2$ .

*Proof.*

$$\mathbf{P}_{\mathcal{N}}\tilde{X}_i \geq 0, \quad \mathbf{P}_{\mathcal{N}}\tilde{X}_i = c_i N$$

$$(4.9) \quad \begin{aligned} (1 - \varepsilon)\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\| &< \|\tilde{X}_i\| < (1 + \varepsilon)\|\mathbf{P}_{\mathcal{N}}(\tilde{X}_i)\| \\ (1 - \varepsilon)\|c_i N\| &< \|\tilde{X}_i\| < (1 + \varepsilon)\|c_i N\| \\ (1 - \varepsilon)c_i &< \|\tilde{X}_i\| < (1 + \varepsilon)c_i \end{aligned}$$

$$\begin{aligned} H_i &= X_{i+1} - X_i \\ &= X_{i+1} - X_i + S - S \\ &= \tilde{X}_i - \tilde{X}_{i+1} \end{aligned}$$

For  $t \in [1, 2]$ ,

$$(4.10) \quad \begin{aligned} \|P(X_i + \alpha_i H_i)\| &\leq \|P(X_i + tH_i)\| \\ \|P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| - \|P(X_i + tH_i)\| &< (c_i + 2c_{i+1})\bar{c}\varepsilon \\ \|P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| &> \|P(X_i + tH_i)\| - (c_i + 2c_{i+1})\bar{c}\varepsilon \\ -\|P(\mathbf{P}_{\mathcal{N}}(X_i + \alpha_i H_i) + \mathbf{P}_{\mathcal{M}}(S))\| &> -\|P(X_i + \alpha_i H_i)\| - (c_i + 2c_{i+1})\bar{c}\varepsilon \\ \|P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| - \|P(\mathbf{P}_{\mathcal{N}}(X_i + \alpha_i H_i) + \mathbf{P}_{\mathcal{M}}(S))\| & \\ &> \|P(X_i + tH_i)\| - \|P(X_i + \alpha_i H_i)\| - 2(c_i + 2c_{i+1})\bar{c}\varepsilon \\ &\geq -2(c_i + 2c_{i+1})\bar{c}\varepsilon \\ \|P(X_i + \alpha_i H_i)\| &< \|P(\mathbf{P}_{\mathcal{N}}(X_i + \alpha_i H_i) + \mathbf{P}_{\mathcal{M}}(S))\| + (c_i + 2c_{i+1})\bar{c}\varepsilon \\ \|P(X_i + tH_i)\| &< \|P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))\| + (c_i + 2c_{i+1})\bar{c}\varepsilon \end{aligned}$$

$$\text{Approach 1. } \dashv \|S - X_i - \alpha_i H_i\| \leq \|\tilde{X}_{i+1}\|$$

$$\begin{aligned} \text{Approach 2. } \dashv \|S - (X_i + \alpha_i H_i)\| &= \|(1 - \alpha_i)\tilde{X}_i + \alpha_i \tilde{X}_{i+1}\| > \|\tilde{X}_{i+1}\| \\ \implies \exists t \in [1, 2] \setminus \{\alpha_i\} ; \quad &\|P(X_i + H_i)\| < \|P(X_i + \alpha_i H_i)\| \end{aligned}$$

$$\text{Approach 3. } \dashv \left| \frac{\|\tilde{X}_{i+1}\|}{\|\tilde{X}_i\|} - \frac{\alpha_i - 1}{\alpha_i} \right| < \gamma = \gamma(\varepsilon)$$

$$\text{Approach 2. Suppose } \|(1 - \alpha_i)\tilde{X}_i + \alpha_i \tilde{X}_{i+1}\| > \|\tilde{X}_{i+1}\|.$$

$$\begin{aligned} P(X_{i+1}) &= -P'[S](\tilde{X}_{i+1}) + O(\|\tilde{X}_{i+1}\|^2) \\ \|P(X_{i+1})\| &\leq c\|\tilde{X}_{i+1}\| < c\|S - X_i - \alpha_i H_i\| \end{aligned}$$

$$\begin{aligned}
P(X_i + tH_i) - P(X_i) &\geq P'[X_i](tH_i) \\
&= -tP'[X_i](P'[X_i]^{-1}(P(X_i))) \\
&= -tP(X_i) \\
P(X_i + tH_i) &\geq (1-t)P(X_i) \quad (??)
\end{aligned}$$

$$\begin{aligned}
\|P(X_i + tH_i)\| &\leq \bar{p}\|(1-t)\tilde{X}_i + t\tilde{X}_{i+1}\| + \bar{c}\|(1-t)\tilde{X}_i + t\tilde{X}_{i+1}\|^2 \\
&\leq (t-1)\bar{p}\|\tilde{X}_i\| + t\bar{p}\|\tilde{X}_{i+1}\| + \bar{c}\|(1-t)\tilde{X}_i + t\tilde{X}_{i+1}\|^2 \\
&< (t-1)\bar{p}\|\tilde{X}_i\| + t\bar{p}\|(1-\alpha_i)\tilde{X}_i + \alpha_i\tilde{X}_{i+1}\| + \bar{c}\|(1-t)\tilde{X}_i + t\tilde{X}_{i+1}\|^2 \\
&\leq (t-1)\bar{p}\|\tilde{X}_i\| + t\bar{p}(\alpha_i - 1)\|\tilde{X}_i\| + t\bar{p}\alpha_i\|\tilde{X}_{i+1}\| + \bar{c}\|(1-t)\tilde{X}_i + t\tilde{X}_{i+1}\|^2
\end{aligned}$$

$$\begin{aligned}
\|P(X_i + tH_i)\| &\leq \bar{p}\|(1-t)\tilde{X}_i + t\tilde{X}_{i+1}\| + \bar{c}\|(1-t)\tilde{X}_i + t\tilde{X}_{i+1}\|^2 \\
&= \bar{p}\|(1-t)\mathbf{P}_{\mathcal{N}}\tilde{X}_i + t\mathbf{P}_{\mathcal{N}}\tilde{X}_{i+1} + (1-t)\mathbf{P}_{\mathcal{M}}\tilde{X}_i + t\mathbf{P}_{\mathcal{M}}\tilde{X}_{i+1}\| \\
&\quad + \bar{c}\|(1-t)\mathbf{P}_{\mathcal{N}}\tilde{X}_i + t\mathbf{P}_{\mathcal{N}}\tilde{X}_{i+1} + (1-t)\mathbf{P}_{\mathcal{M}}\tilde{X}_i + t\mathbf{P}_{\mathcal{M}}\tilde{X}_{i+1}\|^2 \\
&\leq \bar{p}\|(1-t)\mathbf{P}_{\mathcal{N}}\tilde{X}_i + t\mathbf{P}_{\mathcal{N}}\tilde{X}_{i+1}\| + \|(1-t)\mathbf{P}_{\mathcal{M}}\tilde{X}_i\| + \|t\mathbf{P}_{\mathcal{M}}\tilde{X}_{i+1}\| \\
&\quad + \bar{c}\left(\|(1-t)\mathbf{P}_{\mathcal{N}}\tilde{X}_i + t\mathbf{P}_{\mathcal{N}}\tilde{X}_{i+1}\| + \|(1-t)\mathbf{P}_{\mathcal{M}}\tilde{X}_i\| + \|t\mathbf{P}_{\mathcal{M}}\tilde{X}_{i+1}\|\right)^2 \\
&\leq \bar{p}\|(1-t)c_iE + tc_{i+1}E\| + \|(1-t)\mathbf{P}_{\mathcal{M}}\tilde{X}_i\| + \|t\mathbf{P}_{\mathcal{M}}\tilde{X}_{i+1}\| \\
&\quad + \bar{c}\left(\|(1-t)c_iE + tc_{i+1}E\| + \|(1-t)\mathbf{P}_{\mathcal{M}}\tilde{X}_i\| + \|t\mathbf{P}_{\mathcal{M}}\tilde{X}_{i+1}\|\right)^2 \\
&= \bar{p}\|(1-t)c_iE + tc_{i+1}E\| + (t-1)\|\mathbf{P}_{\mathcal{M}}\tilde{X}_i\| + t\|\mathbf{P}_{\mathcal{M}}\tilde{X}_{i+1}\| \\
&\quad + \bar{c}\left(\|(1-t)c_iE + tc_{i+1}E\| + (t-1)\|\mathbf{P}_{\mathcal{M}}\tilde{X}_i\| + t\|\mathbf{P}_{\mathcal{M}}\tilde{X}_{i+1}\|\right)^2
\end{aligned}$$

$$\begin{aligned}
&< \bar{p}|(1-t)c_i + tc_{i+1}| + (t-1)\varepsilon c_i + t\varepsilon c_{i+1} \\
&\quad + \bar{c}(|(1-t)c_i + tc_{i+1}| + (t-1)\varepsilon c_i + t\varepsilon c_{i+1})^2 \\
&= \bar{p}|(1-t)c_i + tc_{i+1}| + (t-1)\varepsilon c_i + t\varepsilon c_{i+1} \\
&\quad + \bar{c}(|(1-t)c_i + tc_{i+1}|^2 + (t-1)^2\varepsilon^2 c_i^2 + t^2\varepsilon^2 c_{i+1}^2 \\
&\quad + 2|(1-t)c_i + tc_{i+1}||t-1|\varepsilon c_i + 2|(1-t)c_i + tc_{i+1}|t\varepsilon c_{i+1} + 2t(t-1)c_i c_{i+1}\varepsilon^2) \\
&= \bar{p}|(1-t)c_i + tc_{i+1}| + ((t-1)c_i + tc_{i+1})\varepsilon \\
&\quad + \bar{c}((1-t)^2 c_i^2 + t^2 c_{i+1}^2 + 2(1-t)tc_i c_{i+1} + (t-1)^2\varepsilon^2 c_i^2 + t^2\varepsilon^2 c_{i+1}^2 \\
&\quad + 2|(1-t)c_i + tc_{i+1}||t-1|\varepsilon c_i + 2|(1-t)c_i + tc_{i+1}|t\varepsilon c_{i+1} + 2t(t-1)c_i c_{i+1}\varepsilon^2) \\
&\leq \bar{p}|(1-t)c_i + tc_{i+1}| + ((t-1)c_i + tc_{i+1})\varepsilon \\
&\quad + \bar{c}((1-t)^2 c_i^2 + t^2 c_{i+1}^2 + 2(1-t)tc_i c_{i+1} + ((t-1)^2 c_i^2 + t^2 c_{i+1}^2 + 2t(t-1)c_i c_{i+1})\varepsilon^2 \\
&\quad + 2(t-1)^2 c_i^2 \varepsilon + 4t(t-1)c_i c_{i+1}\varepsilon + 2t^2 c_{i+1}^2 \varepsilon) \\
&= \bar{p}|(1-t)c_i + tc_{i+1}| + \bar{c}((1-t)^2 c_i^2 + t^2 c_{i+1}^2 + 2(1-t)tc_i c_{i+1}) \\
&\quad + \bar{c}\varepsilon(2(t-1)^2 c_i^2 + 4t(t-1)c_i c_{i+1} + 2t^2 c_{i+1}^2) + ((t-1)c_i + tc_{i+1})\varepsilon \\
&\quad + \bar{c}((t-1)^2 c_i^2 + t^2 c_{i+1}^2 + 2t(t-1)c_i c_{i+1})\varepsilon^2 \\
&= \bar{p}|(1-t)c_i + tc_{i+1}| + \bar{c}((1-t)c_i + tc_{i+1})^2 \\
&\quad + \left(2\bar{c}((t-1)c_i + tc_{i+1})^2 + (t-1)c_i + tc_{i+1}\right)\varepsilon \\
&\quad + \bar{c}((t-1)c_i + tc_{i+1})^2 \varepsilon^2 \\
&= \bar{p}|(1-t)c_i + tc_{i+1}| + \bar{c}((1-t)c_i + tc_{i+1})^2 \\
&\quad + \left(2\bar{c}((t-1)c_i + tc_{i+1})^2 + (t-1)c_i + tc_{i+1}\right)\varepsilon + \bar{c}((t-1)c_i + tc_{i+1})^2 \varepsilon^2 \\
&\leq \bar{p}|(1-t)c_i + tc_{i+1}| + \bar{c}((1-t)c_i + tc_{i+1})^2 \\
&\quad + \left(2\bar{c}(c_i + 2c_{i+1})^2 + c_i + 2c_{i+1}\right)\varepsilon + \bar{c}(c_i + 2c_{i+1})^2 \varepsilon^2
\end{aligned}$$

Approach 3.  $\neg \left| \frac{\|\tilde{X}_{i+1}\|}{\|\tilde{X}_i\|} - \frac{\alpha_i - 1}{\alpha_i} \right| < \gamma = \gamma(\varepsilon).$

$$\begin{aligned}
\left| \frac{\|\tilde{X}_{i+1}\|}{\|\tilde{X}_i\|} - \frac{\alpha_i - 1}{\alpha_i} \right| &= \frac{|\alpha_i \|\tilde{X}_{i+1}\| - (\alpha_i - 1) \|\tilde{X}_i\||}{\alpha_i \|\tilde{X}_i\|} \\
&= \frac{|\alpha_i \|\mathbf{P}_N \tilde{X}_{i+1} + \mathbf{P}_M \tilde{X}_{i+1}\| - (\alpha_i - 1) \|\mathbf{P}_N \tilde{X}_i + \mathbf{P}_M \tilde{X}_i\||}{\alpha_i \|\tilde{X}_i\|} \\
&\leq \frac{|\alpha_i \|\mathbf{P}_N \tilde{X}_{i+1}\| + \alpha_i \|\mathbf{P}_M \tilde{X}_{i+1}\| - (\alpha_i - 1) \|\mathbf{P}_N \tilde{X}_i\| + (\alpha_i - 1) \|\mathbf{P}_M \tilde{X}_i\||}{\alpha_i \|\tilde{X}_i\|} \\
&\leq \frac{|\alpha_i \|\mathbf{P}_N \tilde{X}_{i+1}\| - (\alpha_i - 1) \|\mathbf{P}_N \tilde{X}_i\|| + \alpha_i \|\mathbf{P}_M \tilde{X}_{i+1}\| + (\alpha_i - 1) \|\mathbf{P}_M \tilde{X}_i\|}{\alpha_i \|\tilde{X}_i\|} \\
&< \frac{|\alpha_i c_{i+1} - (\alpha_i - 1) c_i| + \alpha_i \varepsilon \|\mathbf{P}_N \tilde{X}_{i+1}\| + (\alpha_i - 1) \varepsilon \|\mathbf{P}_N \tilde{X}_i\|}{\alpha_i \|\tilde{X}_i\|} \\
&= \frac{|\alpha_i c_{i+1} - (\alpha_i - 1) c_i| + \alpha_i \varepsilon c_{i+1} + (\alpha_i - 1) \varepsilon c_i}{\alpha_i \|\tilde{X}_i\|} \\
&= \frac{|\alpha_i c_{i+1} - (\alpha_i - 1) c_i| + \alpha_i \varepsilon c_{i+1} + (\alpha_i - 1) \varepsilon c_i}{\alpha_i \|\mathbf{P}_N \tilde{X}_i + \mathbf{P}_M \tilde{X}_i\|} \\
&= \frac{|\alpha_i c_{i+1} - (\alpha_i - 1) c_i| + \alpha_i \varepsilon c_{i+1} + (\alpha_i - 1) \varepsilon c_i}{\alpha_i \|\mathbf{P}_N \tilde{X}_i\| - \alpha_i \|\mathbf{P}_M \tilde{X}_i\|} \\
&< \frac{|\alpha_i c_{i+1} - (\alpha_i - 1) c_i| + \alpha_i \varepsilon c_{i+1} + (\alpha_i - 1) \varepsilon c_i}{\alpha_i \|\mathbf{P}_N \tilde{X}_i\| - \alpha_i \varepsilon \|\mathbf{P}_N \tilde{X}_i\|} \\
&= \frac{|\alpha_i c_{i+1} - (\alpha_i - 1) c_i| + \alpha_i \varepsilon c_{i+1} + (\alpha_i - 1) \varepsilon c_i}{\alpha_i (1 - \varepsilon) c_i} \\
&= \frac{|\alpha_i c_{i+1} - (\alpha_i - 1) c_i|}{\alpha_i (1 - \varepsilon) c_i} + \frac{\varepsilon c_{i+1}}{(1 - \varepsilon) c_i} + \frac{(\alpha_i - 1) \varepsilon}{\alpha_i (1 - \varepsilon)}
\end{aligned}$$

a

□

## 5. ADVANCED LINE SEARCH

**Theorem 5.1.** *Define*

$$(5.1) \quad B_l = \sum_{k=l}^n A_k \Phi_{(X_i, H_i)}(k, l)$$

and suppose that there exists  $i_0 \in \mathbb{N}$  such that  $\|\mathbf{P}_M(\tilde{X}_i)\| < \varepsilon \|\mathbf{P}_N(\tilde{X}_i)\|$  for small  $\varepsilon > 0$  and  $i \geq i_0$ . Then,

$$(5.2) \quad \|P(X_i + \alpha_i H_i)\| < \|P(X_{i+i})\| \quad \text{and} \quad \|P(X_i + \alpha_i H_i)\| < \eta,$$

or

$$(5.3) \quad \|X_i + \alpha_i H_i - S\| \leq \|X_{i+i} - S\|,$$

where  $\alpha_i = \min \left\{ t_{jk} \in [1, 2] \left| \sum_{l=0}^n b_{ljk} t_{jk}^l = 0, 1 \leq j \leq m, 1 \leq k \leq m \right. \right\}$  and  $b_{ljk} = [B_l]_{jk}$ .

## 6. NOTE

$$\begin{aligned}
P(\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S)) &= \sum_{k=0}^n A_k (\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))^k \\
&= \sum_{k=0}^n A_k (\mathbf{P}_{\mathcal{N}}(X_i + tH_i) + \mathbf{P}_{\mathcal{M}}(S))^k
\end{aligned}$$

$$\begin{aligned}
&\|D_i\| \\
&= \left\| \sum_{k=1}^n \frac{1}{k!} \sum_{\mathbf{v} \in \{0,1\}^k \setminus \mathbf{0}} P^{(k)}[S] \Psi_{((t-1)\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}), (t-1)\mathbf{P}_{\mathcal{M}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i+1}))}(\mathbf{v}) \right\| \\
&\leq \sum_{k=1}^n \frac{1}{k!} \sum_{\mathbf{v} \in \{0,1\}^k \setminus \mathbf{0}} \|P^{(k)}[S]\| \left\| \Psi_{((t-1)\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}), (t-1)\mathbf{P}_{\mathcal{M}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i+1}))}(\mathbf{v}) \right\| \\
&\leq \sum_{k=1}^n \frac{1}{k!} \sum_{\mathbf{v} \in \{0,1\}^k \setminus \mathbf{0}} \|P^{(k)}[S]\| \left\| (t-1)\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}) \right\|^{k-\mathbf{1}^T \mathbf{v}} \left\| (t-1)\mathbf{P}_{\mathcal{M}}(\tilde{X}_i) - t\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i+1}) \right\|^{\mathbf{1}^T \mathbf{v}} \\
&\leq \sum_{k=1}^n \frac{1}{k!} \sum_{\mathbf{v} \in \{0,1\}^k \setminus \mathbf{0}} \|P^{(k)}[S]\| \left( \left\| (t-1)\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) \right\| + \left\| t\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}) \right\| \right)^{k-\mathbf{1}^T \mathbf{v}} \left( \left\| (t-1)\mathbf{P}_{\mathcal{M}}(\tilde{X}_i) \right\| + \left\| t\mathbf{P}_{\mathcal{M}}(\tilde{X}_{i+1}) \right\| \right)^{\mathbf{1}^T \mathbf{v}} \\
&< \sum_{k=1}^n \frac{1}{k!} \sum_{\mathbf{v} \in \{0,1\}^k \setminus \mathbf{0}} \|P^{(k)}[S]\| \varepsilon^{\mathbf{1}^T \mathbf{v}} \left( \left\| (t-1)\mathbf{P}_{\mathcal{N}}(\tilde{X}_i) \right\| + \left\| t\mathbf{P}_{\mathcal{N}}(\tilde{X}_{i+1}) \right\| \right)^k \\
&= \sum_{k=1}^n \frac{1}{k!} \sum_{l=1}^k \binom{k}{l} \|P^{(k)}[S]\| \varepsilon^l ((t-1)c_i + tc_{i+1})^k \\
&= \sum_{k=1}^n \sum_{l=1}^k \frac{1}{k!} \binom{k}{l} \bar{p}_k \varepsilon^l ((t-1)c_i + tc_{i+1})^k \\
&= \sum_{l=1}^n \varepsilon^l \sum_{k=l}^n \frac{1}{k!} \binom{k}{l} \bar{p}_k ((t-1)c_i + tc_{i+1})^k
\end{aligned}$$

$$\begin{aligned}
S - X &= \mathbf{P}_{\mathcal{N}}(S - X) + \mathbf{P}_{\mathcal{M}}(S - X) \\
&= \mathbf{P}_{\mathcal{N}}(S - X) + E \\
X &= S - \mathbf{P}_{\mathcal{N}}(S - X) - E \\
&= \mathbf{P}_{\mathcal{N}}(X) + \mathbf{P}_{\mathcal{M}}(S) - E
\end{aligned}$$

$$\vdash P(X) \approx P(\mathbf{P}_{\mathcal{N}}(X) + \mathbf{P}_{\mathcal{M}}(S))$$

$$\begin{aligned}
P(X + tH) &= \sum_{k=1}^n \frac{1}{k!} P^{(k)}[S] \left( (t-1)\tilde{X} - t\tilde{X}_+ \right)^k \\
&= \sum_{k=1}^n \frac{1}{k!} P^{(k)}[S] \left( (t-1)\mathbf{P}_{\mathcal{N}}\tilde{X} - t\mathbf{P}_{\mathcal{N}}\tilde{X}_+ + (t-1)\mathbf{P}_{\mathcal{M}}\tilde{X} - t\mathbf{P}_{\mathcal{M}}\tilde{X}_+ \right)^k \\
&= \sum_{k=1}^n \frac{1}{k!} P^{(k)}[S] \left( (t-1)\mathbf{P}_{\mathcal{N}}\tilde{X} - t\mathbf{P}_{\mathcal{N}}\tilde{X}_+ \right)^k + \cdots \\
&= \sum_{k=1}^n \frac{1}{k!} P^{(k)}[S] (\mathbf{P}_{\mathcal{N}}(X + tH) - \mathbf{P}_{\mathcal{N}}(S))^k + \cdots \\
&= \sum_{k=1}^n \frac{1}{k!} P^{(k)}[S] (\mathbf{P}_{\mathcal{N}}(X + tH) + \mathbf{P}_{\mathcal{M}}(S) - S)^k + \cdots \\
&= P(\mathbf{P}_{\mathcal{N}}(X + tH) + \mathbf{P}_{\mathcal{M}}(S)) + \cdots
\end{aligned}$$

$$\begin{aligned}
P(X + tH) &= \sum_{k=0}^n A_k (X + tH)^k \\
&= \sum_{k=0}^n A_k \sum_{l=0}^k \Phi_{(X_i, tH_i)}(k, l) \\
&= \sum_{k=0}^n \sum_{l=0}^k t^l A_k \Phi_{(X_i, H_i)}(k, l) \\
&= \sum_{l=0}^n t^l \sum_{k=l}^n A_k \Phi_{(X_i, H_i)}(k, l)
\end{aligned}$$

$$B_l = \sum_{k=l}^n A_k \Phi_{(X_i, H_i)}(k, l)$$



$$\begin{aligned}
p_i(t) &= \text{tr} \left( \left( \sum_{k=0}^n A_k (X_i + tH_i)^k \right)^T \sum_{k=0}^n A_k (X_i + tH_i)^k \right) \\
&= \text{tr} \left( \left( \sum_{l=0}^n \sum_{k=l}^n t^l A_k \Phi_{(X_i, H_i)}(k, l) \right)^T \left( \sum_{l=0}^n \sum_{k=l}^n t^l A_k \Phi_{(X_i, H_i)}(k, l) \right) \right) \\
&= \text{tr} \left( \left( \sum_{l=0}^n \sum_{k=l}^n t^l \Phi_{(X_i^T, H_i^T)}(k, l) A_k^T \right) \left( \sum_{l=0}^n \sum_{k=l}^n t^l A_k \Phi_{(X_i, H_i)}(k, l) \right) \right) \\
&= \text{tr} \left( \sum_{s=0}^n \sum_{r=s}^n \sum_{l=0}^n \sum_{k=l}^n t^{s+l} \Phi_{(X_i^T, H_i^T)}(r, s) A_r^T A_k \Phi_{(X_i, H_i)}(k, l) \right) \\
&= \sum_{s=0}^n \sum_{r=s}^n \sum_{l=0}^n \sum_{k=l}^n t^{s+l} \text{tr} \left( \Phi_{(X_i^T, H_i^T)}(r, s) A_r^T A_k \Phi_{(X_i, H_i)}(k, l) \right) \\
&= \sum_{s=0}^n \sum_{l=0}^n t^{s+l} \sum_{r=s}^n \sum_{k=l}^n \text{tr} \left( \Phi_{(X_i^T, H_i^T)}(r, s) A_r^T A_k \Phi_{(X_i, H_i)}(k, l) \right) \\
&= \sum_{r=0}^n \sum_{s=0}^r \sum_{k=0}^n \sum_{l=0}^k t^{s+l} \text{tr} \left( \Phi_{(X_i^T, H_i^T)}(r, s) A_r^T A_k \Phi_{(X_i, H_i)}(k, l) \right).
\end{aligned}$$

b

$$p_i(t) = \sum_{s=0}^n \sum_{l=0}^n t^{s+l} \sum_{r=s}^n \sum_{k=l}^n \text{tr} \left( \Phi_{(X_i^T, H_i^T)}(r, s) A_r^T A_k \Phi_{(X_i, H_i)}(k, l) \right)$$

Put  $j = s + l$ case 1)  $j \leq n$ 

$$\sum_{s=0}^j t^j \sum_{r=s}^n \sum_{k=j-s}^n \text{tr} \left( \Phi_{(X_i^T, H_i^T)}(r, s) A_r^T A_k \Phi_{(X_i, H_i)}(k, j-s) \right)$$

case 2)  $j > n$ 

$$\sum_{s=j-n}^n t^j \sum_{r=s}^n \sum_{k=j-s}^n \text{tr} \left( \Phi_{(X_i^T, H_i^T)}(r, s) A_r^T A_k \Phi_{(X_i, H_i)}(k, j-s) \right)$$

## REFERENCES

- [1] Attahiru Sule Alfa. Combined elapsed time and matrix-analytic method for the discrete time  $GI/G/1$  and  $GI^X/G/1$  systems. *Queueing Syst.*, 45:5–25, 2003.
- [2] Nigel G. Bean, Leslie W. Bright, Guy Latouche, Charles E. M. Pearce, Philip K. Pollett, and Peter G. Taylor. The quasi-stationary behavior of quasi-birth-and-death processes. *Ann. Appl. Probab.*, 7(1):134–155, Feb 1997.
- [3] Dario A. Bini, Guy Latouche, and Beatrice Meini. *Numerical Methods for Structured Markov Chains*. Oxford University Press Oxford, 2005.
- [4] Geoffrey. J. Butler, Charles R. Johnson, and Henry Wolkowicz. Nonnegative solutions of a quadratic matrix equation arising from comparison theorems in ordinary differential equations. *SIAM J. Algebraic Discrete Methods*, 6(1):47–53, January 1985.
- [5] D. W. Decker, H. B. Keller, and C. T. Kelley. Convergence rates for Newton's method at singular points. *SIAM J. Numer. Anal.*, 20(2):296–314, 1983.
- [6] D. W. Decker and C. T. Kelley. Newton's method at singular points. I. *SIAM J. Numer. Anal.*, 17(1):66–70, 1980.
- [7] D. W. Decker and C. T. Kelley. Newton's method at singular points. II. *SIAM J. Numer. Anal.*, 17(3):465–471, 1980.

- [8] D. W. Decker and C. T. Kelley. Expanded convergence domains for Newton's method at nearly singular roots. *SIAM J. Sci. Statist. Comput.*, 6(4):951–966, 1985.
- [9] Israel Gohberg, Peter Lancaster, and Leiba Rodman. *Matrix Polynomials*. Academic Press, 1982.
- [10] Chun-Hua Guo. Nonsymmetric algebraic Riccati equations and Wiener-Hopf factorization for M-matrices. *SIAM J. Matrix Anal. Appl.*, 23(1):225–242, 2001.
- [11] Chun-hua Guo and Peter Lancaster. Analysis and modification of Newton's method for algebraic Riccati equations. *Math. Comp.*, 67(223):1089–1105, 1998.
- [12] Chun-Hua Guo and Alan J Laub. On the iterative solution of a class of nonsymmetric algebraic Riccati equations. *SIAM J. Matrix Anal. Appl.*, 22(2):376–391, 2000.
- [13] Qi-Ming He and Marcel F. Neuts. On the convergence and limits of certain matrix sequences arising in quasi-birth-and-death Markov chains. *J. Appl. Probab.*, 38(2):519–541, 2001.
- [14] Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, 1985.
- [15] Roger A. Horn and Charles R. Johnson. *Topics in Matrix Analysis*. Cambridge University Press, 1994.
- [16] L. V. Kantorovich and G. P. Akilov. *Functional Analysis in Normed Linear Spaces*. English translation, Pergamon Press, New York, 1964.
- [17] H.B. Keller. Numerical solution of bifurcation and nonlinear eigenvalue problems. In *Applications of bifurcation theory*, pages 359–384, New York, 1977. Proc. Advanced Sem., Univ. Wisconsin, Madison, Wis., 1976, Publ. Math. Res. Center, No. 38, Academic Press.
- [18] C. T. Kelley. A shamanskii-like acceleration scheme for nonlinear equations at singular roots. *Math. Comp.*, 47(176):609–623, 1986.
- [19] C. T. Kelley and R. Suresh. A new acceleration method for Newton's method at singular points. *SIAM J. Numer. Anal.*, 20(5):1001–1009, 1983.
- [20] Hyun-Min Kim. Convergence of Newton's method for solving a class of quadratic matrix equations. *Honam Math. J.*, 30(2):399–409, 2008.
- [21] Peter Lancaster. *Lambda-matrices and Vibrating Systems*. Pergamon Press, 1966.
- [22] Peter Lancaster and Miron Tismenetsky. *The Theory of Matrices with Applications*. Academic Press, 2 edition, 1985.
- [23] Guy Latouche and Vaidyanathan Ramaswami. *Introduction to Matrix Analytic Methods in Stochastic Modeling*. ASA-SIAM, 1999.
- [24] Jie Meng, Sang-hyup Seo, and Hyun-min Kim. Condition numbers and backward error of a matrix polynomial equation arising in stochastic models. *J. Sci. Comput.*, 76(2):759–776, January 2018. First Online : <https://doi.org/10.1007/s10915-018-0641-x>.
- [25] G. W. Reddien. On Newton's method for singular problems. *SIAM J. Numer. Anal.*, 15(5):993–996, 1978.
- [26] Jong Hyeon Seo and Hyun-Min Kim. Solving matrix polynomials by Newton's method with exact line searches. *J. Korean Soc. Ind. Appl. Math.*, 12(2):55–68, 2008.
- [27] Jong-Hyeon Seo and Hyun-Min Kim. Convergence of pure and relaxed Newton methods for solving a matrix polynomial equation arising in stochastic models. *Linear Algebra Appl.*, 440:34–49, 2014.
- [28] Sang-hyup Seo, Jong-Hyeon Seo, and Hyun-Min Kim. Convergence of a modified Newton method for a matrix polynomial equation arising in stochastic problem. *Electron. J. Linear Algebra*, 34:500–513, October 2018.
- [29] Sang-Hyup Seo, Jong-Hyun Seo, and Hyun-Min Kim. Newton's method for solving a quadratic matrix equation with special coefficient matrices. *Honam Math. J.*, 35(3):417–433, 2013.
- [30] Ivar Stakgold. Branching of solutions of nonlinear equations. *SIAM Rev.*, 13(3):289–332, 1971.
- [31] Gilbert Strang. *Linear Algebra and Its Applications*. Thomson, Brooks/Cole, 4th edition, 2006.

SANG-HYUP SEO, WHERE  
 Email address: [saibie1677@gmail.com](mailto:saibie1677@gmail.com)