


---

---

---

---

---




Jan 5, 2026

Part 1: Integrals

Part 2: Application of Integrals

Part 3: Sequences and Series

## Definite Integrals



$$y = f(x)$$

$$\text{If } f(x) > 0$$

$$\int_a^b f(x) dx = \text{area below the graph}$$

Conceptual Interpretations =  
Continuous accumulation  
of quantity.

---

Infinite Integrals  $\int f(x)$

essentially antiderivatives

$x^2 \rightarrow$  antider.  $\frac{x^3}{3}$  so

the most general antider.

is  $\frac{x^3}{3} + C$ .

$\int f(x) dx = F(x) + C$  means

that most general antider.

of  $f(x)$

$$F'(x) = f(x) \quad \int_0^1 x^2 dx = \frac{1}{3}$$

$$\int x^2 dx = \frac{x^3}{3} + C$$

# FTC: Fundamental Theory of Calculus

$$\text{FTC: } \int_0^1 x^2 dx = \left[ \frac{x^3}{3} \right]_0^1 = \frac{1}{3} - \frac{0}{3} = \frac{1}{3}$$

this always works

January 7, 2026

$$(f(x)g(x))' = f'(x)g(x) + f(x)g'(x)$$

$$\int (f'(x)g(x) + f(x)g'(x)) dx = f(x)g(x) + C$$

$$\int f'(x)g(x) dx + \int f(x)g'(x) = f(x)g(x) + C$$

$$\int f(x)g'(x) = f(x)g(x) - \int f'(x)g(x) dx$$

Integration by parts

$$\int x^2 dx - 7 = \frac{x^3}{3} + (C-7) = \int x^2 dx$$

1)

$$\int x \cdot \cos x dx$$

Integration by  
parts

Solution:

$$\begin{aligned}
 \int x \cdot \cos x \, dx &= \int x \cdot (\sin x)' \, dx = \\
 &= x \cdot \sin x - \int x' \sin x \, dx = x \cdot \sin x - \\
 &- \int \sin x \, dx = x \cdot \sin x + \cos x + C
 \end{aligned}$$

$$\begin{aligned}
 2) \int 2^x \cdot x \, dx &= \int \left( \frac{2^x}{\ln 2} \right)' \cdot x \, dx = \\
 &= \frac{2^x}{\ln 2} \cdot x - \int \frac{2^x}{\ln 2} \cdot 1 \, dx =
 \end{aligned}$$

$$= \frac{2^x \cdot x}{\ln 2} - \frac{1}{\ln 2} \int 2^x \, dx \rightarrow \frac{2^x}{\ln 2}$$

$$= \frac{2^x \cdot x}{\ln 2} - \frac{2^x}{(\ln 2)^2}$$

$$\begin{aligned}
 3) \int x^2 e^x dx &= \int x^2 \cdot (e^x)' dx = \\
 &= x^2 e^x - \int (x^2)' e^x dx = x^2 e^x - 2 \int \\
 &= 2 \int \underbrace{x \cdot e^x}_{\text{blue box}} dx = x^2 e^x - 2 \int x \cdot (e^x)' dx \\
 &= x^2 \cdot e^x - 2 \left( x \cdot e^x - \int \boxed{x'}^1 e^x dx \right) = \\
 &= x^2 \cdot e^x - 2 \cdot x \cdot e^x + 2 \boxed{\int e^x dx} e^x \\
 &= \boxed{x^2 e^x - 2x e^x + 2 e^x + C}
 \end{aligned}$$

$$\begin{aligned}
 4) \int \ln x dx &= \int \boxed{1}^{x'} \cdot \ln x dx = \\
 &= \int x' \ln x dx = x \ln x - \int x (\ln x)' dx \\
 &= x \ln x - \int x \cdot \frac{1}{x} dx = x \ln x - \int 1 dx
 \end{aligned}$$

$$= \boxed{x \cdot \ln x - x + C}$$

$$5) \int \cos^2 x \, dx = \int \cos x \cdot (\sin x)' \, dx$$

$$= \cos x \cdot \sin x - \int (\cos x)' \sin x \, dx$$

$$= \cos x \cdot \sin x + \boxed{\int \sin^2 x \, dx}$$

$$= \cos x \cdot \sin x + \int (1 - \cos^2 x) \, dx$$

$$= \cos x \cdot \sin x + \int 1 \, dx - \int \cos^2 x \, dx$$

$$= \cos x \cdot \sin x + x - \int \cos^2 x \, dx$$

Solve for  $\int \cos^2 x \, dx$ :

$$2 \int \cos^2 x \, dx = \frac{1}{2} (\cos x \sin x + x) + C$$

$$u = x^2$$

$$u = g(x)$$

$$\frac{du}{dx} = 2x$$

$$g'(x)$$

$$du = 2x \, dx$$

$$g'(x) \, dx$$

$$\int \underbrace{f(x)}_u \underbrace{g'(x)}_{dv} \, dx = \underbrace{f(x)}_u \underbrace{g(x)}_v - \int \underbrace{g(x)}_v \underbrace{f'(x)}_{du}$$

$$\int u \, dv = uv - \int v \, du$$

1)

$$\int \underbrace{x}_u \cdot \underbrace{\cos x \, dx}_{dv} :$$

$$u = x \quad v = \sin x$$

$$dv = \cos x \cdot dx$$

$$du = dx$$



$$\int \underbrace{x}_u \cdot \underbrace{\cos x \, dx}_{dv} = \int u \, dv = uv - \int v \, du + C$$

$$= x \cdot \sin x - \int \sin x \, dx = x \cdot \sin x - \cos x + C$$


---

January 12, 2026

Substitution Rule (aka "u-sub")

$$\int (f(x) + g(x)) \, dx = \int f(x) \, dx + \int g(x) \, dx$$

$\int f(x) g(x) \, dx \rightarrow$  integration by parts

$$f(x) g(x) - \int f'(x) g(x) \, dx$$

\* often serves the purpose

$\int f(g(x)) \, dx \quad \text{X} \rightarrow$  no such thing

$\int f(g(x)) g'(x) \, dx \rightarrow$  there is a formula for this, substitution rule

$$\int f(g(x))g'(x) dx = \int \underbrace{F'(g(x))g'(x)}_{(F(g(x)))'} dx$$

$$= \int (F(g(x)))' dx = F(g(x)) + C \quad \text{Let } u = g(x)$$

$$= F(u) + C = \int f(u) du \quad \checkmark$$

$$\text{let } u = x^2$$

$$\frac{du}{dx} = 2x$$

$$du = 2x dx$$

$$u = g(x)$$

$$du = g'(x) dx$$

$$\frac{du}{dx} = g'(x)$$

$$\int \underbrace{f(g(x))}_u \underbrace{g'(x)}_{du} dx = \int f(u) du$$

# Practice

1.

$$\int x \cdot \cos(x^2) dx$$

$$\text{let } u = x^2$$

$$du = 2x dx \rightarrow \frac{du}{2}$$

$$\int x \cdot \cos(x^2) dx = \int \cos(u) \cdot \boxed{\frac{du}{2}} =$$

$$\frac{1}{2} \int \cos(u) du = \frac{1}{2} \sin(u) + C =$$

$$\frac{1}{2} \sin(x^2) + C$$

2.

$$\int x^2 \sqrt[3]{4x^3+3} dx$$

$$\text{let } u = 4x^3+3$$

$$du = 12x^2 dx \mid x^2 dx = \frac{du}{12}$$

$$\int x^2 \sqrt[3]{4x^3+3} = \int \sqrt[3]{u} \frac{du}{12} = \frac{1}{2} \int \sqrt[3]{u} du$$

$$= \frac{1}{12} \int u^{\frac{1}{3}} du = \frac{1}{12} \frac{u^{\frac{4}{3}}}{\frac{4}{3}} + C =$$

$$= \frac{u^{\frac{4}{3}}}{16} + C = \boxed{\frac{(4x^3 + 3)^{\frac{4}{3}}}{16}} + C$$

3.

$$\int \sin(10x) dx \quad \begin{array}{l} \text{let } u = 10x \\ du = 10 dx \\ dx = \frac{du}{10} \end{array}$$

$$\int \sin(10x) dx = \int \sin(u) \frac{du}{10} = \frac{1}{10} \int \sin(u) du$$

$$= -\frac{1}{10} \cos(u) + C = \boxed{-\frac{1}{10} \cos(10x) + C}$$

Useful shortcut:

If  $\int f(x) dx = F(x) + C$ , then

$$\cdot \int f(x+a) dx = F(x+a) + C$$

$$\int \cos(x+7) dx = \sin(x+7) + C$$

$$\int f(a \cdot x) dx = \frac{F(a \cdot x)}{a} + C$$

$$\int \cos(7 \cdot x) dx = \frac{\sin(7x)}{7} + C$$

$$5. \int \frac{x^3}{1+x^4} dx$$

Solution:

$$\text{Let } u = 1+x^4$$

$$du = 4x^3 dx \quad x^3 dx = \frac{du}{4}$$

$$\int \frac{x^3}{1+x^4} dx = \int \frac{du/4}{u} = \frac{1}{4} \int \frac{1}{u} du =$$

$$= \frac{1}{4} \ln|x| + C = \boxed{\frac{1}{4} \ln|1+x^4| + C}$$

$$6. \int \frac{x}{1+x^4} dx$$

$$\int \frac{x}{1+x^4} dx = \int \frac{x}{1+(x^2)^2} dx$$

$$\text{Let } u = x^2$$

$$du = 2x dx \quad x dx = \frac{du}{2}$$

$$\int \frac{1}{1+x^4} x dx = \int \frac{1}{1+u^2} \frac{du}{2} =$$

$$= \frac{1}{2} \int \frac{1}{1+u^2} du = \frac{1}{2} \tan^{-1}(u) + c$$

$$= \boxed{\frac{1}{2} \tan^{-1}(x^2) + c}$$

$$7. \int \frac{1}{x^2 + 14x + 130} dx$$

$$\text{Solution: } x^2 + 14x + 130 = (\quad)^2 + c$$

$$(a+b)^2 = a^2 + 2ab + b^2$$

$$x^2 + 14x + 130 = (x^2 + 2 \cdot x \cdot 7 + 7^2) + 7^2 + 130$$

$$= (x+7)^2 + 81 \rightarrow \text{Completing the square}$$

$$\int \frac{1}{(x+7)^2 + 81} dx \quad \text{Let } u = x+7$$

$$du = dx$$

$$\int \frac{1}{u^2 + 81} du = \int \frac{1}{u^2 + 9^2} du =$$

$$= \frac{1}{9} \tan^{-1}\left(\frac{u}{9}\right) + C = \boxed{\frac{1}{9} \tan^{-1}\left(\frac{x+7}{9}\right) + C}$$

11

$$\int \frac{\sin x \cdot \cos x}{1 + \sin^2 x} dx$$

$$\text{Let } u = \sin x \quad du = \cos x \, dx$$

$$\int \frac{\sin x \cos x}{1 + \sin^2 x} dx = \int \frac{u}{1+u^2} du$$

$$\text{Let } v = 1+u^2$$

$$dv = 2u du$$

$$u \cdot du = \frac{dv}{2}$$

$$\int \frac{1}{v} \frac{dv}{2} = \frac{1}{2} \int \frac{1}{v} dv = \frac{1}{2} \ln|v| + c$$

$$\frac{1}{2} \ln(1 + \sin^2 x) + c$$

Alternative:

$$\text{Let } u = 1 + \sin^2 x \quad du = 2 \sin \cdot \cos x$$

$$\begin{aligned} \int \frac{\sin x \cdot \cos x}{1 + \sin^2 x} &= \frac{1}{2} \int \frac{1}{u} du = \frac{1}{2} \ln|u| + c \\ &= \frac{1}{2} \ln(1 + \sin^2 x) + c \end{aligned}$$



$$12. \int x^5 \cdot \cos(x^3) = \int x^2 \cdot x^3 \cdot \cos(x^3) dx$$

Solution: Let  $u = x^3$   
 $du = 3x^2 dx$

$$\int \underbrace{x^2 x^3 \cos(x^3)}_{u \cdot \cos u} dx = \int u \cdot \cos(u) \frac{du}{3} =$$

$$= \frac{1}{3} \int u \cdot \cos u \, du = \frac{1}{3} \int u \cdot (\sin' u) \, du =$$

$$= \frac{1}{3} \left[ u \cdot \sin(u) - \int u' \sin u \, du \right] = \frac{1}{3} ($$

$$= \frac{1}{3} (u \cdot \sin u - \int \sin u \, du) = \frac{1}{3} (u \cdot \sin u + \cos u) + C = \frac{1}{3} (x^3 \cdot \sin x^3 + \cos(x^3)) + C$$