arXiv:2611.08856v1 [stat.ME] 6 Nov 2026

Competitive Balance Beyond Standings: Bayesian Inference for Latent Tiers and Structural Heterogeneity

Guanyu Hu
Department of Statistics and Probability
Michigan State University
huguanyu@msu.edu

Abstract

Competitive balance has long been a central concern in sports economics, tied to outcome uncertainty, fan engagement, and the way leagues design their policies. Most existing measures of competitive balance are built from standings or winning percentages, which say little about how overall parity relates to the way teams sort into latent competitive tiers. We propose a Bayesian framework based on the Bradley–Terry model that jointly estimates the number of competitive tiers and team memberships. To capture the overall degree of competitive heterogeneity, we introduce a structural balance parameter defined on the unit interval within a hierarchical model, and we establish theoretical results giving this parameter a clear interpretation: it is directly tied to the separation between tiers, and overall heterogeneity depends on both the size and frequency of cross-tier differences. Importantly, structural balance is distinct from the information content of the schedule itself, so leagues with different numbers of teams and games can be compared on equal footing. Inference is carried out with an efficient posterior sampling algorithm that jointly handles team abilities, tier structure, home advantage, and structural balance. Simulation studies validate the proposed method, showing accurate recovery of the latent competitive structure and correct performance of the structural-balance testing procedure. Applied to the NBA and NFL, it reveals that the two leagues exhibit similar global structural balance despite very different competitive organizations: the NBA has three latent tiers, while the NFL has two broader ones, and the NFL's sparser schedule produces considerably more posterior uncertainty. Overall parity, it turns out, can mask quite different forms of competitive stratification—something standings-based measures alone cannot detect.

Keywords: Bayesian nonparametrics; Bradley–Terry model; pairwise comparison; sports analytics.

1 Introduction

Competitive balance sits at the heart of professional sports leagues and has long occupied sports economists and league managers. Teams closely matched in underlying strength make for less predictable outcomes and seasons that stay meaningful into their final weeks, and that unpredictability is a large part of what keeps fans engaged — engagement that, in turn, underwrites much of a league's commercial value. Persistent gaps in team quality work the opposite way: they harden into a hierarchical structure where a handful of clubs dominate and the uncertainty of outcome that draws fans in the first place starts to erode. That's why competitive balance comes up so often in debates over league governance, talent allocation, revenue sharing, and the long-run sustainability of competition. Leagues have responded with an array of parity-promoting policies, including player drafts, salary caps, luxury taxes, revenue sharing, and schedule design, and

much of the classical literature in sports economics examines these instruments in exactly this light (Vrooman, 1995; Fort and Quirk, 1995; Szymanski, 2001, 2003; Zimbalist, 2002; Fort and Maxcy, 2003). Measuring competitive balance well matters for a reason beyond description: it's how we tell whether these policies are actually delivering the parity they're designed to produce. The standard toolkit for measuring competitive balance, which includes the dispersion of winning percentages, concentration indices, and related measures (Utt and Fort, 2002; Owen et al., 2007; Manasis et al., 2013; Manasis and Ntzoufras, 2014; Manasis et al., 2022), summarizes realized outcomes well. But it has two real limitations. It mixes persistent differences in underlying team strength with plain game-level noise, and it says almost nothing about how teams sort into latent competitive groups or how uncertain that grouping itself is. As a result, two leagues can look equally balanced on paper while their underlying competitive organization differs substantially, and what looks like a difference in standings may just reflect how much schedule information each league happens to provide.

A related but distinct feature of league competition is the presence of *competitive tiers*. Teams need not fall into clearly distinguishable positions along a complete ranking — several clubs may simply be of comparable latent strength and together form a common competitive level. That means competitive balance and competitive stratification capture different things: two leagues can show similar overall heterogeneity while organizing teams into very different numbers of tiers, and two leagues with the same number of tiers can still differ sharply in how far apart those tiers sit. We treat these as separate inferential problems.

Basini et al. (2023) make the value of latent grouping concrete, using a Bayesian stochastic block model to trace more than four decades of English football and showing how shifts in the number, sizes, and interaction patterns of latent groups surface features of competition that conventional balance indices miss. The natural home for this kind of question is the Bradley–Terry–Luce model (Bradley and Terry, 1952; Luce, 1959), which represents game outcomes through differences in latent ability and underlies a large literature on sports ranking, forecasting, Bayesian inference, and scalable computation (Davidson, 1970; Glickman, 1999; Hunter, 2004; Caron and Doucet, 2012; Mollica and Tardella, 2021; Tian et al., 2024; Pearce and Erosheva, 2025b). But the standard version of that model assigns every competitor a continuous latent ability, so no two competitors ever share a strength level with positive probability — every team is distinguishable from every other whether or not that distinction means anything.

Several lines of work relax this. The ranking lasso groups Bradley–Terry effects through penalized fusion (Masarotto and Varin, 2012); a separate body of Bayesian and mixture-based work handles clustered rankings, partial orders, and related structures more generally (Marden, 1996; Murphy and Martin, 2003; Busse et al., 2007; Dunson and Peddada, 2008; Deng et al., 2014; Li et al., 2022). Two recent papers push this further by inferring latent rank groups directly: Pearce and Erosheva (2025a) propose a rank-clustering Bradley–Terry–Luce model that jointly estimates the number and memberships of latent rank clusters, and Santi and Friel (2026) combine paired-comparison ranking with latent block structure to recover tied competitive levels.

What none of this tells you is how far apart the tiers actually are. A league could have three tightly bunched tiers and a league could have two tiers separated by a canyon, and a method built to answer "who belongs together" has no way to distinguish those two situations — yet they represent very different competitive environments. That's the gap we're after: we infer the tiers, but we also put a number on how much separation exists between them, on a scale that lets you compare that separation across leagues regardless of how many tiers or teams each one has.

We develop a Bayesian extension of the Bradley–Terry model that studies competitive tier structure and global structural balance jointly. A mixture of finite mixtures prior (Miller and Harrison, 2018) is placed on team abilities, so the number of occupied tiers and their memberships

are both learned from the observed games. Our work builds on recent approaches that characterize competitive structure through latent groups: Bayesian stochastic block models describe competition through the number, composition, and interaction patterns of latent blocks (Basini et al., 2023), spike and slab fusion priors identify competitors with tied latent ranks (Pearce and Erosheva, 2025a), and Bradley–Terry stochastic block models jointly infer tied rank groups, their strengths, and their memberships (Santi and Friel, 2026). These methods describe how competitors are grouped, but the grouping alone does not say how strongly the resulting competitive levels are separated. A league with several closely spaced tiers can be more balanced than one with only two tiers sitting far apart. We therefore introduce a global structural balance parameter that measures the dispersion of tier-specific abilities separately from the latent partition. The partition captures competitive stratification, while the balance parameter quantifies how large the heterogeneity among tiers actually is, which gives a common scale for comparing leagues with very different competitive structures. Our theoretical development makes this distinction precise. We characterize the balance parameter through the exact distribution of average between-tier separation, and we show that team-level heterogeneity decomposes into the magnitude of between-tier differences and the prevalence of cross-tier pairs. Greater structural balance, we further show, corresponds to fewer practically distinguishable competitive levels at a fixed resolution, which clarifies why tier count by itself is not an adequate measure of balance. The common scale also lets us move beyond descriptive comparison. We develop Bayesian procedures for comparing structural balance between systems directly, including directional posterior probabilities and practical-equivalence assessments, even when the systems differ in their numbers of teams and comparison designs. Since leagues also differ markedly in their schedules, we study identification under incomplete comparison graphs and separate the underlying structural quantity from the amount of information available to estimate it. Posterior computation combines the MFM representation with Pólya–Gamma augmentation (Polson et al., 2013) and jointly propagates uncertainty in team abilities, tier memberships, home advantage, structural balance, and cross-league comparisons. Our application to the NBA and NFL shows why both the measure and the comparison procedure matter. Despite very different observed standings and schedule densities, the posterior analysis offers little directional evidence that either league is globally more balanced, which points to similar levels of structural balance. The latent competitive organizations are nonetheless quite different: the NBA has three competitive tiers, while the NFL has two broader ones. The NFL's much sparser schedule also leads to greater posterior uncertainty, both in league-specific inference and in the cross-league comparison. Similar overall parity, it turns out, can coexist with different forms of competitive stratification, and formal uncertainty quantification can separate genuine differences in structural balance from differences that merely reflect schedule information.

The remainder of the paper is organized as follows. Section 2 introduces the mixture of finite mixtures model for team abilities, defines the balance parameter B, and describes the MCMC algorithm used for posterior sampling and post MCMC inference procedure. Section 3 develops the theoretical results behind B, and Section 4 turns that interpretation into a practical procedure for comparing balance across two systems. Section 5 reports extensive simulations that check recovery and the comparison procedure under schedules of varying density, and Section 6 applies the framework to the NBA and NFL. We conclude in Section 7 with a summary of the findings and directions for future work.

2 Methodology

2.1 Paired comparisons and latent tier structure

Let $i = 1, \ldots, n$ index observational units and $e = 1, \ldots, M$ index paired comparisons. For comparison e, let a_e and b_e denote the two participating units, and let $Y_e = 1$ if unit a_e is preferred to unit b_e and $Y_e = 0$ otherwise. We assume the Bradley–Terry model (Bradley and Terry, 1952)

$$Y_e \mid \boldsymbol{\theta}, h \sim \text{Bernoulli}(p_e), \qquad \text{logit}(p_e) = \theta_{a_e} - \theta_{b_e} + h, \tag{1}$$

where θ_i is the latent ability of unit i. The parameter h represents a systematic first-position effect, with $h \sim N(0, \sigma_h^2)$ when estimated and $h = 0$ under a symmetric comparison design.

The comparison structure can be represented by an oriented incidence matrix. Let $\boldsymbol{q}_e \in \mathbb{R}^n$ contain a 1 in position a_e, a -1 in position b_e, and zeros elsewhere, and define

$$Q_M = (\boldsymbol{q}_1, \ldots, \boldsymbol{q}_M)^\top.$$

Then the linear predictor is $\boldsymbol{q}_e^\top \boldsymbol{\theta} + h$. This representation connects identifiability of the latent effects to the comparison graph and its incidence matrix (Dong et al., 2026).

We assume that units with comparable latent abilities may share a common tier. Let K denote the nominal number of mixture components and assign the mixture-of-finite-mixtures prior (Miller and Harrison, 2018)

$$\begin{aligned} K &\sim 1 + \text{Poisson}(\lambda_K), \\ \boldsymbol{\pi} \mid K &\sim \text{Dirichlet}(\gamma, \ldots, \gamma), \\ z_i \mid \boldsymbol{\pi}, K &\sim \text{Categorical}(\boldsymbol{\pi}), \qquad \theta_i = \mu_{z_i}. \end{aligned} \tag{2}$$

Conditional on a global precision parameter ϕ,

$$\mu_k \mid \phi, K \overset{\text{iid}}{\sim} N\left(0, \frac{\tau_0^2}{\phi}\right), \qquad k = 1, \ldots, K, \tag{3}$$

where $\tau_0 > 0$ is a fixed reference scale.

Let

$$\mathcal{P} = \{C_1, \ldots, C_T\}$$

denote the occupied partition after empty mixture components are removed, with $T = |\mathcal{P}|$. Thus T is the number of occupied tiers and $\mathcal{P}$ determines their memberships. Mixture labels have no substantive meaning and no ordering constraint is imposed during posterior sampling.

For a fixed partition, let $Z_{\mathcal{P}}$ be the $n \times T$ membership matrix and define

$$D_{\mathcal{P}} = Q_M Z_{\mathcal{P}}.$$

Then

$$\boldsymbol{\theta} = Z_{\mathcal{P}} \boldsymbol{\mu}, \qquad \boldsymbol{\psi} = D_{\mathcal{P}} \boldsymbol{\mu} + h \mathbf{1}_M.$$

For a connected comparison graph, $\text{rank}(D_{\mathcal{P}}) = T - 1$, so the occupied tier effects are identifiable up to a common location. A precise rank condition is given in Section 3.1.

2.2 Structural balance

We assign

$$\phi \sim \text{Gamma}(a_\phi, b_\phi),$$

using the shape-rate parameterization, and define

$$B = \frac{\phi}{1+\phi}, \qquad \phi = \frac{B}{1-B}, \qquad 0 < B < 1. \tag{4}$$

The parameter B measures the global magnitude of structural balance. Larger values of B imply greater precision and hence smaller separation among the latent tier effects.

For two distinct tiers c and d,

$$\mu_c - \mu_d \mid B \sim N\left(0, 2\tau_0^2 \frac{1-B}{B}\right),$$

and therefore

$$E\left\{(\mu_c - \mu_d)^2 \mid B\right\} = 2\tau_0^2 \frac{1-B}{B}.$$

For $T \geq 2$, define the average squared separation

$$D_T = \frac{2}{T(T-1)} \sum_{c<d} (\mu_c - \mu_d)^2. \tag{5}$$

Then

$$E(D_T \mid B, T) = 2\tau_0^2 \frac{1-B}{B}, \tag{6}$$

which does not depend on T. Thus the latent partition $(T, \mathcal{P})$ describes the tier structure, whereas B measures the magnitude of between-tier heterogeneity.

The interpretation also carries to the paired-comparison scale. In the absence of a position effect,

$$p_{cd} = \text{expit}(\mu_c - \mu_d).$$

As $B \to 1$, the tier contrasts shrink toward zero and $p_{cd} \to 1/2$. Hence larger values of B correspond to greater structural balance.

2.3 Posterior computation

We develop a semi-collapsed Gibbs sampler (Neal, 2000) that combines the MFM partition representation with Pólya–Gamma augmentation (Polson et al., 2013). The mixture weights and nominal component count are integrated out when updating the occupied partition, while the effect of a proposed new tier is integrated analytically. The sampler therefore avoids reversible-jump moves and does not require the number of occupied tiers to be fixed in advance. Full derivations of the sampling updates are provided in the Supporting Materials.

For the current partition, define

$$\psi_e = \boldsymbol{d}_e^\top \boldsymbol{\mu} + h, \qquad \kappa_e = Y_e - \frac{1}{2},$$

where $\boldsymbol{d}_e^\top$ is row e of $D_\mathcal{P}$. Introduce the Pólya–Gamma latent variables

$$\omega_e \mid \psi_e \sim \text{PG}(1, \psi_e), \qquad e = 1, \ldots, M. \tag{7}$$

Let

$$\Omega = \mathrm{diag}(\omega_1, \ldots, \omega_M), \qquad \boldsymbol{\kappa} = (\kappa_1, \ldots, \kappa_M)^\top.$$

Conditional on $\boldsymbol{\omega}$,

$$\begin{pmatrix} \boldsymbol{\mu} \\ h \end{pmatrix} \Big| \text{ rest} \sim N\left(\mathcal{Q}^{-1}\boldsymbol{b}, \mathcal{Q}^{-1}\right), \tag{8}$$

where

$$\mathcal{Q} = \begin{pmatrix} D_{\mathcal{P}}^\top \Omega D_{\mathcal{P}} + \phi\tau_0^{-2} I_T & D_{\mathcal{P}}^\top \Omega \mathbf{1}_M \\ \mathbf{1}_M^\top \Omega D_{\mathcal{P}} & \mathbf{1}_M^\top \Omega \mathbf{1}_M + \sigma_h^{-2} \end{pmatrix},$$

and

$$\boldsymbol{b} = \begin{pmatrix} D_{\mathcal{P}}^\top \boldsymbol{\kappa} \\ \mathbf{1}_M^\top \boldsymbol{\kappa} \end{pmatrix}.$$

When $h = 0$, the last row and column are omitted.

Integrating out $(K, \boldsymbol{\pi})$ gives

$$p(\mathcal{P}) = V_n(T) \prod_{c=1}^{T} \gamma^{(n_c)}, \qquad V_n(T) = \sum_{k=T}^{\infty} \frac{k_{[T]}}{(\gamma k)^{(n)}} p_K(k), \tag{9}$$

where $n_c = |C_c|$, $x^{(r)}$ denotes a rising factorial, and $x_{[r]}$ denotes a falling factorial.

To update z_i, remove unit i from its current tier and let t denote the number of remaining occupied tiers. For an existing tier c,

$$W_{ic} = (n_{-i,c} + \gamma) \exp\{\ell_i(c)\}, \tag{10}$$

where $\ell_i(c)$ is the Pólya–Gamma augmented log-likelihood contribution from comparisons involving unit i.

For a proposed new tier, define

$$A_i = \frac{\phi}{\tau_0^2} + \sum_{e \in \mathcal{E}_i} \omega_e, \qquad C_i = \sum_{e \in \mathcal{E}_i} s_{ie}(\kappa_e - \omega_e r_{ie}),$$

and

$$H_i = \sum_{e \in \mathcal{E}_i} \left(\kappa_e r_{ie} - \frac{\omega_e}{2} r_{ie}^2\right).$$

The integrated contribution of the proposed tier is

$$m_i^{\text{new}} = \exp(H_i) \left(\frac{\phi/\tau_0^2}{A_i}\right)^{1/2} \exp\left(\frac{C_i^2}{2A_i}\right),$$

giving

$$W_i^{\text{new}} = \gamma \frac{V_n(t+1)}{V_n(t)} m_i^{\text{new}}. \tag{11}$$

If a new tier is selected,

$$\mu_{\text{new}} \mid \text{rest} \sim N\left(\frac{C_i}{A_i}, \frac{1}{A_i}\right). \tag{12}$$

Conditional on the occupied tier effects,

$$\phi \mid \boldsymbol{\mu}, T \sim \text{Gamma}\left(a_\phi + \frac{T}{2},\ b_\phi + \frac{1}{2\tau_0^2} \sum_{c=1}^{T} \mu_c^2\right), \tag{13}$$

after which $B = \phi/(1+\phi)$.

Algorithm 1 summarizes the sampler.

Algorithm 1 Semi-collapsed Gibbs sampler

Initialize $\mathcal{P}^{(0)}$, $\boldsymbol{\mu}^{(0)}$, $h^{(0)}$, and $\phi^{(0)}$.
for $s = 1, \ldots, S$ **do**
  Compute $\psi_e = \boldsymbol{d}_e^\top \boldsymbol{\mu} + h$ and sample ω_e according to (7).
  for $i = 1, \ldots, n$ **do**
    Remove unit i from its current tier and delete the tier if it becomes empty.
    Compute the existing-tier weights in (10) and the new-tier weight in (11).
    Sample z_i from the normalized allocation weights.
    if a new tier is selected **then**
      Sample μ_{new} from (12).
    end if
  end for
  Construct the updated tier design matrix $D_{\mathcal{P}}$.
  Sample $(\boldsymbol{\mu}, h)$ from (8).
  Sample ϕ from (13) and set $B = \phi/(1+\phi)$.
  Store $\mathcal{P}$, T, $\boldsymbol{\mu}$, h, ϕ, and B.
end for

2.4 Posterior summaries

The MCMC output supports inference on everything the model estimates: the number of occupied tiers, their memberships, the tier effects, and the structural balance parameter. The posterior distribution of the occupied tier count is estimated by

$$\widehat{\Pr}(T = t \mid \boldsymbol{Y}) = \frac{1}{S} \sum_{s=1}^{S} \mathbf{1}\{T^{(s)} = t\}.$$

Because any single draw of the partition can be misleading, we summarize partition uncertainty through the posterior similarity matrix

$$\widehat{P}_{ij} = \frac{1}{S} \sum_{s=1}^{S} \mathbf{1}\left\{z_i^{(s)} = z_j^{(s)}\right\},$$

so that $\widehat{P}_{ij}$ is the posterior probability that units i and j belong to the same tier.

When a single representative partition is needed, we use the least-squares criterion of Dahl (2006). Let

$$A_{ij}^{(s)} = \mathbf{1}\left\{z_i^{(s)} = z_j^{(s)}\right\}$$

be the co-clustering matrix from posterior draw s. The Dahl partition is

$$\widehat{\mathcal{P}}_{\text{Dahl}} = \mathcal{P}^{(s^*)}, \qquad s^* = \arg\min_{1 \le s \le S} \sum_{i<j} \left(A_{ij}^{(s)} - \widehat{P}_{ij}\right)^2. \tag{14}$$

Since the criterion involves only pairwise co-clustering indicators, it is invariant to mixture labels and no post-processing for label switching is required.

For interpretation, the tiers in $\widehat{\mathcal{P}}_{\text{Dahl}}$ are ordered by their posterior tier effects. Structural balance is summarized by the posterior distribution of B, with its posterior median and credible interval as the primary summaries. When several paired-comparison populations are analyzed together, the posterior draws of B feed directly into the Bayesian comparisons of Section 4.

3 Theoretical Properties

This section studies two theoretical properties of the proposed framework. We first consider identifiability under incomplete comparison designs and then provide a structural interpretation of the balance parameter. These results clarify when the model can learn team differences from an incomplete schedule and how the balance parameter can be interpreted in practical comparisons of competitive structure. Proofs of all results are provided in the Supporting Materials.

3.1 Identification under incomplete comparison designs

The Bradley–Terry likelihood involves the latent abilities only through pairwise differences. Therefore, shifting all tier effects by a common constant leaves the likelihood unchanged. The common location is not identified, but under a connected comparison design all contrasts among occupied tiers are identifiable.

Proposition 1 (Graph identification of tier contrasts)**.** *Suppose the observed comparison graph is connected and $\mathcal{P} = \{C_1, \dots, C_T\}$ contains T nonempty blocks. Then*

$$\operatorname{rank}(D_{\mathcal{P}}) = T - 1,$$

or equivalently,

$$\operatorname{null}(D_{\mathcal{P}}) = \operatorname{span}\{\mathbf{1}_T\}.$$

Thus all contrasts among the occupied tier effects are identifiable.

If the first-position effect h is included, let $H_{\mathcal{P}}$ be a $T \times (T-1)$ contrast matrix satisfying

$$H_{\mathcal{P}}^{\top}\mathbf{1}_T = 0, \qquad H_{\mathcal{P}}^{\top}H_{\mathcal{P}} = I_{T-1}.$$

Then $(\boldsymbol{\beta}_{\mathcal{P}}, h)$ is identifiable whenever

$$\operatorname{rank}\left[D_{\mathcal{P}}H_{\mathcal{P}}, \mathbf{1}_M\right] = T.$$

Proposition 1 distinguishes identifiability from estimation precision. Connectivity ensures that occupied tier contrasts are identifiable, whereas the number and distribution of observed comparisons determine how precisely these contrasts can be estimated.

To characterize estimation under increasing comparison information, suppose the true occupied partition is $\mathcal{P}^*$ and consider a fixed-n regime with $M \to \infty$. Assume that the limiting comparison graph is connected, the true comparison probabilities are bounded away from zero and one, and

$$\frac{1}{M}X_{\mathcal{P}^*,M}^{\top}W_M^*X_{\mathcal{P}^*,M} \longrightarrow \mathcal{I}^*, \tag{15}$$

where $\mathcal{I}^*$ is positive definite.

Proposition 2 (Conditional posterior concentration)**.** *Suppose the prior density for the identifiable tier contrasts and h is positive and continuous in a neighborhood of the true parameter. Conditional on $\mathcal{P} = \mathcal{P}^*$, for any sequence $r_M \to \infty$,*

$$\Pi\left[\|\boldsymbol{\vartheta} - \boldsymbol{\vartheta}^*\| > \frac{r_M}{\sqrt{M}} \;\middle|\; Y^{(M)}, \mathcal{P}^*\right] \longrightarrow 0$$

in P^*-probability.

Proposition 2 shows that, conditional on the occupied partition, increasing the number of informative comparisons improves estimation of the tier contrasts at the usual parametric rate. Thus the amount of schedule information affects estimation precision but does not change the definition of structural balance.

3.2 Structural interpretation of competitive balance

From (4),

$$\phi = \frac{B}{1-B}, \qquad 0 < B < 1,$$

and the occupied tier effects satisfy

$$\mu_c \mid B \overset{\text{iid}}{\sim} N\left(0, \tau_0^2 \frac{1-B}{B}\right).$$

Therefore, for two distinct tiers,

$$\mu_c - \mu_d \mid B \sim N\left(0, 2\tau_0^2 \frac{1-B}{B}\right).$$

Recall the average squared separation D_T defined in (5).

Theorem 1 (Exact distribution of tier dispersion)**.** *Conditional on B and $T \geq 2$,*

$$\frac{(T-1)B}{2\tau_0^2(1-B)} D_T \sim \chi^2_{T-1}. \tag{16}$$

Consequently,

$$E(D_T \mid B, T) = 2\tau_0^2 \frac{1-B}{B},$$

and

$$\mathrm{Var}(D_T \mid B, T) = \frac{8\tau_0^4}{T-1} \left(\frac{1-B}{B}\right)^2.$$

Moreover, if $0 < B_2 < B_1 < 1$, then

$$|\mu_c - \mu_d| \mid B_1 \leq_{\text{st}} |\mu_c - \mu_d| \mid B_2.$$

Theorem 1 provides a direct interpretation of B. The expected average between-tier separation does not depend on the number of occupied tiers, the number of competitors, or the number of observed comparisons. Therefore, B provides a common measure of between-tier heterogeneity across different competitive systems.

To connect this quantity with team-level heterogeneity, define

$$D_n^{\text{unit}} = \frac{2}{n(n-1)} \sum_{1 \leq i < j \leq n} (\theta_i - \theta_j)^2$$

and

$$q(\mathcal{P}) = 1 - \frac{\sum_{c=1}^{T} n_c(n_c-1)}{n(n-1)}.$$

Corollary 1 (Population-level dispersion decomposition)**.** *Conditional on B and $\mathcal{P}$,*

$$E\left(D_n^{\text{unit}} \mid B, \mathcal{P}\right) = q(\mathcal{P})\, 2\tau_0^2 \frac{1-B}{B}.$$

Corollary 1 shows that overall team-level heterogeneity depends on two features: the magnitude of separation between occupied tiers, measured by $2\tau_0^2(1-B)/B$, and the proportion of team pairs belonging to different tiers, measured by $q(\mathcal{P})$. Therefore, two systems may have similar structural balance while exhibiting different levels of overall heterogeneity because their tier compositions differ.

3.3 Structural balance and effective tier complexity

The occupied tier count T is learned from the posterior partition and is not determined by B. However, B is related to the number of competitive levels that remain distinguishable at a given practical resolution.

Fix $\delta > 0$ and define a threshold graph on the occupied tiers by connecting c and d whenever

$$|\mu_c - \mu_d| \le \delta.$$

Let $T_\delta(B)$ denote the number of connected components of this graph.

Proposition 3 (Balance and effective tier complexity)**.** *Fix $T \ge 2$ and $\delta > 0$, and write*

$$\mu_c(B) = \tau_0 \left(\frac{1-B}{B}\right)^{1/2} Z_c, \qquad Z_c \overset{\text{iid}}{\sim} N(0,1).$$

Under this common-Z coupling, if $B_1 > B_2$, then

$$T_\delta(B_1) \le T_\delta(B_2) \qquad \textit{almost surely},$$

and consequently

$$E\{T_\delta(B_1) \mid T\} \le E\{T_\delta(B_2) \mid T\}.$$

For any two distinct occupied tiers,

$$\Pr\left(|\mu_c - \mu_d| > \delta \mid B\right) = 2\Phi\left[-\frac{\delta}{\sqrt{2}\tau_0}\left(\frac{B}{1-B}\right)^{1/2}\right],$$

which is strictly decreasing in B.

Proposition 3 shows that greater structural balance leads to smaller differences among occupied tiers and therefore fewer practically distinguishable competitive levels at a fixed resolution δ. This result concerns effective tier complexity and does not imply that the raw occupied MFM tier count T must decrease monotonically with B in every finite data set.

Finally, for a neutral comparison between tiers c and d, let

$$p_{cd} = \operatorname{expit}(\mu_c - \mu_d).$$

Then

$$E\left[\{\operatorname{logit}(p_{cd})\}^2 \mid B\right] = 2\tau_0^2 \frac{1-B}{B},$$

so larger values of B correspond to smaller variation in pairwise log-odds contrasts.

4 Bayesian Comparison of Structural Balance Across Systems

The results in Section 3.2 provide a common scale for comparing competitive balance across paired-comparison systems. Consider two systems indexed by $\ell = 1, 2$, with balance parameters B_1 and B_2. The systems may differ in the number of competitors, occupied tiers, and observed comparisons, but are analyzed using the same reference scale τ_0.

By Theorem 1, define

$$d(B) = E(D_T \mid B, T) = 2\tau_0^2 \frac{1-B}{B}.$$

Because $d(B)$ is strictly decreasing on $(0, 1)$, larger values of B correspond to smaller between-tier separation and therefore greater structural balance. Importantly, $d(B)$ does not depend on the number of competitors, the number of occupied tiers, or the number of observed comparisons. These features affect posterior uncertainty and the realized competitive structure, but not the interpretation of structural balance. Therefore, the same scale can be used to compare systems with different competitive structures and schedules.

4.1 Directional comparison of structural balance

Define

$$\Delta_B = B_1 - B_2.$$

Because $d(B)$ is strictly decreasing,

$$\Delta_B > 0 \quad \Longleftrightarrow \quad d(B_1) < d(B_2).$$

Thus $\Delta_B > 0$ indicates greater structural balance in system 1, whereas $\Delta_B < 0$ indicates greater structural balance in system 2.

Given posterior draws,

$$\Delta_B^{(s)} = B_1^{(s)} - B_2^{(s)}.$$

We summarize the posterior distribution of Δ_B using its median, credible interval, and the directional probability

$$P_{12} = \Pr(B_1 > B_2 \mid Y_1, Y_2).$$

Values near one provide evidence that system 1 is more balanced, values near zero favor system 2, and values near one half indicate little directional evidence.

4.2 Practical differences in structural balance

A directional comparison does not indicate whether the difference in structural balance is practically meaningful. We therefore define

$$\Delta_d = d(B_2) - d(B_1),$$

so that

$$\Delta_d > 0 \quad \Longleftrightarrow \quad B_1 > B_2.$$

For a prespecified practical margin $\epsilon_d > 0$, consider

$$H_- : \Delta_d < -\epsilon_d, \qquad H_0 : |\Delta_d| \leq \epsilon_d, \qquad H_+ : \Delta_d > \epsilon_d.$$

Here, H_- represents meaningfully greater balance in system 2, H_0 represents practical similarity, and H_+ represents meaningfully greater balance in system 1.

The corresponding posterior probabilities are

$$\begin{aligned}
\pi_- &= \Pr(\Delta_d < -\epsilon_d \mid Y_1, Y_2), \\
\pi_0 &= \Pr(|\Delta_d| \leq \epsilon_d \mid Y_1, Y_2), \\
\pi_+ &= \Pr(\Delta_d > \epsilon_d \mid Y_1, Y_2),
\end{aligned}$$

where

$$\pi_- + \pi_0 + \pi_+ = 1.$$

The vector (π_-, π_0, π_+) distinguishes evidence for a meaningful difference from evidence for practical similarity. In particular, a directional probability near one half does not imply practical equivalence.

For posterior draws,

$$\Delta_d^{(s)} = 2\tau_0^2 \left\{ \frac{1 - B_2^{(s)}}{B_2^{(s)}} - \frac{1 - B_1^{(s)}}{B_1^{(s)}} \right\}.$$

For calibration, the root mean squared tier separation is

$$s(B) = \{d(B)\}^{1/2} = \sqrt{2}\tau_0 \left(\frac{1 - B}{B} \right)^{1/2},$$

which is expressed on the latent Bradley–Terry log-odds scale. This provides a direct way to specify ϵ_d in terms of a practically meaningful difference in tier separation.

Evidence for practical similarity can also be summarized using the interval Bayes factor

$$BF_{01}^{(\epsilon)} = \frac{\Pr(H_0 \mid Y_1, Y_2) / \Pr(H_1 \mid Y_1, Y_2)}{\Pr(H_0) / \Pr(H_1)}, \qquad H_1 : |\Delta_d| > \epsilon_d.$$

Values greater than one indicate increased support for practical similarity relative to the prior, whereas values below one favor a meaningful difference. We report this quantity as a complementary posterior summary together with (π_-, π_0, π_+).

5 Simulation study

We conducted a factorial simulation study to evaluate the proposed framework under comparison structures and competitive configurations motivated by professional sports applications. The simulations address two related questions. First, we examine how accurately the model recovers latent team abilities, competitive tiers, and structural balance under schedules with different amounts of comparison information. Second, we evaluate whether the proposed posterior comparison of structural balance remains informative when two systems differ substantially in their numbers of teams and observed comparisons.

5.1 Estimation and latent-tier recovery

The simulations use $n = 30$ teams under three comparison-graph designs. The first is the observed NBA regular-season schedule with 1,230 games. To see what happens as comparison information is stripped away while keeping the graph connected, we add a complete double round-robin with 870 games and a complete single round robin with 435 games. The three designs step down in informativeness, which lets us watch how schedule density affects recovery of the latent competitive structure.

Each design is crossed with two structural settings,

$$(B, T) = (0.2, 4) \qquad \text{and} \qquad (B, T) = (0.5, 3),$$

a less balanced system with four tiers and a more balanced one with three. The second setting is deliberately the harder one for tier recovery: since larger B means smaller between-tier separation, its tiers sit closer together even though there are fewer of them. Within each setting we try both balanced and unbalanced tier allocations,

$$(8, 8, 7, 7) \qquad \text{and} \qquad (3, 12, 12, 3)$$

for four tiers, and

$$(10, 10, 10) \qquad \text{and} \qquad (5, 20, 5)$$

for three. The unbalanced layouts test whether recovery suffers when most teams pile into one or two levels instead of spreading across tiers.

Team labels are randomly assigned to tiers in each replicate. With $\tau_0 = 1$, the tier abilities are scaled so that the implied structural balance matches the target exactly, and games are generated from

$$\Pr(Y_{ij} = 1 \mid \theta, h) = \text{logit}^{-1}(\theta_i - \theta_j + h), \qquad h = 0.35.$$

We ran 100 independent replicates for each of the 12 simulation settings. For each replicate, the MFM–BT model was fitted using two independently initialized MCMC chains. Each chain consisted of 6,000 iterations, with the first 1,500 iterations discarded as burn-in and every third subsequent iteration retained, yielding 1,500 posterior draws per chain. We used $\lambda_K = 1$, $\tau_0 = 1$, and $\gamma = 1$ throughout the simulation study. The number of occupied tiers is inferred from the data rather than fixed at the truth. Clustering accuracy is measured by the adjusted Rand index (ARI) (Hubert and Arabie, 1985) of the Dahl representative partition, and tier-count recovery by

$$100R^{-1} \sum_{r=1}^{R} I(\widehat{T}_r = T).$$

Ability estimation is judged by mean squared error (MSE), ranking by the Spearman correlation between estimated and true ability ranks, and the system-level parameters by B-MSE and h-MSE from their posterior means. These criteria deliberately separate recovery of the discrete tier structure from estimation of the continuous balance parameter.

We compare MFM–BT with three two-stage benchmark procedures: BT–GMM, which clusters ordinary Bradley–Terry ability estimates using a Gaussian mixture model (McLachlan and Peel, 2000); BT–DP, which applies a DP-means clustering procedure to the Bradley–Terry estimates (Kulis and Jordan, 2012); and BT–pairwise-fusion, which uses an all-pairs L_1 fusion penalty motivated by the ranking-lasso method (Masarotto and Varin, 2012), all fitted to exactly the same graphs and outcomes. Each benchmark selects its own number of clusters from its fitted model, since in practice neither the number nor the composition of competitive tiers is known in advance.

Figure 1 summarizes clustering recovery across the 12 simulation settings. The upper row reports the adjusted Rand index (ARI), and the lower row reports the percentage of replicates that recover the true number of tiers. Within each graph design, the four bars correspond to the two (B, T) settings under balanced and unbalanced tier allocations. For MFM–BT and BT–DP, the reported partitions are obtained from the Dahl-selected posterior representative partition.

MFM–BT achieves the best clustering performance across nearly all settings. Its advantage is especially clear for the sparse single-round-robin design and the more difficult $(B, T) = (0.5, 3)$ setting, where the tier differences are smaller. BT–DP is generally the strongest benchmark in terms of ARI, but it tends to overestimate the number of tiers. On average, it estimates approximately 5.13 tiers when the true number is either 3 or 4, which explains its relatively low true-tier recovery rate. BT–GMM performs moderately well under the denser designs but loses accuracy as comparison information decreases. It tends to merge some of the true tiers, leading to fewer estimated tiers than the truth. BT–pairwise-fusion has the lowest ARI and consistently underestimates the number of tiers, recovering the true tier count in none of the simulation replicates.

Table 1 summarizes performance across the 12 simulation settings, with 100 replicates for each setting. MFM–BT achieves the highest average ARI, 0.858, and recovers the true number of tiers in 81.8% of replicates. The genuine Bayesian BT–DP benchmark performs second, with an average




Figure 1: Clustering performance across the 12 simulation designs.

ARI of 0.774 and a true-tier recovery rate of 26.2%. BT–GMM recovers the true tier count in 29.7% of replicates, while BT–pairwise-fusion never recovers the true count. MFM–BT also has the smallest ability MSE and the highest rank correlation. Thus, its advantage is not limited to partition recovery: jointly modeling tier memberships and abilities also improves continuous ability estimation and the induced team ranking.

Table 1: Average performance over the 12 simulation settings and 100 replicates per setting. Entries are means with standard deviations in parentheses.

Method	ARI	True T (%)	Ability MSE	Rank correlation
MFM–BT	0.858 (0.182)	81.8 (38.6)	0.137 (0.162)	0.959 (0.067)
BT–GMM	0.610 (0.290)	29.7 (45.7)	0.378 (1.811)	0.915 (0.054)
BT–DP	0.774 (0.250)	26.2 (44.0)	0.144 (0.164)	0.952 (0.069)
BT–pairwise-fusion	0.433 (0.168)	0.0 (0.0)	1.597 (1.136)	0.913 (0.059)

This gap between methods matters for the application we have in mind. A competitive-balance study needs more than a ranking of teams; it needs a stable account of which teams are practically indistinguishable in latent strength, and the much higher ARI of MFM-BT says its posterior partition captures that organization far more faithfully. At the same time, the strong rank correlations across several methods remind us that producing a ranking is much easier than recovering the tier structure beneath it, so high ranking accuracy on its own says little about stratification.

Table 2 shows the MFM-BT estimation results by design. Under the NBA schedule recovery is excellent, with ARI between 0.948 and 0.991 and true tier counts recovered in 79% to 97% of replicates. Performance slips moderately under the double round robin and more sharply under the single round robin, worst of all at $(B, T) = (0.5, 3)$, where ARI falls to 0.577–0.614. The pattern is exactly what the structure predicts: $B = 0.5$ puts the tiers closer together, and with fewer games it becomes harder to say which side of a boundary a team belongs on.

The balance parameter is far more stable than individual tier memberships. B-MSE stays small in nearly every design even where ARI drops substantially: under the single-round-robin settings with $(B, T) = (0.5, 3)$, ARI sits below 0.62 while B-MSE remains between 0.003 and 0.006. B summarizes the overall scale of heterogeneity rather than the fate of any one team, so it doesn't need every team to land in its exact true tier to stay well estimated.

Table 2: MFM-BT performance by graph design, structural setting, and tier allocation.

Graph design	(B, T)	Allocation	ARI	True T (%)	Ability MSE	B-MSE	h-MSE
NBA	(0.2,4)	Balanced	0.990	97	0.058	0.002	0.007
NBA	(0.2,4)	Unbalanced	0.991	96	0.036	0.003	0.004
NBA	(0.5,3)	Balanced	0.948	79	0.032	0.005	0.004
NBA	(0.5,3)	Unbalanced	0.955	81	0.026	0.004	0.004
Double RR	(0.2,4)	Balanced	0.951	85	0.104	0.003	0.012
Double RR	(0.2,4)	Unbalanced	0.964	86	0.082	0.004	0.009
Double RR	(0.5,3)	Balanced	0.891	76	0.064	0.005	0.007
Double RR	(0.5,3)	Unbalanced	0.898	82	0.050	0.004	0.006
Single RR	(0.2,4)	Balanced	0.752	73	0.425	0.003	0.025
Single RR	(0.2,4)	Unbalanced	0.768	67	0.329	0.010	0.015
Single RR	(0.5,3)	Balanced	0.614	85	0.229	0.006	0.014
Single RR	(0.5,3)	Unbalanced	0.577	74	0.209	0.003	0.010

Losing games mainly hurts local structure. Ability MSE rises noticeably as schedules thin out, and tier assignments near close levels become ambiguous, while the global balance signal survives because it pools information across the whole league – exactly the separation between partition uncertainty and B-uncertainty the framework is built around.

Unbalanced tier sizes cause no systematic damage, either. In several settings the unbalanced allocation does as well as, or slightly better than, the balanced one, so the method does not rely on roughly equal groups – a good sign for sports data, where tiers routinely contain very different numbers of teams.

Taken together, the simulations bear out the separate roles of the partition and the balance parameter. With an informative schedule, MFM-BT recovers abilities, rankings, and tiers accurately; as comparisons grow scarce, uncertainty about individual abilities and memberships rises, especially when tiers sit close together, while B remains comparatively steady. B tracks the magnitude of structural heterogeneity; the comparison graph only governs how precisely that structure can be learned from the games at hand – which is exactly the division of labor the theory predicts.

5.2 Validation of structural-balance comparisons

We separately evaluated the proposed posterior procedure for comparing structural balance across two systems. The purpose of this experiment is to determine whether comparisons on the common B-scale remain meaningful when the systems being compared differ substantially in both population size and schedule density.

The first design was a matched comparison in which both systems contained 30 teams and used the same NBA schedule with 1,230 comparisons. The second was a heterogeneous comparison in which system 1 used the NBA design ($n_1 = 30, M_1 = 1230$), while system 2 used the NFL design ($n_2 = 32, M_2 = 272$). The heterogeneous setting is particularly relevant to the empirical analysis because it separates differences in the underlying balance parameter from differences in the amount of information available to estimate it.

For every replicate, system 1 was generated with $(B_1, T_1) = (0.5, 3)$, while system 2 was generated with $(B_2, T_2) = (0.2, 4)$. Because $B_1 > B_2$, system 1 is more structurally balanced under the data-generating model. We generated 100 replicates for each design and fitted the same MFM-BT

model independently to the two systems. Posterior draws were used to calculate

$$\Delta_B = B_1 - B_2, \qquad \Delta_d = d(B_2) - d(B_1), \qquad d(B) = 2\tau_0^2 \frac{1-B}{B}.$$

Positive values of either contrast favor system 1. We summarized the directional probability $\Pr(B_1 > B_2 \mid Y)$, the corresponding probability on the dispersion scale, and the three practical-testing probabilities (π_-, π_0, π_+). The practical-equivalence margin was set to $\epsilon_d = 0.4$ on the dispersion scale.

The results are summarized in Table 3. For the matched design, the average posterior probability that system 1 was more balanced was 0.864, compared with 0.829 for the heterogeneous NBA–NFL comparison. The corresponding probabilities that system 1 was more balanced by more than the practical-equivalence margin were 0.836 and 0.795. The posterior median dispersion contrasts were 3.46 and 3.15, respectively. Hence, the procedure identifies the correct direction of the balance difference under both comparison designs.

Table 3: Validation of the Bayesian structural-balance comparison over 100 replicates.

Design	$\Pr(B_1 > B_2)$	$\Pr(\Delta_d > 0)$	π_-	π_0	π_+	$BF_{01}^{(\epsilon)}$	Median Δ_d
Matched NBA–NBA	0.864	0.864	0.113	0.050	0.836	0.317	3.46
Heterogeneous NBA–NFL	0.829	0.829	0.143	0.062	0.795	0.397	3.15

The contrast between the two designs is itself informative. Swapping the NBA schedule for the much sparser NFL schedule does not flip the inferred ordering of the two systems, but it does weaken the posterior evidence. The average probability assigned to the correct practical direction falls from 0.836 to 0.795, while the mass given to the opposite direction and to the equivalence region grows. Schedule density, in other words, changes how confident the comparison is, not what it concludes on the common structural-balance scale.

The interval Bayes factors average 0.317 under the matched design and 0.397 under the heterogeneous one. Both sit below one, so the posterior does not favor practical equivalence relative to the prior in this directionally separated experiment, though neither comes close to certainty. That is about what one should expect from a single season, especially when one system supplies only 272 comparisons. It also explains why posterior probabilities serve the purpose better than a flat declaration that one league is or is not more balanced.

In sum, the second simulation supports the framework's main comparative claim. Structural balance can be compared across systems with different numbers of teams, tier configurations, and comparison graphs, because B is defined independently of all of these. Schedule differences surface instead through posterior uncertainty. This is precisely the situation the sports application presents: two leagues can be placed on the same structural scale even though one of them offers far more information about its latent competitive structure.

6 Comparing Competitive Balance and Tier Structure in the NBA and NFL

We apply the proposed MFM–BT model to the 2025–2026 National Basketball Association (NBA) regular season and the 2025 National Football League (NFL) regular season. Each game enters the analysis as a paired comparison between the home and away teams, and each league has its own home-advantage parameter on the common reference scale $\tau_0 = 1$. NFL postseason games are excluded, as is the additional Knicks–Spurs special event in the NBA.

For each league, we run two independently initialized MCMC chains. Each chain contains 4,000 iterations, with the first 1,000 iterations discarded as burn-in and every third subsequent iteration retained. This yields 1,000 posterior draws for posterior summaries. We use $\lambda_K = 1$ and $\gamma = 0.4$ for the MFM prior, and assess convergence using trace plots, running posterior means, and Gelman–Rubin diagnostics for B and the home-advantage parameter.

The two schedules differ sharply in density. The NBA season has each of its 30 teams play 82 games, giving 1,230 regular-season comparisons in total. The NFL, with 32 teams, plays only 17 games apiece, for 272 comparisons. The NBA therefore provides more than four times as many pairwise observations on relative team strength, while the NFL season is short enough that a single game carries considerable weight. This contrast between a long, dense schedule and a short, sparse one makes the pairing well suited to studying how schedule information affects what can be learned about competitive structure.

6.1 Structural balance and cross-league comparison

Posterior summaries for the two leagues appear in Table 4. The posterior mean of B is 0.482 for the NBA and 0.473 for the NFL. The NBA estimate is nominally higher, but the difference is small relative to posterior uncertainty: the 90% credible intervals are $[0.240, 0.681]$ and $[0.205, 0.693]$, respectively. The posterior mean of the home-advantage parameter is 0.274 for the NBA and 0.196 for the NFL. The corresponding 90% credible intervals are $[0.172, 0.381]$ and $[-0.030, 0.423]$. Thus, the NBA shows a larger estimated home advantage, although the intervals overlap substantially and the NFL interval includes zero.

Table 4: Posterior summaries of structural balance, home advantage, and latent tier structure for the NBA and NFL.

League	n	M	$\mathbb{E}(B \mid Y)$	90% CrI for B	$\mathbb{E}(h \mid Y)$	90% CrI for h	Dahl T
NBA	30	1,230	0.482	[0.240, 0.681]	0.274	[0.172, 0.381]	3
NFL	32	272	0.473	[0.205, 0.693]	0.196	[-0.030, 0.423]	2

To compare the leagues formally, we use the dispersion-scale procedure from Section 4. Define

$$\Delta_d = d(B_{\text{NFL}}) - d(B_{\text{NBA}}),$$

so positive values indicate greater structural balance in the NBA. We calibrate the practical margin by allowing a 0.25-unit difference in the RMS tier separation

$$s(B) = \sqrt{d(B)}$$

around $B_0 = 0.5$, which with $\tau_0 = 1$ gives the conservative dispersion-scale margin $\epsilon_d = 0.77$.

The posterior median of Δ_d is 0.04, with a 90% credible interval of $[-3.91, 5.49]$. The posterior probability that the NBA is more structurally balanced is

$$\Pr(\Delta_d > 0 \mid Y) = 0.510,$$

essentially a coin flip. Posterior mass spreads across all three practical regions,

$$(\pi_-, \pi_0, \pi_+) = (0.315, 0.340, 0.345),$$

rather than concentrating on a meaningful difference in either direction, and the interval Bayes factor $BF_{01} = 1.42$ offers only modest evidence for practical similarity relative to the prior. The

Table 5: Bayesian comparison of NBA and NFL structural balance on the dispersion scale.

Median Δ_d	90% CrI	$\Pr(\Delta_d > 0)$	π_-	π_0	π_+	BF_{01}
0.04	[-3.91,5.49]	0.510	0.315	0.340	0.345	1.42

data, in other words, support neither a directional difference in structural balance nor strong practical equivalence; one season simply does not contain enough information for the latter.

The comparison also highlights a useful property of the balance measure. The NBA supplies more than four times as many regular-season comparisons as the NFL, yet this does not change what B means, only how well it is estimated. The NFL interval is noticeably wider at a similar posterior mean, which is exactly what one expects when less data are available to separate persistent strength differences from game-level noise. A league should therefore not be called more or less balanced merely because its schedule happens to produce a sharper or noisier standings table.

6.2 Comparison with traditional competitive-balance measures

For context, we also compute conventional competitive-balance measures used in sports economics from regular-season win percentages, including the standard deviation, top-to-bottom range, Herfindahl–Hirschman index (HHI), and Gini coefficient (Fort and Maxcy, 2003; Utt and Fort, 2002; Owen et al., 2007; Manasis et al., 2013). Lower values indicate greater observed balance. These measures summarize realized standings and therefore address a different question from the structural balance parameter. In particular, they do not separate persistent differences in underlying team strength from game-level variation, and they do not quantify uncertainty in the latent tier structure.

Table 6: Traditional competitive-balance measures calculated from regular-season standings.

League	SD of win percentage	Range	HHI	Gini
NBA	0.169	0.573	0.0370	0.188
NFL	0.204	0.647	0.0363	0.230

The conventional measures mostly point toward greater observed parity in the NBA (Table 6). The standard deviation of win percentages is 0.169 in the NBA against 0.204 in the NFL, and the ranges are 0.573 and 0.647. The Gini coefficients, 0.188 and 0.230, tell the same story. Only the HHI departs slightly, at 0.0370 for the NBA and 0.0363 for the NFL. The conventional toolkit thus does not even agree with itself about the ranking, though most of its members see more dispersion in the NFL standings.

The model-based analysis cuts across this. Despite the wider spread of observed NFL win percentages under most conventional measures, the posterior distributions of B give little reason to think the leagues differ in underlying structural balance. Some of what the standings show as an NFL disadvantage is plausibly the product of a much shorter schedule and the game-level variation that comes with it, rather than genuinely larger persistent heterogeneity. Observed parity and latent structural balance, it seems, need not tell the same story.

6.3 Latent competitive tiers and team rankings

The structural balance parameter measures overall heterogeneity, but it does not describe how teams are organized within a league. We therefore also examine the estimated competitive tiers. Because mixture labels are arbitrary, we order the tiers according to the Dahl-selected tier abilities, with Tier 1 denoting the strongest competitive tier. The Dahl representative partition identifies

three tiers in the NBA with 6, 14, and 10 teams, respectively, and two tiers in the NFL with 16 teams each (Table 7).

Table 7: MFM–BT Dahl representative tier composition for the NBA and NFL. Tier 1 denotes the strongest competitive tier.

League and tier	Teams
NBA Tier 1 (higher)	Boston Celtics; Denver Nuggets; Detroit Pistons; New York Knicks; Oklahoma City Thunder; San Antonio Spurs.
NBA Tier 2 (middle)	Atlanta Hawks; Charlotte Hornets; Cleveland Cavaliers; Golden State Warriors; Houston Rockets; LA Clippers; Los Angeles Lakers; Miami Heat; Minnesota Timberwolves; Orlando Magic; Philadelphia 76ers; Phoenix Suns; Portland Trail Blazers; Toronto Raptors.
NBA Tier 3 (lower)	Brooklyn Nets; Chicago Bulls; Dallas Mavericks; Indiana Pacers; Memphis Grizzlies; Milwaukee Bucks; New Orleans Pelicans; Sacramento Kings; Utah Jazz; Washington Wizards.
NFL Tier 1 (higher)	Buffalo Bills; Chicago Bears; Denver Broncos; Detroit Lions; Green Bay Packers; Houston Texans; Indianapolis Colts; Jacksonville Jaguars; Los Angeles Chargers; Los Angeles Rams; Minnesota Vikings; New England Patriots; Philadelphia Eagles; Pittsburgh Steelers; San Francisco 49ers; Seattle Seahawks.
NFL Tier 2 (lower)	Arizona Cardinals; Atlanta Falcons; Baltimore Ravens; Carolina Panthers; Cincinnati Bengals; Cleveland Browns; Dallas Cowboys; Kansas City Chiefs; Las Vegas Raiders; Miami Dolphins; New Orleans Saints; New York Giants; New York Jets; Tampa Bay Buccaneers; Tennessee Titans; Washington Commanders.

The NBA shows a more differentiated competitive structure. Six teams make up Tier 1, a large middle group of 14 sits in Tier 2, and 10 teams fall to Tier 3. Much of the league, in other words, is packed into the middle, with a relatively small group pulling away at the top. This is also where the tier structure and the global balance parameter part company: a league can be highly balanced overall and still contain a recognizable group of leading teams. The tier compositions also say things the raw standings do not. Tier 1 puts the defending champion Oklahoma City Thunder and the usual contenders, Boston and Denver, alongside two teams a standings-only reading would overlook. Detroit spent much of the season playing better than its win total suggested, and San Antonio belonged to the elite group on latent strength well before its record caught up. The Knicks sit in that group too, which shows the model looking past short streaks to how teams actually compare. Working the other way, a few famous names, the Lakers and Warriors among them, land in the broad middle tier rather than the elite one, and a win-percentage table flattens that distinction into one long ranking. The practical reading for a front office is simple: about half the league is a middle band where games are close to a coin flip, and a six-team group forms the realistic championship tier. That distinction shapes how a season should be judged, and no standings table makes it visible.

The NFL partition is coarser, splitting the league evenly into two tiers of 16. That should not be read as a simpler competitive structure. With only 272 regular-season games against the NBA's 1,230, the data simply carry far less information about teams of similar ability. The broad NFL tiers reflect both the organization the data support and the heavier uncertainty that a sparse schedule brings. The NFL tiers carry the same kind of information. Tier 1 contains the eventual Super

Bowl champion Seattle and the AFC champion New England, along with teams whose seasons ended earlier than their strength deserved, Buffalo and the Chargers being the clearest cases. It also holds the season's surprise division winners, including Chicago, whose arrival in the top group the model caught even though the franchise had spent years near the bottom. Tier 2, meanwhile, collects teams whose reputations run ahead of their present strength. Kansas City and Baltimore both fall below the midpoint on latent ability despite championship pedigrees that say otherwise. In a seventeen-game season, a couple of close losses or one injury at the wrong time can push a team like that out of January, and the tiers capture the fact that they never separated from the league's middle. The honest summary of an NFL season, it turns out, is a large top group of broadly comparable clubs, a handful of genuine stragglers, and a lot of doubt in between, not the clean thirty-two-step ordering the standings imply.

The comparison also shows why the number of tiers should not be used as a measure of overall competitive balance. The NBA has three estimated tiers and the NFL has two, while their posterior estimates of B are very similar. The tier structure describes how teams are organized into competitive levels, whereas B measures the magnitude of heterogeneity among those levels. Therefore, more tiers do not necessarily imply less competitive balance.

Figure 2 provides a more detailed view of the posterior ranking and clustering results. The rank intervals describe uncertainty in the relative positions of individual teams, while the posterior mean co-clustering matrices summarize uncertainty in whether pairs of teams belong to the same latent competitive tier.

The posterior mean co-clustering matrices provide information that is not available from a single representative partition. Teams near a tier boundary may have lower posterior co-clustering probabilities with teams in their assigned tier and nonnegligible probabilities of clustering with teams in a neighboring tier. The Dahl tiers should therefore be interpreted as representative summaries of posterior clustering uncertainty rather than as fixed classifications of team quality.

6.4 Agreement with observed standings and practical implications

As a descriptive check, we compare posterior median ranks with observed regular-season ranks based on winning percentage, using point differential as a transparent tie-breaker. Since both rankings come from the same games, this measures agreement with the observed season rather than out-of-sample prediction.

Table 8: Agreement between MFM–BT posterior ranks and observed regular-season ranks.

League	Spearman ρ	Kendall τ	Pearson r	Mean absolute rank error	Top-five overlap
NBA	0.978	0.924	0.949	2.35	5/5
NFL	0.990	0.944	0.959	2.38	4/5

The model ranks teams much the way the standings do. The Spearman correlation is 0.978 for the NBA and 0.990 for the NFL, and the Kendall correlations are 0.924 and 0.944. The model's top five also matches the observed NBA top five exactly, and four of the observed NFL top five. So the model keeps the ordering we see in the season results, but adds something the standings cannot: uncertainty intervals and information about which teams belong to the same competitive group.

The results leave us with a few observations. First, the usual measures cannot even settle which league is more balanced. Most of the dispersion measures point to the NBA, the HHI points slightly to the NFL, and our model finds no meaningful difference in structural balance. Much depends on what one chooses to look at: the results on the court and field, or the strength of the

teams underneath them. It also becomes clear that equal balance does not mean the same kind of competition. The NBA breaks into three tiers, with most clubs crowded into a broad middle and a small elite group above them, while the NFL settles into two wide groups. A policy aimed at parity might close the gaps between teams without changing how they bunch together, or it might reshuffle the groups without narrowing the gaps, and it is worth knowing which of the two a given policy actually does. Finally, the schedule puts a ceiling on what any of this can tell us. An NBA season simply contains far more information about how teams compare, whereas an NFL season leaves real doubt about teams of similar strength and about where one tier gives way to the next. A comparison built only on standings risks mistaking that doubt for a genuine difference in balance. On the whole, the NBA and NFL look much alike in global structural balance but quite different in how their competition is organized, and in how confidently that organization can be described. Balance, tier structure, and schedule information are separate features of a league. Treating them as such makes for better comparisons than leaning on standings, conventional indices, or tier counts alone.

7 Discussion and Conclusion

We have developed a Bayesian paired-comparison framework for studying competitive balance that separates global structural heterogeneity from latent competitive tier structure. The MFM prior learns the number and membership of occupied tiers, while the balance parameter B quantifies how strongly those tiers are separated. Our theoretical results give B a direct interpretation in terms of between-tier dispersion and show how overall team-level heterogeneity depends jointly on the size of tier separation and the prevalence of cross-tier comparisons. Because the framework separates the underlying competitive structure from the information contained in the observed schedule, balance can be compared across systems with different numbers of teams and games. The simulations show that the method recovers latent competitive structure accurately when the comparison design is informative enough, and that estimates of structural balance remain comparatively stable even as schedules thin out. The NBA and NFL application reinforces the distinction between global balance and competitive stratification. Conventional standings-based measures generally suggest greater observed parity in the NBA, yet the posterior analysis finds little evidence of a meaningful difference in underlying structural balance. The two leagues are organized differently all the same: the NBA splits into three latent competitive tiers, while the NFL falls into two broader ones, and the greater uncertainty in the NFL reflects its much sparser schedule.

Several extensions are worth exploring. A longitudinal version of the model could let B, team abilities, and tier memberships drift across seasons, which would make it possible to characterize changes in parity, movement between competitive tiers, and the effects of league policies on competitive structure. On the theory side, it would be useful to understand how the topology of the comparison graph shapes estimation. Beyond plain connectivity, properties such as graph density, degree heterogeneity, spectral connectivity, and bottlenecks may govern how precisely abilities, tiers, and structural balance can be learned, and could inform the design of schedules that carry more information. One further direction is to relax the transitive structure of the Bradley–Terry model. Sports competitions often show systematic matchup patterns: team A tends to beat team B, team B tends to beat team C, but team C tends to beat team A. A one-dimensional latent ability cannot represent such cycles. A natural next step is to augment the global ability component with structured matchup effects that capture nontransitive patterns while keeping measures of overall strength and balance interpretable. That would give a richer account of both hierarchical and cyclic structure in competitive systems.

References

Basini, F., Tsouli, V., Ntzoufras, I., and Friel, N. (2023). Assessing competitive balance in the English Premier League for over forty seasons using a stochastic block model. *Journal of the Royal Statistical Society Series A: Statistics in Society*, 186(3):530–556.

Bradley, R. A. and Terry, M. E. (1952). Rank analysis of incomplete block designs: I. the method of paired comparisons. *Biometrika*, 39(3/4):324–345.

Busse, L. M., Orbanz, P., and Buhmann, J. M. (2007). Cluster analysis of heterogeneous rank data. In *Proceedings of the 24th international conference on Machine learning*, pages 113–120.

Caron, F. and Doucet, A. (2012). Efficient bayesian inference for generalized bradley–terry models. *Journal of Computational and Graphical Statistics*, 21(1):174–196.

Dahl, D. B. (2006). Model-based clustering for expression data via a dirichlet process mixture model. In Do, K.-A., Müller, P., and Vannucci, M., editors, *Bayesian Inference for Gene Expression and Proteomics*, pages 201–218. Cambridge University Press.

Davidson, R. R. (1970). On extending the bradley-terry model to accommodate ties in paired comparison experiments. *Journal of the American Statistical Association*, 65(329):317–328.

Deng, K., Han, S., Li, K. J., and Liu, J. S. (2014). Bayesian aggregation of order-based rank data. *Journal of the American Statistical Association*, 109(507):1023–1039.

Dong, P., Han, R., Jiang, B., and Xu, Y. (2026). Statistical ranking with dynamic covariates. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 88(1):221–238.

Dunson, D. B. and Peddada, S. D. (2008). Bayesian nonparametric inference on stochastic ordering. *Biometrika*, 95(4):859–874.

Fort, R. and Maxcy, J. (2003). Competitive balance in sports leagues: An introduction. *Journal of Sports Economics*, 4(2):154–160.

Fort, R. and Quirk, J. (1995). Cross-subsidization, incentives, and outcomes in professional team sports leagues. *Journal of Economic Literature*, 33(3):1265–1299.

Glickman, M. E. (1999). Parameter estimation in large dynamic paired comparison experiments. *Journal of the Royal Statistical Society Series C: Applied Statistics*, 48(3):377–394.

Hubert, L. and Arabie, P. (1985). Comparing partitions. *Journal of Classification*, 2(1):193–218.

Hunter, D. R. (2004). MM algorithms for generalized Bradley-Terry models. *The Annals of Statistics*, 32(1):384–406.

Kulis, B. and Jordan, M. I. (2012). Revisiting K-means: New algorithms via bayesian nonparametrics. In *Proceedings of the 29th International Conference on Machine Learning*, pages 1131–1138.

Li, X., Yi, D., and Liu, J. S. (2022). Bayesian analysis of rank data with covariates and heterogeneous rankers. *Statistical Science*, 37(1):1–23.

Luce, R. D. (1959). *Individual Choice Behavior: A Theoretical Analysis*. Wiley.

Manasis, V., Avgerinou, V., Ntzoufras, I., and Reade, J. J. (2013). Quantification of competitive balance in European football: Development of specially designed indices. *IMA Journal of Management Mathematics*, 24(3):363–375.

Manasis, V. and Ntzoufras, I. (2014). Between-seasons competitive balance in European football: Review of existing and development of specially designed indices. *Journal of Quantitative Analysis in Sports*, 10(2):139–152.

Manasis, V., Ntzoufras, I., and Reade, J. J. (2022). Competitive balance measures and the uncertainty of outcome hypothesis in european football. *IMA Journal of Management Mathematics*, 33(1):19–52.

Marden, J. I. (1996). *Analyzing and modeling rank data*. CRC Press.

Masarotto, G. and Varin, C. (2012). The ranking lasso and its application to sport tournaments. *The Annals of Applied Statistics*, 6(4):1949–1970.

McLachlan, G. J. and Peel, D. (2000). *Finite Mixture Models*. Wiley Series in Probability and Statistics. John Wiley & Sons, New York.

Miller, J. W. and Harrison, M. T. (2018). Mixture models with a prior on the number of components. *Journal of the American Statistical Association*, 113(521):340–356.

Mollica, C. and Tardella, L. (2021). Bayesian analysis of ranking data with the extended plackett–luce model. *Statistical Methods & Applications*, 30(1):175–194.

Murphy, T. B. and Martin, D. (2003). Mixtures of distance-based models for ranking data. *Computational statistics & data analysis*, 41(3-4):645–655.

Neal, R. M. (2000). Markov chain sampling methods for Dirichlet process mixture models. *Journal of Computational and Graphical Statistics*, 9(2):249–265.

Owen, P. D., Ryan, M., and Weatherston, C. R. (2007). Measuring competitive balance in professional team sports using the herfindahl–hirschman index. *Review of Industrial Organization*, 31(4):289–302.

Pearce, M. and Erosheva, E. A. (2025a). Bayesian rank-clustering. *Psychometrika*, 90(3):904–931.

Pearce, M. and Erosheva, E. A. (2025b). Modeling preferences: A bayesian mixture of finite mixtures for rankings and ratings. *Journal of the American Statistical Association*, 120(551):1621–1632.

Polson, N. G., Scott, J. G., and Windle, J. (2013). Bayesian inference for logistic models using pólya–gamma latent variables. *Journal of the American Statistical Association*, 108(504):1339–1349.

Santi, L. and Friel, N. (2026). The bradley–terry stochastic block model. *The Annals of Applied Statistics*, 20(2):1766–1787.

Szymanski, S. (2001). Income inequality, competitive balance and the attractiveness of team sports. *Economic Journal*, 111(469):69–84.

Szymanski, S. (2003). The economic design of sporting contests. *Journal of Economic Literature*, 41(4):1137–1187.

Tian, X., Shi, J., Shen, X., and Song, K. (2024). A spectral approach for the dynamic bradley–terry model. *Stat*, 13(3):e722.

Utt, J. and Fort, R. (2002). Pitfalls to measuring competitive balance with gini coefficients. *Journal of Sports Economics*, 3(4):367–373.

Vrooman, J. (1995). A general theory of professional sports leagues. *Southern Economic Journal*, 61(4):971–990.

Zimbalist, A. (2002). Competitive balance in sports leagues: An introduction. *Journal of Sports Economics*, 3(2):111–121.




Figure 2: Posterior team rankings, Dahl representative tiers, and posterior mean co-clustering matrices for the NBA and NFL. Circles denote posterior median ranks, triangles denote ranks from the Dahl representative partition, and horizontal bars represent 90% posterior credible intervals for team ranks. Team colors indicate the Dahl representative tiers, with Tier 1 denoting the strongest competitive tier. The heatmaps display the posterior mean co-clustering matrices, where each entry is the posterior mean of the indicator that two teams belong to the same latent tier. Rank 1 denotes the strongest team.