

Geographical Data and Computational Tools for Determining Restoration Priorities in Mangrove Ecosystems

Ainur Smailova, Laura Valentina Sierra, Pablo Ramos-Henao, Saif Ur Rehman, Yan-Yu Lin

* Department of Computer Science, University of Liverpool

Abstract

In this review article

Introduction

Mangrove Forest Ecosystems

Mangrove forests are unique, highly productive coastal ecosystems characterized by salt-tolerant trees and shrubs that thrive in the harsh intertidal zones. These plant species are also named mangroves, given that they are the dominant species of their environment. The areas between land and sea where conditions fluctuate widely in salinity and water level, which creates complex niches for the distinct species that can tap into the vast resources that these areas possess; these forests are vital for global ecological balance, providing critical biodiversity services and supporting complex food webs [1].

A primary function of mangroves is their significant role in global climate change mitigation. They possess an exceptional capacity for carbon sequestration, storing up four times more carbon than comparable terrestrial forests [2]. Furthermore, ecologically, mangroves serve as natural coastal defense barriers, providing protection against tidal erosion and storm surges, while also functioning as critical nurseries and habitats for numerous species of fish, crustaceans, and birds; thus supporting coastal communities worldwide.

Despite these benefits, mangrove forests are facing rapid decline: deforestation is primarily driven by anthropogenic pressures such as urbanization, aquaculture expansion, and agricultural development, leading to reported loss rates of 1 to 2 percent of the total area per annum [2]. The complexity of developing effective preservation strategies is amplified by the diversity of mangrove species and the vast range of global climate conditions they inhabit, necessitating comprehensive, globally available datasets and advanced analytical tools to determine where particular species may or may not thrive. Having

access to sound ecological modelling of species distributions allows for more effective restoration efforts, I.E. planting the right mangrove species after logging or climate disasters. Additionally, real time updates on forest coverage can issue warnings to NGOs and local authorities, which lead to faster, more effective action [1].

To address the challenge of global mangrove monitoring, the Global Mangrove Watch (GMW) platform provides open access to remote-sensing data and analytical instruments. This platform delivers near real-time information on mangrove extent and global change dynamics Global Mangrove Watch. A key component of GMW is the mangrove loss alert system, which utilizes Copernicus Sentinel-2 satellite data processed at a 20-meter resolution. While Sentinel-2 typically provides weekly image acquisitions, the alert process is subject to delays caused by cloud cover and the requirement for multiple observations to confirm genuine ecosystem change. Consequently, an alert is typically transmitted approximately three months after a loss event occurs [1].

Despite having modern technology and open data access, achieving global conservation goals for coastal ecosystems has been difficult because consistent, worldwide data on their past and present health is lacking. Now, access to resources like the decades-old Landsat satellite archive and newer, higher-resolution satellites allows us to analyze changes over long periods. This high-resolution data helps us accurately map scattered or fragmented ecosystems like mangrove forests. This new data is combined with powerful computing systems and cloud platforms (e.g., Google Earth Engine), allowing us to quickly process satellite data for the entire planet. This combination has encouraged many global-scale studies that can help answer critical questions for mangrove conservation. Still it is important to note that these data sets notably merge data from different countries and institutions, which is often patchy, has different resolutions, and may be incomplete. As seen in the Global Mangrove Watch official documentation, data fusion is needed to some extent, depending on the

kind of analysis one may wish to run. It will become evident when the results of the referenced studies are discussed, this poses challenges when running machine learning algorithms, and when interpreting the results from a holistic and ecological perspective.

Aim

The aim of this paper is to assess the status of the technologies and analysis techniques used in determining the conservation status and restoration needs of mangrove forests; as well as identifying research gaps.

Objectives

The first objective is to review and assess the current completeness and robustness of various available datasets. Several factors can affect data access and analysis, such as cloud cover and shadows when comparing satellite data. These limitations can hinder the consistency of images required for continuous change monitoring. [3] Secondly, we aim to identify the technologies used for mangrove identification and conservation, as well as the computational modules used to analyse these datasets and gauge their performance. Additionally, we aim to outline the current challenges of measurement tools and computational analysis methods to elaborate on the main existing research gaps.[4]

Methodology

Article Selection Process

In the first stage of this scoping review, records from four major databases were retrieved. The initial search produced a combined set of studies, with the number of records obtained from each source presented in prisma chart (Figure X).

During screening, we identified a number of non-English records and further duplicates. Because our institution offers full-text access to all searched databases, none of the records were excluded due to retrieval limitations.

The remaining studies were then divided among the research team, and each member screened titles and abstracts to assess their relevance to the research objectives. Studies were excluded if they did not involve geospatial data, if they relied solely on simple descriptive or statistical analyses, or if they did not use the modelling approaches central to our review, such as Random Forest or CNN-based methods, they fell outside the scope of this study.

Findings

This is the section for findings

Geological Survey Tools

This is the section for geological survey tools

Computational Analysis Techniques

This is the section for computational analysis techniques

Research Gaps

«add more here»>

Convolutional Neural Networks (CNNs) are the most widely used models in the field and are the foundation of most mangrove classification approaches. However, they struggle to detect small and fragmented mangroves resulting in incorrect classifications. [5]

Conclusion

References

- [1] Maricé Leal and Mark D. Spalding. *The State of the World's Mangroves 2024*. Global Mangrove Alliance, July 2024. DOI: 10.5479/10088/119867. URL: <https://repository.si.edu/handle/10088/119867> (visited on 11/24/2025).
- [2] Thomas A. Worthington et al. "Harnessing Big Data to Support the Conservation and Rehabilitation of Mangrove Forests Globally". In: *One Earth* 2.5 (May 22, 2020), pp. 429–443. ISSN: 2590-3322. DOI: 10.1016/j.oneear.2020.04.018. URL: <https://www.sciencedirect.com/science/article/pii/S2590332220302050> (visited on 10/29/2025).
- [3] Uday Pimple et al. "Enhancing monitoring of mangrove spatiotemporal tree diversity and distribution patterns". In: *Land Degradation & Development* 34.5 (Mar. 1, 2023), pp. 1265–1282. ISSN: 1085-3278. DOI: 10.1002/ldr.4537. URL: <https://research.ebsco.com/linkprocessor/plink?id=1ac97efb-b02e-32b6-8ca3-f1f271c277bc> (visited on 11/25/2025).

- [4] Qiaosi Li, Frankie Kwan Kit Wong, and Tung Fung. "Mapping multi-layered mangroves from multispectral, hyperspectral, and LiDAR data". In: *Remote Sensing of Environment* 258 (June 2021), p. 112403. ISSN: 00344257. DOI: 10.1016/j.rse.2021.112403. URL: <https://linkinghub.elsevier.com/retrieve/pii/S0034425721001218> (visited on 11/18/2025).
- [5] Ruoxin Zhang et al. "FragMangro: A cross-domain zero-shot model for monitoring fragmented mangrove ecosystems". In: *Journal of King Saud University Computer and Information Sciences* 37.4 (May 20, 2025), p. 46. ISSN: 2213-1248. DOI: 10.1007/s44443-025-00053-y. URL: <https://doi.org/10.1007/s44443-025-00053-y> (visited on 11/18/2025).