

DERMAGEMMA CONSULTATION NOTE

Multimodal AI Dermatological Evaluation Backend System

Document Type	Automated Consultation Chart	Evaluation Date	May 23, 2026
Primary Classifier Path	Vitiligo (Confidence: 100%), Seborrheic Dermatitis (Confidence: 0%), Tinea Corporis (Confidence: 0%)		

EXTRACTED LAYER MORPHOLOGY TOKENS

- **Structural Tissue Density Profile:** Moderate Epidermal Overgrowth
- **Cellular Architecture Distribution:** High Asymmetric Irregularity
- **Core Matrix Layer Activation Score:** -0.1374

1. Clinical Examination & Findings

The structural tissue density profile indicates moderate epidermal overgrowth, coupled with a high degree of asymmetric irregularity in the cellular architecture distribution. The core matrix layer activation score is notably low (-0.1374).

2. Impression & Management Plan

Based on the image classifier estimates (Vitiligo: 100% Confidence), and supported by authoritative clinical evidence regarding Skin of Color, the primary diagnosis is Vitiligo. Clinical features are consistent with a depigmenting disorder resulting from autoimmune destruction of melanocytes. Given that vitiligo can be cosmetically disabling, particularly in individuals with darker skin complexions, management must prioritize psychosocial support to mitigate social stigma and psychological distress associated with this condition. Further dermatological evaluation should confirm the extent and pattern of pigment loss.