



# FÖRSÄTTSBLAD TENTAMEN/ EXAMINATION COVER

*Jag intygar att mobiltelefon och annan otillåten elektronisk utrustning är avstängd och förvaras på anvisad plats. / I hereby confirm that mobile phones and other unauthorized electronic equipment is shut off and placed according to instructions*

MARKERA MED "X"/  
MARK WITH "X"

IFYLLES AV STUDENT OCH TENTAMENSVAKT/  
TO BE FILLED IN BY THE STUDENT AND THE INVIGILATOR:

KURSKOD / COURSE CODE	D D 2 4 2 3	EFTERNAMN / FAMILY NAME	FILOTHEOU																
KURSNAMN / COURSE NAME	Bildbehandling och datorseende																		
PROVKOD / TEST CODE	T E N 1																		
TENTAMENSDATUM / EXAMINATION DATE																			
Y/Y/Y/Y	M/M	D/D																	
2 0 1 6	- 0 1	- 1 5																	
PROGRAMKOD / PROGRAM CODE: <i>5CR</i>	INLÄMNINGSTID / TIME SUBMITTED: <i>1807</i>	SIGNATUR TENTAMENSVAKT / SIGNATURE INVIGILATOR: <i>SD</i>	ANTAL SIDOR / NO OF PAGES: <i>27</i>																
MARKERA BEHANDLADE UPPGIFTER MED "X" OCH EJ BEHANDLADE UPPGIFTER MED "-" / MARK WITH "X" PROBLEMS SOLVED. MARK WITH "-" PROBLEMS NOT ATTEMPTED																			
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
X	X	X	X	-	X														

IFYLLES AV INSTITUTIONEN / TO BE FILLED IN BY THE DEPARTMENT:

BEDÖMNING / ASSESSMENT																				
A r2 ≥ r2	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
	4	6	4.5	4	-	5	+ 5	= 28.5												

BONUSPOÄNG/  
BONUS POINTS:

SLUTSUMMA /  
FINAL POINTS:

BETYG/  
GRADE:


0406186500

Godkänts av examiner /  
approved by Examiner.....

*Mårten Björn*

1)

object



Room with  
no light  
coming in  
except from  
the pinhole

Light comes in through an infinitesimally small hole in one wall. The object is projected to the opposite wall, scaled and inverted both horizontally and vertically, due to the intersection of the light rays with it.

It is not possible for a real-pinhole camera to be manufactured due to the infinitesimality of the hole.



FILOTHEOU

ALEXANDROS

871108-5590 SCR

2 A2

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

2) Edge detection can help shape features, or produce, or verify segmentation methods. ~~that~~. Also, they denote ~~a~~ the boundaries of objects, hence they are valuable ~~for semantic~~ if we want to produce semantically rich results through computer vision or ~~in~~ image analysis.



FILÓTHEOU ALEXANDROS 871108-5590 SZR

Family name, first name

Personal Registration Number

Programme

3

A.3.

Sheet no.

Problem no.

3) We can represent points located at infinity.

If  $(kx, ky, kz, k)$  are the homogeneous coordinates of a point, then if  $k=0$ , the point is considered to be at infinity. ( $(x, y, z)$  are its coordinates in the Cartesian system).



FN 07 THEOU ALEXANDROS 871108-5590 SZR

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

A A4

4) The magnitude holds ~~the~~ the information that is comprehensible to humans via their vision system. The phase has no ~~an~~ immediate visual value to humans, although I found a study where the researchers used the ~~the~~ magnitude of one image ~~and~~ ~~con~~ in combination with the phase of another, and the combined "reconstructed" image was visually similar to the image from where the phase was taken.

The phase component holds the information from where each sinusoid that makes up the signal is bound to start.



FILOTHEOU ALEXANDRIS

871108-5590 SDR

5

A6

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

6) ~~Q~~ The camera sensor ~~must be~~ has to be somehow mounted to the inside of the camera and ~~aligned with~~ its center, aligned with the center of the camera's lens. It is ~~usual~~ expected that the sensor may ~~be not~~ not be perfectly ~~para~~ aligned with the plane glued to, ~~it either~~ meaning it can be skewed ~~it either~~ the ~~Z~~ along either the Z-axis, rotated along the Z axis or its center may not be in line with the one of the lens. These types of imperfections ~~are~~ are tackled by the intrinsic camera parameters.



FILOTHEOU ALEXANDROS

871108-5590 SCR

6

A8

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

8) We assume that the image is one period of an infinitely repeated image

We assume that ~~any~~ image is the representation over one period of a periodic signal. This means that the periodic signal is the infinitely repeated x-wise and y-wise image.



FILOTHEOU

ALEXANDROS

871108-5590

SCR

7

A9

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

9) If  $W_{max}$  is the highest frequency present in an image, then aliasing occurs when we try to sample it at a sampling rate lower than  $2 \cdot W_{max}$ .



FILOTHEOU ALEXANDRIS 871108-5590 SCR 8 A10

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

10.) Gaussian filters are linear and shift invariant.



FILOTHEOU ALEXANDROS 871108-5590 SCR

9

All

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

II. ~~We compute the~~

First, we compute the x-wise and y-wise first order derivatives,  $f_x$  and  $f_y$  respectively. Then, the second moment matrix is formed by

$$\Sigma = \sum \begin{bmatrix} f_x^2 & f_x f_y \\ f_x f_y & f_y^2 \end{bmatrix}$$

that is, summing over all values in  $f_x$  squared,  $f_y$  squared and the product of the  $f_x$  and  $f_y$  for every corresponding image position.



FILOTHEOU ALEXANDROS

871108-5590 SZR

10 A19

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

12) It contains information about  
~~non features~~ & ~~sum of corners~~  
differences of Gaussian of feature  
sum at corner corners and combinations  
of features

-1p

Not, the feature descriptor.



FILΟΤΗΕΩΝ ΑΛΕΞΑΝΔΡΟΣ 871108-5590 SCR

Family name, first name

Personal Registration Number

Programme

11

A13

Sheet no.

Problem no.

13) KANSAC works through these 5 steps:

1. First, pick a minimal set of points
2. Fit a ~~line~~ (line in this case) model to those points.
3. Count the number of points that whose distance is lower than a set threshold.
4. Repeat steps 1-3 for a predefined number of iterations.
5. Choose the model ~~with~~ with the highest number of points in step 3.



FROTHEN

ALEXANDROS

Family name, first name

871108-5590

Personal Registration Number

Programme

19

A14

Sheet no.

Problem no.

14. Template-based methods are not robust because they look for a fixed template, or image, inside another one and ~~that is the latter~~ in order for it to be recognised it has to be in exactly the same dimension, orientation and brightness in it. There is little to no invariance to scale, point of view, occlusion, noise or illumination conditions with <sup>this</sup> ~~these~~ type of recognition method.



FILOTHEOU ALEXANDROS 871108-5590 SCR 13. A15

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

15) In binocular disparities it is easier because the cameras' ~~are~~ positions are fixed relatively to each other, and their baseline is known and fixed over time. However, in optical flow the structure of the scene might change along with the position of the camera, hence ~~are~~ there are two types of motions going on: egomotion and scene motion. For depth to be computed, or disparities, the relative translation and rotation for each type of motion needs to be computed, and depth cannot be computed without translation.


-S.

-TP

-T

-SP

12/15



FILOTHEOU ALEXANDRIS 871109-5590 SZR

14 B.I

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

1.) ⑩ For the first camera:

$$X_{1A} = P_A \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ 1 \end{bmatrix}$$

$$X_{2A} = P_A \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 2 \end{bmatrix} = \begin{bmatrix} 0,5 \\ 3 \\ 1 \end{bmatrix}$$

$$X_{3A} = P_A \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 7 \\ 2 \end{bmatrix} = \begin{bmatrix} 1,5 \\ 3,5 \\ 1 \end{bmatrix}$$

$$X_{4A} = P_A \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 1 \end{bmatrix} =$$

⑪ For the second camera,

$$X_{1B} = P_B \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ 1 \end{bmatrix}$$


$$X_{2B} = P_B \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 6 \\ 1 \end{bmatrix}$$

$$X_{3B} = P_B \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 7 \\ 1 \end{bmatrix}$$


$$X_{4B} = P_B \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 1 \end{bmatrix}$$

Where we have assumed homogeneous coordinates. Then in 2D:

► For the first camera:



► For the second camera:





FILOTHEOU ALEXANDROS 871108-5590 SIR.

16 B1

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

The second camera is the affine one, since parallel lines in 3D are maintained in their projection in 2D. Hence the first one is the perspective one.

④ By looking at the projection matrices, we can tell that the second one is the affine one, since its last row is all zeros except for the last element, which is one.



FILOTHEOU ALEXANDROS 871108-5590

SZR.

17

B.9

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

(circled)

1) Fourier transform:

$$\hat{f}(w) = \int_{-\infty}^{+\infty} f(x) e^{-jwx} dx = \int_{-\infty}^{+\infty} e^{-jwx} dx = \frac{-1}{jw} [e^{-jwx}]_{-\infty}^{+\infty}$$

$$\hat{f}(w) = \frac{-1}{jw} (e^{-\frac{jw}{2}} - e^{\frac{jw}{2}})^{-1/2} = \frac{2}{w} \cdot \frac{e^{\frac{jw}{2}} - e^{-\frac{jw}{2}}}{2j}$$

$$\hat{f}(w) = \frac{2}{w} \cdot \sin \frac{w}{2} = \frac{\sin \frac{w}{2}}{\frac{w}{2}}$$

~~Unfortunately, a box filter introduce~~  
Unfortunately, convolving an image with ~~it's~~ a box filter introduces ringing effects, and, hence, it degrades the quality of the image. This is due to the sinc's structure.



FLOTHEIM ALEXANDER

871108-5590

SIR

18

B.2

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

2). In 1D, a convolution of function  $f$  with function  $g$  is defined as:

$$f(x) * g(x) = \int_{-\infty}^{+\infty} f(a)g(x-a)da.$$

② In our case:

$$f(x) * f(x) = \int_{-\infty}^{+\infty} f(a)f(x-a)da$$

$$= \int_{-1/2}^{1/2} 1 \cdot f(x-a)da. \quad ①$$

$$f(a) = \begin{cases} 1, & -1/2 \leq a \leq 1/2 \\ 0, & \text{otherwise} \end{cases}$$

$$f(-a) = \begin{cases} 1, & -1/2 \leq -a \leq 1/2 \\ 0, & \text{otherwise} \end{cases}$$

$$f(-a) = \begin{cases} 1, & a \leq 1/2 \text{ and } a \geq -1/2 \\ 0, & \text{otherwise} \end{cases}$$

$$f(x-a) = \begin{cases} 1, & x-1/2 \leq x-a \leq x+1/2 \\ 0, & \text{otherwise.} \end{cases}$$



FILMTHEATER ALEXANDER

Family name, first name

871108-5590

Personal Registration Number

SUR

Programme

19

B2

Sheet no.


Problem no.

or, substitute  $x-a=y$

then:  $da = -dy$

$$a = -\frac{1}{2} \rightarrow y = x + \frac{1}{2}$$

$$a = \frac{1}{2} \rightarrow y = x - \frac{1}{2}$$




-1p

and ①:

$$x = \frac{1}{2} \quad x + \frac{1}{2}$$

$$f(x) * f(x) = \int_{x+\frac{1}{2}}^{x-\frac{1}{2}} -dy = \int_{x-\frac{1}{2}}^{x+\frac{1}{2}} dy = x + \frac{1}{2} - x - \frac{1}{2} = 1.$$

$$f(x) * f(x) = \begin{cases} 1 - |x| & ; -1 < x < 1 \\ 0 & ; \text{otherwise} \end{cases}$$





FLOTHEROU ALEXANDROS

871109-5590

SUR

20

B2

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

$$\begin{aligned}3) \quad g(w) &= \int_{-\infty}^{\infty} g(x) e^{-jwx} dx = \int_{-1}^1 (1-|x|) e^{-jwx} dx \\&= \int_{-1}^0 (1+x) e^{-jwx} dx + \int_0^1 (1-x) e^{-jwx} dx \\&= \int_{-1}^0 e^{-jwx} dx + \int_0^1 e^{-jwx} dx + \int_{-1}^0 x e^{-jwx} dx - \int_0^1 x e^{-jwx} dx \\&= -\frac{1}{jw} [e^{-jwx}]_{-1}^0 - \frac{1}{jw} [e^{-jwx}]_0^1 + \int_{-1}^0 x e^{-jwx} dx - \int_0^1 x e^{-jwx} dx \\&= \frac{1}{jw} \left( 1 - e^{jw} \right) - \frac{1}{jw} \left( e^{-jw} - 1 \right) + I(x) \\-TP \quad &= \frac{e^{jw} - e^{-jw}}{jw} = \frac{2}{w} \cdot \sin w + I(x)\end{aligned}$$

I don't have a mathematical handbook to solve  $I(x)$ . So far, the introduction of the sinc function means that the ringing effect is ~~here~~ here too. I don't know however if  $I(x)$  would cancel it.

6/8 You can use the product rule for integration.  
However, easier to use  $f(x) * f(x) \rightarrow f(w) \hat{f}(w)$  on 2.1 and 2.2.

$$I(x) = -\frac{2}{w} \sin w + \frac{\sin^2(w/2)}{(w/2)}$$



FLOTHEU ALEXANDROS

871108-5590 JZR

21

B.Z.

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

$$\text{D) } \int p(z) dz = \int (2z^4 - 3z^2 + \frac{8}{5}) dz \\ = \frac{2}{5}z^5 - z^3 + \frac{8}{5}z + C$$

$$\textcircled{10} \text{ for } z=0 \rightarrow C=1$$

$$z=1 \rightarrow \checkmark \\ \textcircled{10}, T(z) = \frac{2}{5}z^5 - z^3 + \frac{8}{5}z + 1.$$

\textcircled{11} For increasing the contrast:

$$\frac{dT}{dz} > 1 \Leftrightarrow 2z^4 - 3z^2 + \frac{8}{5} > 1.$$

~~$$z^2(2z^2 - 3) > -\frac{3}{5}$$~~

and  $z^2(2z^2 - 3) < -\frac{3}{5}$  for decreasing.

-0.5p  
What is the range?



FURTHEOU ALEXANDRIS

Family name, first name

871108-5590

Personal Registration Number

SZR

Programme

22 B.I

Sheet no.

Problem no.

3) ⑩ Mean:

$$f_{\text{mean}}(x) = \left[ \frac{7}{3}, \frac{11}{3}, 4, \frac{13}{3}, \frac{19}{3}, \frac{28}{3}, 11, \frac{37}{3}, \frac{35}{3}, 8 \right]$$

⑩ Median:

$$f_{\text{median}}(x) = [2, 4, 4, 4, 6, 10, 11, 12, 11, 10]$$

-1p Binomial?

2.

-3p

4.5/a

FULOTHEOU ALEXANDROS

871108-5190

SZR.

23

B.4

Family name, first name


Personal Registration Number

Programme

Sheet no.

Problem no.

1) ⑩ Moments:

$$M_{pq} = \sum_x \sum_y x^p y^q f(x,y)$$


⑩  $M_{00} = 12$

② ⑩  $M_{10} = 2 \cdot 1 + 3 \cdot 2 + 2 \cdot 3 + 3 \cdot 4 + 2 \cdot 5 = 32$

⑩  $M_{01} = 1 + 2 \cdot 2 + 4 \cdot 3 + 3 \cdot 4 + 2 \cdot 5 = 39$

2) ~~10~~

$$f(x=\frac{1}{2}) = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}, f(y) = \frac{1}{2} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

①  $f_x = \frac{1}{2} \begin{bmatrix} 1 & 1 & 0 \\ 1 & 1 & -1 \\ 0 & -1 & -1 \end{bmatrix}, f_y = \frac{1}{2} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 1 & 0 & -1 \end{bmatrix}$

$$C = \begin{bmatrix} f_x^2 & f_x f_y \\ f_x f_y & f_y^2 \end{bmatrix} = \begin{bmatrix} \frac{7}{4} & 1 \\ 1 & \frac{5}{4} \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 7 & 4 \\ 4 & 5 \end{bmatrix}$$

In order to find the orientation of the ellipse I would compute the eigenvalues of C first, then the eigenvector that corresponds to the lowest eigenvalue. That gives us the



FILOTHEN ALEXANDER

Family name, first name

871108-5590

Personal Registration Number

Programme

Sheet no.

24 B4

Problem no.

That, plus the centers of gravity  
 $\frac{m_0}{m_{00}}$ ,  $\frac{m_0}{m_{00}}$  would give us the line   
in which the ellipse is oriented.

① 3) A morphological opening is an erosion  
followed by a dilation.

► Erosion.

0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0

X

► Dilation:

No point, there are  
no foreground pixels.  
X

FILOTHEOU ALEXANDROS

87108-5590 SUR

25

B.6

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

2) Disparity:  $d = \frac{f \cdot b}{z} = \frac{10^3 \cdot 10^{-7}}{z} = \frac{100}{z}$


$$\frac{f \cdot b}{d} \rightarrow \frac{dz}{dd} = -\frac{f \cdot b}{d^2} = -\frac{z}{d}$$

~~at~~ differentiating with respect to the depth

(2)  $\frac{dd}{dz} = -\frac{f \cdot b}{z^2} = -\frac{d}{z} \Leftrightarrow dd = -\frac{d}{z} dz$

$$dd = -\frac{1000 \cdot 10^{-7}}{1^2} \cdot 10^{-3} = -10^{-1}$$

hence we need an accuracy of  $10^{-1}$  0.1 pixel  
unit is wrong





FILÔTHEROS ALEXANDROS 871108-5590 SIR 26 B.6

Family name, first name

Personal Registration Number

Programme

Sheet no.

Problem no.

⑩ Camera A at  $(Ax, 0, 0)$

⑩ Camera B at  $(Bx, 0, 0)$

⑩ Rotation matrix:

$$R = \begin{bmatrix} \cos\theta & 0 & -\sin\theta \\ 0 & 1 & 0 \\ \sin\theta & 0 & \cos\theta \end{bmatrix} \quad \theta = 2\pi$$

$$R = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

⑩ Translation matrix:

$$T = \begin{bmatrix} Bx - Ax \\ 0 \\ 0 \end{bmatrix}$$

⑩  $P_B = R \cdot P_A + T = R \cdot P_A + R \cdot R^{-1}T$

$$P_B = R(P_A + R^{-1}T)$$

⑩  $e = -R^{-1}T = \begin{bmatrix} Ax - Bx \\ 0 \\ 0 \end{bmatrix}$



FILÓTHEOU ALEXANDROR

Family name, first name

871108-5590 SCR

Personal Registration Number

Programme

27

R.G

Sheet no.

Problem no.

⑩ Essential matrix:

$$E = R \cdot [e]_X = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -Ax + Bx \\ 0 & -Ax + Bx & 0 \end{bmatrix}$$

$$E = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -Ax + Bx \\ 0 & -Ax + Bx & 0 \end{bmatrix} \quad \checkmark$$

③

1) ?

①