

PAPER

The impact of anisotropy on neutron star properties: insights from — universal relations


To cite this article: Sailesh Ranjan Mohanty *et al* JCAP03(2024)054

View the [article online](#) for updates and enhancements.

You may also like

- [Dynamical Formation Scenarios for GW190521 and Prospects for Decihertz Gravitational-wave Astronomy with GW190521-like Binaries](#)
A. Miguel Holgado, Alexis Ortega and Carl L. Rodriguez
- [Diffusive Tidal Evolution for Migrating Hot Jupiters](#)
Yanqin Wu and
- [Decay orbital period of the binary system on gravitational waves' detection](#)
P Y D Sagita, A Suroso and F P Zen

The impact of anisotropy on neutron star properties: insights from I – f – C universal relations

Sailesh Ranjan Mohanty^{^a}, Sayantan Ghosh^{^a}, Pinku Routaray^{^a}, H.C. Das^{^b}
and Bharat Kumar^{^{a,*}}

^aDepartment of Physics & Astronomy, National Institute of Technology,
Rourkela 769008, India

^bINFN Sezione di Catania, Dipartimento di Fisica,
Via S. Sofia 64, Catania 95123, Italy

E-mail: saileshranjanmohanty@gmail.com, sayantanghosh1999@gmail.com,
routaraypinku@gmail.com, harishdas.physics@gmail.com,
kumarbh@nitrkl.ac.in

ABSTRACT: Anisotropy in pressure within a star emerges from exotic internal processes. In this study, we incorporate pressure anisotropy using the Quasi-Local model. Macroscopic properties, including mass (M), radius (R), compactness (C), dimensionless tidal deformability (Λ), the moment of inertia (I), and oscillation frequency (f), are explored for the anisotropic neutron star. Magnitudes of these properties are notably influenced by anisotropy degree. Universal I – f – C relations for anisotropic stars are explored in this study. The analysis encompasses various EOS types, spanning from relativistic to non-relativistic regimes. Results show the relation becomes robust for positive anisotropy, weakening with negative anisotropy. The distribution of f -mode across M – R parameter space as obtained with the help of C – f relation was analyzed for different anisotropic cases. Using tidal deformability data from GW170817 and GW190814 events, a theoretical limit for canonical f -mode frequency is established for isotropic and anisotropic neutron stars. For isotropic case, canonical f -mode frequency for GW170817 event is $f_{1.4} = 2.606^{+0.457}_{-0.484}$ kHz; for GW190814 event, it is $f_{1.4} = 2.097^{+0.124}_{-0.149}$ kHz. These relationships can serve as reliable tools for constraining nuclear matter EOS when relevant observables are measured.

KEYWORDS: gravitational waves / theory, neutron stars

ARXIV EPRINT: [2305.15724](https://arxiv.org/abs/2305.15724)

*Corresponding author.

Contents

1	Introduction	1
2	Theoretical framework	3
2.1	Stellar structure equations	3
2.2	Anisotropy model	4
2.3	Slowly rotating NS and moment of inertia	5
2.4	Tidal deformability parameters	6
2.5	Non-radial oscillation in Cowling approximation	7
3	Universal relations	8
3.1	I – f relation	8
3.2	C – f relation	9
3.3	C – I relation	11
3.4	f -Love relation	12
3.5	Comparison study	14
4	Conclusion	15

1 Introduction

The detection of gravitational waves (GWs) is a crucial objective in astrophysics today, with significant efforts being devoted to this problem globally. Several ground-based experiments and space missions have already been devised and are poised to yield significant discoveries in the near future [1–5]. Furthermore, ongoing efforts are underway to develop a third-generation GW telescope, such as the Einstein Telescope [6] and Cosmic Explorer [7], which promise even higher levels of sensitivity. The reason why the detection of GWs is so challenging is that they are incredibly weak, necessitating detectors with very high sensitivity, as well as the precise waveform of the signal emitted from astrophysical objects.

Neutron stars (NSs) exhibit oscillations that are regarded as potential sources of GWs, manifesting in various forms including radial [8–13] and non-radial [14–17] modes. When a NS experiences external or internal disturbances, it emits GWs through different oscillation modes known as quasi-normal modes (QNMs), each characterized by the restoring force that brings them back to their equilibrium state. Notable QNMs include the fundamental mode (f -mode) [18–20], pressure mode (p -mode) [15, 21], gravity mode (g -mode) [22–29], rotational mode (r -mode) [30–35], space-time mode (w -mode) [36, 37], and other modes [38–41]. The frequencies of these oscillations are directly linked to the internal structure and composition of the stars [24]. Theoretical studies indicate that the f -mode possesses the highest likelihood of being detected initially, with approximately 10% of the gravitational radiation attributed to f -mode oscillations for $l = 2$ [42].

Different modes of oscillation exhibit distinct behaviors depending on the type of star, offering valuable insights. For instance, the signature of the hadron-quark phase transition

can be inferred through the observation of both f and g -modes in hybrid stars [39]. A study by Flores and Lugones [40] suggests that compact objects emitting GWs within the frequency range of 0–1 kHz may be hybrid stars, while frequencies exceeding 7 kHz could indicate strange stars. However, the nature of compact objects and their GW emissions still pose challenges due to the limitations of our terrestrial detectors, which are unable to detect certain frequency ranges. Nonetheless, constraints on the frequency of various modes can be established by establishing relationships between the frequency and specific properties of NSs. Several approaches have been proposed to link the f -mode frequency with various NS properties, including compactness [43], the moment of inertia [44], and tidal deformability [21, 45, 46]. In this work, we aim to derive such relationships in a model-independent manner, known as universal relations (URs). These URs provide valuable insights into the properties of NSs and their associated oscillation modes.

In literature, there are several URs have already been established between different properties of the NS [47–57]. However, the focus of this study primarily lies on the I – f – C relations specifically for anisotropic NSs. Previous URs proposed thus far have mainly been formulated for isotropic NSs, assuming a matter-energy distribution characterized by an isotropic perfect fluid. However, at extremely high-density regions, the presence of nuclear matter can induce deviations between the tangential and radial components of pressure, leading to the emergence of an anisotropic fluid. To obtain more accurate results, we aim to include the effects of anisotropy in our analysis. Anisotropy in NSs can arise from various factors, such as the influence of a high magnetic field [58–65], pion condensation [66], phase transitions [67], relativistic nuclear interactions [68, 69], core crystallization [70], and the presence of superfluid cores [71–73], among others. Several models have been proposed to incorporate anisotropy within NSs, such as the Bowers-Liang model [74], Horvat et al. model [75], Cosenza et al. model [76], and others. In this study, we will primarily focus on the Quasi-Local (QL) model, as proposed by Horvat et al. [75], to describe anisotropy within NSs. More details about the QL-model will be elaborated in section 2.2.

The presence of pressure anisotropy within NSs has a significant impact on various macroscopic properties, including the mass-radius relation, compactness, surface redshift, moment of inertia, tidal deformability, and non-radial oscillations [77–94]. The specific impact on these quantities varies depending on the degree of anisotropy and the choice of the model employed. Biswas and Bose constrained the degree of anisotropy within stars utilizing tidal deformability data from the GW170817 event [90]. More recently, the I -Love- C universal relation for anisotropic NSs has been proposed. By incorporating observational data from GW170817, constraints have been placed on the moment of inertia and radius of canonical anisotropic stars for different degrees of anisotropy [95].

In this study, we introduce the I – f – C URs for anisotropic NSs for the first time, utilizing a range of models describing unified EOSs such as RMF models (for $npe\mu$, hyperonic $npe\mu Y$, and strange $npe\mu Ys$ matter), density-dependent RMF models, and Skyrme-Hartree-Fock (SHF) models [15, 96–102]. The unified EOSs closely reproduce the properties of finite nuclei, nuclear matter, and NSs and support $\geq 2.0 M_\odot$ star. A total of 60 EOSs (35 from RMF models and 25 from SHF models) are considered in this paper. By employing these EOSs, we calculate various macroscopic properties of NSs with varying degrees of anisotropy using the QL-model. The primary objective of this work is to constrain the f -mode frequency

of anisotropic stars by leveraging recent observational data. Detailed formalism, results, and discussions are presented in the subsequent sections, shedding light on the intricate relationship between anisotropy, macroscopic properties, and the f -mode frequency in NSs.

2 Theoretical framework

2.1 Stellar structure equations

The line element that describes the space-time inside a spherically symmetric star is [103, 104]

$$ds^2 = -e^{2\psi} (dt)^2 + e^{2\lambda} dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \quad (2.1)$$

where, ψ and λ are metric functions that depend on r . Introducing anisotropy in the matter-energy distribution of the system, we obtain the following stress-energy tensor [83]

$$T_{\mu\nu} = (\mathcal{E} + P_t) u_\mu u_\nu + P_t g_{\mu\nu} - \sigma k_\mu k_\nu, \quad (2.2)$$

where u^μ represents the four-velocity of the fluid, and k^μ is a unit space-like four-vector. In addition, \mathcal{E} denotes the energy density, and σ represents the anisotropic pressure ($\sigma = P_t - P_r$), where P_t and P_r are the tangential and radial pressure, respectively. The four-vectors u^μ and k^μ must fulfill the following conditions:

$$u_\mu u^\mu = -1, \quad k_\mu k^\mu = 1, \quad u_\mu k^\mu = 0. \quad (2.3)$$

By solving Einstein field equations [105] for the anisotropic matter-energy distribution in spherically symmetric space-time, one can obtain the modified Tolman-Oppenheimer-Volkoff (TOV) [106] equations, which describes the stellar structure of an anisotropic NS given as

$$\begin{aligned} \frac{dm}{dr} &= 4\pi r^2 \mathcal{E}, \\ \frac{dP_r}{dr} &= -(P_r + \mathcal{E}) \left(\frac{m}{r^2} + 4\pi r P_r \right) e^{2\lambda} + \frac{2}{r} \sigma, \\ \frac{d\psi}{dr} &= -\frac{1}{P_r + \mathcal{E}} \frac{dP_r}{dr} + \frac{2\sigma}{r(P_r + \mathcal{E})}, \end{aligned} \quad (2.4)$$

where $m(r)$ is the enclosed mass corresponding to radius r and $\lambda(r)$ is the metric function defined as

$$e^{-2\lambda} = 1 - \frac{2m}{r}.$$

In order to numerically solve the aforementioned set of coupled ordinary differential equations (ODEs), specific boundary conditions need to be established. Conventionally, the surface of the star is set at $r = R$, where the radial pressure becomes zero ($P_r = 0$). As the equilibrium system exhibits spherical symmetry, the Schwarzschild metric is employed to describe the exterior space-time. This choice ensures metric continuity at the surface of the anisotropic neutron star (NS) and imposes a boundary condition on ψ . Specifically, the value of ψ at $r = R$ must coincide with the value of ψ in the Schwarzschild metric at $r = R$.

$$\psi(r = R) = \frac{1}{2} \ln \left[1 - \frac{2M}{R} \right].$$

By making a selection of an EOS governing the radial pressure (P_r) and adopting an anisotropic model for σ , it becomes feasible to numerically solve eq. (2.4). This numerical solution involves specifying a central energy density $\mathcal{E}(r=0) = \mathcal{E}_c$, while enforcing the initial condition $m(r=0) = 0$.

2.2 Anisotropy model

For anisotropic NS, we use the Quasi-Local (QL) model proposed by Horvat et al. [75] describing the quasi-local nature of anisotropy in the following

$$\sigma = P_t - P_r = \frac{\beta_{\text{QL}}}{3} P_r \mu = \frac{\beta_{\text{QL}}}{3} P_r (1 - e^{-2\lambda}), \quad (2.5)$$

where the factor β_{QL} depicts the measurement of the degree of anisotropy in the fluid and ($\mu = 1 - e^{-2\lambda} = \frac{2m(r)}{r}$) also known as local compactness is the quasi-local variable. In order to maintain their spherically symmetric configuration, anisotropic NSs must adhere to specific conditions, as outlined in references [83, 94]. These conditions include:

- Absence of anisotropy at the center of the NS i.e. $\sigma = 0$, or equivalently, $P_r = P_t$ at $r = 0$.
- Positivity of P_r and P_t throughout the entire star.
- Positivity of the null energy density (\mathcal{E}), dominant energy density ($\mathcal{E} + P_r, \mathcal{E} + P_t$), and strong energy density ($\mathcal{E} + P_r + 2P_t$) within the star.
- Non-negativity of the sound speed (c_s^2) inside the star, with the c_s^2 in the radial and tangential directions satisfying the following constraints: $0 < c_{s,r}^2, c_{s,t}^2 < 1$. It is also essential to ensure that the speed of sound does not exceed the speed of light ($c = 1$ in this study).

Therefore, the conditions mentioned above are crucial for maintaining the spherical symmetry and physical consistency of anisotropic NSs.

The advantage of using the QL-model with anisotropy is that it ensures the fluid remains isotropic at the center of the star due to the behavior of $(1 - e^{-2\lambda} = 2m/r) \sim r^2$ when $r \rightarrow 0$, while also being applicable only to relativistic configurations where anisotropy may arise at high densities [89]. For the 60 EOS-ensembles considered in this study, the QL-model with anisotropy parameter ranging from $-2 < \beta_{\text{QL}} < 2$ satisfies all the necessary conditions to maintain spherical symmetry in an anisotropic NS configuration [83, 94]. Figure 1 shows that the speed of sound in the tangential direction ($c_{s,t}^2 = \frac{\partial P_t}{\partial \mathcal{E}}$) for maximum mass configuration corresponds to DD2 EOS, satisfies the causality condition throughout the star for $-2 < \beta_{\text{QL}} < 2$.

The mass-radius (MR) profile for a given EOS can be obtained by solving the TOV eqs. (2.4) for various central densities, which generate a sequence of mass and radius. Figure 2 illustrates the MR profiles for the anisotropic star for the DD2 EOS. Adjusting the value of β_{QL} influences the maximum mass and the corresponding radius of the NS. The positive value of β_{QL} increases the maximum mass and its associated radius, and vice-versa for β_{QL} . Observational data from different observations, such as X-ray, NICER, and GW (GW170817




Figure 1. The radial profile of the sound speed ($c_{s,t}^2$) for different β_{QL} values of the maximum mass NS corresponding to the DD2 EOS.




Figure 2. *Left:* mass-radius profiles for anisotropic NSs with $-2 < \beta_{\text{QL}} < 2$ for the DD2 EOS. The black dashed line represents the isotropic case. The limits on mass and radius from the PSR J0030+0451 [108, 109] are shown in the light pink boxes, and the revised NICER data [110] is shown in the light blue boxes. The green horizontal bar represents the mass range observed in the GW190814 event [112]. *Right:* the MI as a function of mass for different values of β_{QL} . The error bars were calculated based on the results of several pulsar analyses as done in ref. [97].

and GW190814), to constrain the degree of anisotropy within NS [108–112]. For example, values of $1 < \beta_{\text{QL}} < 2$ satisfy the mass constraint ($2.50\text{--}2.67 M_{\odot}$) of the GW190814 event, suggesting that one of the merger companions may have been a highly anisotropic NS [81].

2.3 Slowly rotating NS and moment of inertia

The MI of a slowly rotating anisotropic NS can be expressed as [86]

$$I = \frac{8\pi}{3} \int_0^R r^4 e^{\lambda-\psi} (\mathcal{E} + P_r + \sigma) \frac{\bar{\omega}}{\Omega} dr, \quad (2.6)$$

where, $\bar{\omega}$ is the frame-dragging angular frequency [113], and Ω is the angular velocity of a uniformly rotating NS. The MI of an anisotropic NS is shown as a function of its mass in the left panel of figure 2. As the NS mass increases, the MI also increases until a stable




Figure 3. *Left:* the Λ as a function of the mass for different values of the β_{QL} corresponds to DD2 EOS. The error bars in both panels represent the observational constraints from GW170817 [111] and GW190814 events [112]. *Right:* the f -mode frequency as a function mass. The error bars represent the theoretical limits that we obtained in subsection 3.4.

configuration is reached, after which it starts to decrease. Furthermore, both the mass and MI of the NS increase with positive values of β_{QL} , while the opposite trend is observed for negative values of β_{QL} . The impact of anisotropy on the MI is more pronounced for high-mass NSs compared to low-mass ones. Kumar and Landry [97] have established constraints on the MI inferred from various sources such as double neutron stars (DNS), millisecond pulsars (MSP), and low-mass X-ray binaries (LMXB). The error bars in the figure represent the possible range of values for these constraints.

2.4 Tidal deformability parameters

When the NS is present in the external field (ϵ_{ij}) created by its companion star, it acquires a quadrupole moment (Q_{ij}). The magnitude of the quadrupole moment is linearly proportional to the tidal field and is given by [114, 115]

$$Q_{ij} = -\alpha\epsilon_{ij}, \quad (2.7)$$

where α is the tidal deformability of a star. α can be defined in terms of tidal Love number k_2 as $\alpha = \frac{2}{3}k_2R^5$. The dimensionless tidal deformability of the star is defined as $\Lambda = \alpha/M^5 = \frac{2}{3}k_2C^{-5}$. The detailed derivation of k_2 for an anisotropic star can be found in refs. [90, 94, 95].

The dimensionless tidal deformability of anisotropic NSs is shown in figure 3. As the anisotropy parameter β_{QL} increases, the magnitude of the Love number k_2 and its corresponding tidal deformability Λ decrease, while they increase with decreasing β_{QL} . The impact of anisotropy on tidal deformability, as mentioned above, reverses after attaining maximum mass configuration, beyond which the star becomes unstable. The GW170817 (NS-NS merger) event constrains $\Lambda_{1.4}$ to be 190^{+390}_{-120} [111]. In the case of the GW190814 merger event, one should not rule out the possibility of the lower mass component being a massive NS. With the spectral EOS distribution and conditioned used for GW170817 [116], each EOS is reweighted by the probability that its maximum mass is at least as large as the

mass of the secondary component ($2.6 M_\odot$), which put a limit on $\Lambda_{1.4} = 616_{-158}^{+273}$ [112] (in the NS-BH scenario). In this case, the predicted value of $\Lambda_{1.4}$ satisfies the GW190814 limit for almost all values of β_{QL} , whereas, for DD2 EOS, the GW170817 limit is met in the range of $0.5 < \beta_{\text{QL}} < 2$. However, Λ sharply decreases once the stable configuration is exceeded.

2.5 Non-radial oscillation in Cowling approximation

The Cowling approximation, initially proposed by Cowling [117] for Newtonian stars and later extended to the case of NSs by McDermott et al. [14]. Under this approximation, the metric perturbations are neglected, keeping the space-time metric fixed. We will provide a brief explanation of the derivation of the perturbation equations in the Cowling formalism in the following, while more comprehensive details can be found in [79]. One can obtain the oscillation equations in the Cowling approximation by considering a harmonic time dependence for the perturbation function $W(r, t) = W(r)e^{i\omega t}$ and $V(r, t) = V(r)e^{i\omega t}$, where ω represents the oscillation frequency in the following [79, 118]

$$\begin{aligned} W' &= \frac{d\mathcal{E}}{dP_r} \left[\omega^2 \frac{\mathcal{E} + P_t}{\mathcal{E} + P_r} \left(1 + \frac{\partial\sigma}{\partial P_r} \right)^{-1} e^{\lambda-2\psi} r^2 V + \psi' W \right] - l(l+1)e^\lambda V \\ &\quad - \frac{\sigma}{\mathcal{E} + P_r} \left[\frac{2}{r} \left(1 + \frac{d\mathcal{E}}{dP_r} \right) W + l(l+1)e^\lambda V \right], \\ V' &= 2V\psi' - \left(1 + \frac{\partial\sigma}{\partial P_r} \right) \frac{\mathcal{E} + P_r}{\mathcal{E} + P_t} \frac{e^\lambda}{r^2} W - \left[\frac{\sigma'}{\mathcal{E} + P_t} + \left(\frac{d\mathcal{E}}{dP_r} + 1 \right) \frac{\sigma}{\mathcal{E} + P_t} \left(\psi' + \frac{2}{r} \right) \right. \\ &\quad \left. - \frac{2}{r} \frac{\partial\sigma}{\partial P_r} - \left(1 + \frac{\partial\sigma}{\partial P_r} \right)^{-1} \left(\frac{\partial^2\sigma}{\partial P_r^2} P_r' + \frac{\partial^2\sigma}{\partial P_r \partial \mu} \mu' \right) \right] V. \end{aligned} \quad (2.8)$$

To solve the equations mentioned earlier, it is necessary to consider boundary conditions at the center and surface of the star in the following

$$\omega^2 \frac{\mathcal{E} + P_t}{\mathcal{E} + P_r} \left(1 + \frac{\partial\sigma}{\partial P_r} \right)^{-1} e^{-2\psi} V + \left(\psi' - \frac{2}{r} \frac{\sigma}{\mathcal{E} + P_r} \right) e^{-\lambda} \frac{W}{r^2} = 0, \quad (2.9)$$

and the boundary condition at the star center ($r = 0$) satisfies

$$\tilde{W} = -l\tilde{V}$$

where, the functions \tilde{W} and \tilde{V} are defined as $W = \tilde{W}r^{l+1}$ and $V = \tilde{V}r^l$. In this work, we focus on the quadrupolar modes, which correspond to $l = 2$. In figure 3, we show the f -mode frequency of a NS as a function of its mass by varying the anisotropic parameter for the DD2 EOS as a representative case. For a specific mass NS, the frequency decreases for a positive value of β_{QL} while increases for a negative β_{QL} till maximum mass is attained, after which the star becomes unstable. Using the tidal deformability limit from GW170817 and GW190814, one can impose constraints on the canonical f -mode frequency for isotropic and anisotropic stars, as discussed in subsection 3.4. We also overlaid the derived theoretical limit on the figure to assess its consistency.

3 Universal relations

The main purpose of UR is to explore the star properties that are difficult to measure through observations. Several URs have already been proposed to estimate the properties of NS, but they are primarily focused on isotropic cases [47, 49, 119–122]. However, very few studies have been dedicated to URs for anisotropic stars, which are more realistic than isotropic ones. Hence, in this study, we aim to explore various types of URs between the moment of inertia, tidal deformability, compactness, and f -mode frequency for anisotropic NSs.

Although some known URs for anisotropic NS have been proposed in refs. [48, 90, 95], our primary focus will be on the URs between the moment of inertia, f -mode frequency, and compactness (I – f – C) as well as the f –Love relation of the anisotropic NS. The NS oscillates with different modes, emitting gravitational waves (GWs). The oscillation frequencies, such as the f -mode, p -mode, etc., might be detectable in the near future with our terrestrial detectors. However, to interpret these observations effectively, we require prior theoretical knowledge. Therefore, approximate URs for anisotropic NSs hold great significance in astrophysical observations.

Before delving into different URs, it is necessary to normalize/dimensionless certain key parameters of NSs that are required to obtain the URs. Here, we chose the units of MI and f -mode frequency are $\text{kg}\cdot\text{m}^2$ and kHz , respectively. Therefore, these quantities need to be normalized, and their normalized values are given as

- Normalized MI (η) = $\sqrt{M^3/I}$
- Normalized f -mode frequency ($\bar{\omega}$) = ωM

Here, we calculate the URs between I – f , C – f , I – C , and f –Love for anisotropic NSs. For this study, we chose 60 EOSs, as mentioned in the introduction. Regarding anisotropy, we adopt the same QL-model with different degrees of anisotropy, varying from -2 to $+2$. Alternatively, one may opt for other models, such as Bower-Liang’s, as used in ref. [95].

3.1 I – f relation

The I – f UR for isotropic NS with a few EOSs was first calculated by Lau et al. [44]. Breu and Rezzolla [49] studied the universal behavior of dimensionless MI, which is defined as $\bar{I} = I/M^3$, and is more accurate than the dimensionless MI defined earlier, $\bar{I} = I/MR^2$. Lau and Leung replaced $\bar{I} = I/M^3$ with $\eta = \sqrt{M^3/I}$, called as normalized MI/effective compactness, due to its proportionality with compactness for stars. Therefore, in this study, we use $\eta = \sqrt{M^3/I}$ rather than $\bar{I} = I/MR^2$. The relation between I – f for anisotropic NSs is performed using the least-squares fit with the approximate formula

$$\eta = \sum_{n=0}^{n=4} a_n (\bar{\omega})^n. \quad (3.1)$$

The normalized MI (η) is plotted as the function of normalized f -mode frequency ($\bar{\omega}$) in figures 4–5 with $\beta_{\text{QL}} = -2, 0, +2$ for anisotropic NS. The residuals are computed with the formula,

$$\Delta\eta = \frac{\eta - \eta_{\text{fit}}}{\eta_{\text{fit}}}. \quad (3.2)$$




Figure 4. I – f relation with anisotropy parameter $\beta_{\text{QL}} = 0$ for selected EOSs. The black-dashed line shows the least-squares fit using eq. (3.1). The lower panel displays the residuals of the fitting obtained using the formula in eq. (3.2).




Figure 5. *Left:* same as figure 4, but with $\beta_{\text{QL}} = +2$. *Right:* for $\beta_{\text{QL}} = -2$.

We enumerated the coefficients (a_n) with their corresponding reduced chi-squared (χ_r^2) errors in table 1. An increase in anisotropy results in a decrease in the value of χ_r^2 error, indicating stronger EOS insensitive relation, and vice versa.

3.2 C – f relation

Andersson and Kokkotas [38] first established the correlation between C and f -mode frequency. Here, we calculate the C – f relations for anisotropic NSs, using the approximate formula obtained through least-squares fitting

$$C = \sum_{n=0}^{n=4} b_n(\bar{\omega})^n. \quad (3.3)$$

Compactness is plotted as the function of normalized f -mode frequency ($\bar{\omega}$) in figures 6–7 with $\beta_{\text{QL}} = -2, 0, +2$ for anisotropic NS. The coefficients (b_n) with χ_r^2 -error are enumerated in table 1. The magnitude of b_n increases with increasing β_{QL} , implying that the fitting

<i>I</i> – <i>f</i>						<i>C</i> – <i>f</i>					
$\beta_{\text{QL}} =$	–2.0	–1.0	0.0	+1.0	+2.0	$\beta_{\text{QL}} =$	–2.0	–1.0	0.0	+1.0	+2.0
$a_0 (10^{-2}) =$	2.825	2.848	2.918	2.886	2.903	$b_0 (10^{-3}) =$	4.007	4.093	4.084	4.197	4.249
$a_1 =$	3.945	3.919	3.865	3.874	3.856	$b_1 =$	2.232	2.220	2.223	2.212	2.209
$a_2 (10^1) =$	–2.436	–2.303	–2.139	–2.092	–2.011	$b_2 =$	–8.752	–8.143	–7.963	–7.531	–7.269
$a_3 (10^2) =$	1.364	1.218	1.059	0.982	0.889	$b_3 (10^1) =$	3.066	2.656	2.629	2.371	2.211
$a_4 (10^2) =$	–2.852	–2.457	–2.064	–1.852	–1.613	$b_4 =$	4.836	6.740	–3.179	5.173	–7.423
$\chi_r^2 (10^{-6}) =$	13.511	8.514	5.338	3.541	2.517	$\chi_r^2 (10^{-6}) =$	2.228	1.702	1.846	2.282	2.871

<i>C</i> – <i>I</i>					
$\beta_{\text{QL}} =$	–2.0	–1.0	0.0	+1.0	+2.0
$c_0 (10^{-3}) =$	–8.651	–8.244	–8.941	–7.860	–7.637
$c_1 (10^{-1}) =$	4.473	4.348	4.477	4.236	4.171
$c_2 =$	1.360	1.465	1.389	1.557	1.612
$c_3 =$	–3.986	–4.288	–4.080	–4.516	–4.668
$c_4 =$	4.892	5.272	5.147	5.610	5.827
$\chi_r^2 (10^{-6}) =$	6.338	6.382	6.436	6.313	6.210

Table 1. The fitting coefficients are listed for *I*–*f*, *C*–*f*, and *C*–*I* relations with $\beta_{\text{QL}} = -2.0, -1.0, 0.0, +1.0, +2.0$. The reduced chi-squared (χ_r^2) is also given for all cases.




Figure 6. *C*–*f* relation with anisotropy parameter $\beta_{\text{QL}} = 0$ for assumed EOSs. The black-dashed line is fitted with eq. (3.3). The lower panel shows the residuals for the fitting are calculated.

is more robust for the isotropic case. Additionally, χ_r^2 also increases with the inclusion of anisotropy, and it is the minimum for the isotropic case. Therefore, the inclusion of anisotropy (whether positive or negative) weakens the EOS insensitive *C*–*I* UR.

One of the primary applications of *C*–*f* UR involves determining *M* and *R* based on the analysis of observed mode data, as articulated by Andersson and Kokkotas [43]. For a unique choice of *f*-mode frequency the *C*–*f* UR can be exploited to construct a *M*–*R* relation. This constrained relationship, accounting for uncertainties represented by standard deviations in UR, yields *M*–*R* bands, illustrated in the left panel of figure 8. In this representation, the orange *M*–*R* band delineates a region where neutron stars are anticipated to exhibit a




Figure 7. *Left:* same as figure 6, but with $\beta_{\text{QL}} = +2$. *Right:* for $\beta_{\text{QL}} = -2$.

frequency of $f = 2.606^{+0.457}_{-0.484}$ kHz, with the solid dashed line denoting $f = 2.606$ kHz. Similarly, the pink band corresponds to a region where neutron stars are expected to possess a frequency of $f = 2.097^{+0.124}_{-0.149}$ kHz, and the solid dashed line represents $f = 2.097$ kHz. It is noteworthy that the frequency constraints employed for plotting M – R bands align with the canonical f -mode frequency constraints for isotropic neutron stars determined in this study for the GW170817 and GW190814 events. The horizontal error bars in the left panel of figure 8 indicate radius limits imposed by the M – R bands of the respective events, considering a canonical mass neutron star.

The right panel of figure 8 portrays the distribution of f -mode frequencies across the M – R parameter space for isotropic neutron stars based on C – f UR. The black dashed line represents a specific set of mass and radius values for isotropic stars, anticipated to exhibit the mentioned frequency according to C – f UR. The figure also depicts variations in these M – R lines resulting from the inclusion of anisotropy. NSs with frequencies $f < 1.5$ kHz lie in the low compactness region and suffer minimal changes in mass and radius due to the inclusion of anisotropy. Through observing M – R lines as depicted in figure 8, we can conclude that for a constant mass NS having a fixed frequency with $f \geq 1.5$ kHz, the radius would tend to decrease with the presence of positive anisotropy and increase for negative anisotropy, altering the compactness of the star in order to maintain its natural frequency till a certain critical point/set of mass and radius is reached. After this, the effect of anisotropy on M – R lines reverses. This kind of behavior, in which the effects of anisotropy on the NS parameter reverses, is likely to originate due to an unstable core. We suspect that this kind of instability arises due to presence of central density beyond the maximal stable configuration ($\rho_c > \bar{\rho}_c$) following the standard stability criterion $\left(\frac{\partial M}{\partial \rho_c}\right)_{\rho_c=\bar{\rho}_c} = 0$ [123]. This suggests that the critical mass-radius point in the M – R curves is the maximum stability point, beyond which the NSs become unstable in nature.

3.3 C – I relation

The relationship between the dimensionless MI ($\bar{I} = I/M^3$) and compactness has been established as a lower-order polynomial fit by Ravenhall and Pethick [124]. Since then, this




Figure 8. *Left:* mass-radius profiles of isotropic NSs for EOS ensembles that we have considered in this study. The orange and pink colored MR bands correspond to limits on NS’s mass and radius imposed by isotropic C – f UR for the canonical f -mode frequency ($f_{1.4}$) that was obtained through f -Love UR with the help of tidal deformability constraints ($\Lambda_{1.4}$) of GW170817 [111] and GW190814 [112] events, respectively. The horizontal error bars illustrate the radius limits for a canonical mass NS imposed by the frequency bands for respective events. *Right:* the frequency distribution contour plot across the M – R parameter space. This distribution is imposed by the isotropic C – f UR. M – R lines for isotropic cases corresponding to a set of frequencies are shown in black dashed-dot lines. The red and blue dashed lines represent the M – R curves for anisotropy parameter $\beta_{\text{QL}} = +2$ and -2 , respectively.

relation has been studied and modified by various authors, including for the double pulsar system with higher-order polynomial fitting [125], scalar-tensor theory and R^2 gravity [122, 126], rotating stars [49], and strange stars [127]. In this work, we investigate the C – I relations for anisotropic NS using the normalized moment of inertia ($\eta = \sqrt{M^3/I}$) instead of the dimensionless one. We use the approximate formula to perform a least-squares fit

$$C = \sum_{n=0}^{n=4} c_n(\eta)^n. \quad (3.4)$$

We display the relationship between C and η for anisotropic NSs with $\beta_{\text{QL}} = -2, 0, +2$, respectively in figures 9–10. We observe that the inclusion of anisotropy has little effect on the χ_r^2 error, indicating that the C – I UR is conserved even when anisotropy is present, especially for NSs with low compactness.

3.4 f -Love relation

An important tool for studying the oscillation of NSs through observational exploration is a UR between the non-radial f -mode frequency (a promising source of GWs) and the tidal deformability (a parameter that can be extracted from the GW data). The exploration of multi-polar universal relations between the f -mode frequency and tidal deformability of compact stars was first explored by Chan et al. [45] and further improvised by Pradhan et al. [46]. Recently, Sotani and Kumar [21] introduced a UR between the quasi-normal modes and tidal deformability for isotropic NSs. In this work, we calculate the f -Love relations for




Figure 9. C – I relation with anisotropy parameter $\beta_{\text{QL}} = 0$ for assumed EOSs. The black-dashed line is fitted with eq. (3.4). The lower panel shows the residuals for the fitting are calculated.




Figure 10. *Left:* same as figure 9, but with $\beta_{\text{QL}} = +2$. *Right:* same as figure 9, but with $\beta_{\text{QL}} = -2$.

anisotropic NSs and perform a least-squares fit using the approximate formula

$$\bar{\omega} = \sum_{n=0}^{n=4} d_n (\log(\Lambda))^n. \quad (3.5)$$

The relationship between $\bar{\omega}$ and Λ for $\beta_{\text{QL}} = -2, 0, +2$ cases are depicted in figures 11–12. The coefficients (d_n) with χ_r^2 errors are listed in table 2. For positive values of anisotropy, the errors in χ_r^2 decrease, indicating that the EOS-insensitive relations become stronger with the addition of anisotropy. Conversely, for negative values, the errors increase. Therefore, positive values of anisotropy strengthen the f -Love UR. With the help of tidal deformability constraints of a canonical mass NS for GW170817 [111] ($\Lambda_{1.4} = 190_{-120}^{+390}$), and GW190814 [112] ($\Lambda_{1.4} = 616_{-158}^{+273}$) events, we establish theoretical limits on the f -mode frequency for each event with different degrees of anisotropy, utilizing the f -Love URs obtained in this study. We assume the later event (GW190814) to be a NS-BH merger event as explained in ref. [81] to support our findings. The pink-dash and orange-dot regions in figure 11–12 represent the tidal constraints for events GW190814 and GW170817 respectively, and the corresponding




Figure 11. The f -Love relation for anisotropic NSs with $\beta_{\text{QL}} = 0$ for various assumed EOSs. The black-dashed line represents the best fit using eq. (3.5). The light-pink-shaded region and the orange-shaded region represent the range of canonical tidal deformability data obtained from the GW190814 [112] and GW170817 [111] papers, respectively.




Figure 12. *Left:* same as figure 11, but with $\beta_{\text{QL}} = +2$. *Right:* for $\beta_{\text{QL}} = -2$.

vertical lines represent the values of constrained $\bar{\omega}$. The canonical f -mode frequency imposed by events GW170817 and GW190814 for different degrees of anisotropy as obtained in this study is enumerated in table 3.

3.5 Comparison study

We constrain the canonical f -mode frequency for GW170817 [111] and GW190814 [112] events across different degrees of anisotropy, as outlined in table 3. The canonical f -mode frequency is also compared with previous studies, focusing on isotropic NS, as listed in table 4. Notably, the f -mode frequency obtained in this study is approximately 30–35% more than the findings of Chan et al. [45], Pradhan et al. [46], and Sotani and Kumar [21]. This difference in the f -mode was anticipated, given that the aforementioned authors employed a full-GR formalism for their f -mode calculations, in contrast to our use of the Cowling approximation in this study.

$\beta_{\text{QL}} =$	-2.0	-1.0	0.0	1.0	+2.0
$d_0 (10^{-1}) =$	2.001	2.037	2.077	2.131	2.169
$d_1 (10^{-2}) =$	-0.964	-1.998	-2.722	-3.607	-4.158
$d_2 (10^{-2}) =$	-1.857	-1.443	-1.215	-0.874	-0.699
$d_3 (10^{-3}) =$	3.795	3.207	2.975	2.406	2.260
$d_4 (10^{-4}) =$	-2.155	-1.873	-1.815	-1.538	-1.464
$\chi_r^2 (10^{-6}) =$	4.149	2.311	1.269	0.917	0.735

Table 2. The fitting coefficients are listed for f -Love relation with $-2.0 < \beta_{\text{QL}} < +2.0$. The reduced chi-squared (χ_r^2) is also given for all cases.

β_{QL}	GW170817		GW190814	
	$\bar{\omega}_{1.4}$	$f_{1.4}$	$\bar{\omega}_{1.4}$	$f_{1.4}$
-2.0	$0.121^{+0.020}_{-0.022}$	$2.790^{+0.453}_{-0.499}$	$0.098^{+0.006}_{-0.007}$	$2.265^{+0.130}_{-0.157}$
-1.0	$0.116^{+0.020}_{-0.021}$	$2.680^{+0.451}_{-0.487}$	$0.094^{+0.005}_{-0.007}$	$2.168^{+0.126}_{-0.152}$
0.0	$0.113^{+0.020}_{-0.021}$	$2.606^{+0.457}_{-0.484}$	$0.091^{+0.005}_{-0.006}$	$2.097^{+0.124}_{-0.149}$
+1.0	$0.111^{+0.020}_{-0.021}$	$2.550^{+0.461}_{-0.481}$	$0.089^{+0.005}_{-0.006}$	$2.044^{+0.123}_{-0.147}$
+2.0	$0.109^{+0.020}_{-0.021}$	$2.508^{+0.467}_{-0.482}$	$0.087^{+0.005}_{-0.006}$	$2.001^{+0.122}_{-0.146}$

Table 3. The canonical normalized f -mode frequency ($\bar{\omega}_{1.4}$), and f -mode frequency ($f_{1.4}$ in kHz) inferred from GW170817 and GW190814 data.

Ref.	GW170817	GW190814
	$f_{1.4}$	$f_{1.4}$
Chan et al. [45]	$2.120^{+0.445}_{-0.446}$	$1.652^{+0.111}_{-0.130}$
Pradhan et al. [46]	$2.120^{+0.444}_{-0.445}$	$1.653^{+0.111}_{-0.130}$
Sotani and Kumar [21]	$2.124^{+0.440}_{-0.446}$	$1.656^{+0.112}_{-0.132}$
This Work	$2.606^{+0.457}_{-0.484}$	$2.097^{+0.124}_{-0.149}$

Table 4. The canonical f -mode frequency ($f_{1.4}$ in kHz) inferred from GW170817 and GW190814 data using f -Love UR obtained in different literature for isotropic NS.

4 Conclusion

In this study, we have explored the properties of anisotropic NS with the help of the QL-model proposed by Horvat et al. [75]. The main motivation for taking the QL-model is that it ensures that $r \rightarrow 0$, the anisotropy must vanish, and in other parts of the star, the anisotropy must be there. Different fluid conditions are also studied for varieties of EOSs, and it found that all conditions are perfectly satisfied for the QL model. The speed of sound is also non-negative with any degree of anisotropy for the QL-model in comparison to the BL-model, as mentioned in refs. [90, 95]. Therefore, one can vary the limit of the QL-model from negative to positive values to calculate various properties of the NS.

Different macroscopic properties of the star have been calculated with different degrees of anisotropy with the help of a variety EOSs spanning from relativistic to non-relativistic cases. It has been observed that the magnitude of the macroscopic properties increases (decreases) for positive (negative) values of β_{QL} . Almost all the considered EOSs satisfy the different observational limits provided by different observations such as X-ray, pulsar, NICER, GWs, etc. One can impose strong constraints on them with the help of these observational data. Furthermore, we found that positive and negative anisotropy affects tidal deformability parameters and quadrupolar non-radial f -mode frequency significantly, which suggests that

the star with higher anisotropy sustains more life in the inspiral-merger phase, while the star with lower anisotropy is more likely to collapse.

In addition, we have studied the I - f - C UR for anisotropic NSs for five values of $\beta_{\text{QL}} = -2.0, -1.0, 0.0, +1.0$, and $+2.0$. This analysis considered almost 60 tabulated EOS-ensembles spanning a wide range of stiffness, complying with multimessenger constraints. Moreover, one can use the I - f - C universal relation for anisotropic stars to extract information about different properties that are not directly observable with current detectors and telescopes. By varying the anisotropy value, we calculated the I - f , C - f , and C - I universal relations and fitted them with the polynomial equation using the least-square method. Our results showed that the reduced chi-square errors for the I - f , C - f , and C - I relations were 5.4863×10^{-6} , 2.0599×10^{-6} , and 6.8402×10^{-6} , respectively, for isotropic stars. In addition to the I - f - C universal relations, we calculated the f -Love universal relation to constrain the canonical f -mode frequency for anisotropic stars. We observed that the sensitivity of the C - f universal relation is weaker for anisotropic stars in comparison to the isotropic case. However, the relation between I - f and f -Love became stronger with increasing anisotropy. The C - I relation barely changed with the inclusion of anisotropy compared to the other universal relations. The distribution of f -mode across mass-radius parameter space of NSs as obtained by utilizing the C - f relation studied for different anisotropic cases.

With the help of various observational data for dimensionless tidal deformability, such as GW170817 and GW190814, we established a theoretical constraint on the canonical f -mode frequency for both isotropic and anisotropic stars, which is presented in table 3. As our main objective in this paper was to analyze variations in I - f - C URs resulting from the inclusion of anisotropy, we adhered to the Cowling approximation formalism for computing the f -mode. This choice was necessitated by the absence of a comprehensive and reliable full GR formalism for determining QNM in anisotropic NSs. Consequently, for compensation, we calculated constraints on the canonical f -mode frequency for isotropic stars, relying on URs obtained by researchers in refs. [45, 46, 128], which followed a full-GR formalism, and summarized the outcomes in table 4. This constraint can be refined by incorporating different anisotropy models and considering various phenomena such as magnetic fields, quarks in the core, and dark matter in detail in future work. Therefore, our findings provide avenues for investigating the various mechanisms that generate anisotropy within compact stars and for constraining its degree with observational data.

Acknowledgments

I would like to thank P. Landry and T. Zhao for their fruitful discussions regarding universal relations and fitting procedures. B.K. acknowledges partial support from the Department of Science and Technology, Government of India, with grant no. CRG/2021/000101.

References

- [1] B.P. Abbott et al., *LIGO: The Laser interferometer gravitational-wave observatory*, *Rept. Prog. Phys.* **72** (2009) 076901 [[arXiv:0711.3041](#)] [[INSPIRE](#)].
- [2] G.M. Harry, *Advanced LIGO: The next generation of gravitational wave detectors*, *Class. Quant. Grav.* **27** (2010) 084006 [[INSPIRE](#)].

- [3] F. Acernese et al., *The Virgo status*, *Class. Quant. Grav.* **23** (2006) S635 [INSPIRE].
- [4] VIRGO collaboration, *Calibration and sensitivity of the Virgo detector during its second science run*, *Class. Quant. Grav.* **28** (2011) 025005 [Erratum *ibid.* **28** (2011) 079501] [[arXiv:1009.5190](#)] [INSPIRE].
- [5] F. Antonucci et al., *From laboratory experiments to LISA Pathfinder: achieving LISA geodesic motion*, *Class. Quant. Grav.* **28** (2011) 094002 [[arXiv:1012.5968](#)] [INSPIRE].
- [6] M. Punturo et al., *The third generation of gravitational wave observatories and their science reach*, *Class. Quant. Grav.* **27** (2010) 084007 [INSPIRE].
- [7] E.D. Hall, *Cosmic Explorer: A Next-Generation Ground-Based Gravitational-Wave Observatory*, *Galaxies* **10** (2022) 90 [INSPIRE].
- [8] S. Chandrasekhar, *The Dynamical Instability of Gaseous Masses Approaching the Schwarzschild Limit in General Relativity*, *Astrophys. J.* **140** (1964) 417 [Erratum *ibid.* **140** (1964) 1342] [INSPIRE].
- [9] G. Chanmugam, *Radial oscillations of zero-temperature white dwarfs and neutron stars below nuclear densities*, *Astrophys. J.* **217** (1977) 799.
- [10] K.D. Kokkotas and J. Ruoff, *Radial oscillations of relativistic stars*, *Astron. Astrophys.* **366** (2001) 565 [[gr-qc/0011093](#)] [INSPIRE].
- [11] P. Routaray, H.C. Das, S. Sen, B. Kumar, G. Panotopoulos and T. Zhao, *Radial oscillations of dark matter admixed neutron stars*, *Phys. Rev. D* **107** (2023) 103039 [[arXiv:2211.12808](#)] [INSPIRE].
- [12] S. Sen et al., *Radial Oscillations in Neutron Stars from Unified Hadronic and Quarkyonic Equation of States*, *Galaxies* **11** (2023) 60 [[arXiv:2205.02076](#)] [INSPIRE].
- [13] P. Routaray, A. Quddus, K. Chakravarti and B. Kumar, *Probing the impact of WIMP dark matter on universal relations, GW170817 posterior, and radial oscillations*, *Mon. Not. Roy. Astron. Soc.* **525** (2023) 5492 [[arXiv:2202.04364](#)] [INSPIRE].
- [14] P.N. McDermott, H.M. van Horn and C.J. Hansen, *Nonradial oscillations of neutron stars*, *Astrophys. J.* **325** (1988) 725.
- [15] A. Kunjipurayil, T. Zhao, B. Kumar, B.K. Agrawal and M. Prakash, *Impact of the equation of state on f- and p-mode oscillations of neutron stars*, *Phys. Rev. D* **106** (2022) 063005 [[arXiv:2205.02081](#)] [INSPIRE].
- [16] H.C. Das, A. Kumar, S.K. Biswal and S.K. Patra, *Impacts of dark matter on the f-mode oscillation of hyperon star*, *Phys. Rev. D* **104** (2021) 123006 [[arXiv:2109.01851](#)] [INSPIRE].
- [17] P. Routaray et al., *Investigating dark matter-admixed neutron stars with NITR equation of state in light of PSR J0952-0607*, *JCAP* **10** (2023) 073 [[arXiv:2304.05100](#)] [INSPIRE].
- [18] T. Zhao and J.M. Lattimer, *Universal relations for neutron star f-mode and g-mode oscillations*, *Phys. Rev. D* **106** (2022) 123002 [[arXiv:2204.03037](#)] [INSPIRE].
- [19] B.K. Pradhan and D. Chatterjee, *Effect of hyperons on f-mode oscillations in Neutron Stars*, *Phys. Rev. C* **103** (2021) 035810 [[arXiv:2011.02204](#)] [INSPIRE].
- [20] B.K. Pradhan, D. Chatterjee, M. Lanoye and P. Jaikumar, *General relativistic treatment of f-mode oscillations of hyperonic stars*, *Phys. Rev. C* **106** (2022) 015805 [[arXiv:2203.03141](#)] [INSPIRE].
- [21] H. Sotani and B. Kumar, *Universal relations between the quasinormal modes of neutron star and tidal deformability*, *Phys. Rev. D* **104** (2021) 123002 [[arXiv:2109.08145](#)] [INSPIRE].

- [22] L.S. Finn, *g-modes in zero-temperature neutron stars*, *Mon. Not. Roy. Astron. Soc.* **227** (1987) 265.
- [23] A. Reisenegger and P. Goldreich, *A new class of g-modes in neutron stars*, *Astrophys. J.* **395** (1992) 240.
- [24] T. Zhao, C. Constantinou, P. Jaikumar and M. Prakash, *Quasinormal g modes of neutron stars with quarks*, *Phys. Rev. D* **105** (2022) 103025 [[arXiv:2202.01403](#)] [[INSPIRE](#)].
- [25] N. Lozano, V. Tran and P. Jaikumar, *Temperature Effects on Core g-Modes of Neutron Stars*, *Galaxies* **10** (2022) 79 [[arXiv:2207.13488](#)] [[INSPIRE](#)].
- [26] C. Constantinou, S. Han, P. Jaikumar and M. Prakash, *g modes of neutron stars with hadron-to-quark crossover transitions*, *Phys. Rev. D* **104** (2021) 123032 [[arXiv:2109.14091](#)] [[INSPIRE](#)].
- [27] P. Jaikumar, A. Semposki, M. Prakash and C. Constantinou, *g-mode oscillations in hybrid stars: A tale of two sounds*, *Phys. Rev. D* **103** (2021) 123009 [[arXiv:2101.06349](#)] [[INSPIRE](#)].
- [28] W. Wei, M. Salinas, T. Klähn, P. Jaikumar and M. Barry, *Lifting the Veil on Quark Matter in Compact Stars with Core g-mode Oscillations*, *Astrophys. J.* **904** (2020) 187 [[arXiv:1811.11377](#)] [[INSPIRE](#)].
- [29] V. Tran, S. Ghosh, N. Lozano, D. Chatterjee and P. Jaikumar, *g-mode oscillations in neutron stars with hyperons*, *Phys. Rev. C* **108** (2023) 015803 [[arXiv:2212.09875](#)] [[INSPIRE](#)].
- [30] B. Haskell, K. Glampedakis and N. Andersson, *A new mechanism for saturating unstable r-modes in neutron stars*, *Mon. Not. Roy. Astron. Soc.* **441** (2014) 1662 [[arXiv:1307.0985](#)] [[INSPIRE](#)].
- [31] B. Haskell, *R-modes in neutron stars: Theory and observations*, *Int. J. Mod. Phys. E* **24** (2015) 1541007 [[arXiv:1509.04370](#)] [[INSPIRE](#)].
- [32] O.P. Jyothilakshmi, P.E.S. Krishnan, P. Thakur, V. Sreekanth and T.K. Jha, *Hyperon bulk viscosity and r-modes of neutron stars*, *Mon. Not. Roy. Astron. Soc.* **516** (2022) 3381 [[arXiv:2208.14436](#)] [[INSPIRE](#)].
- [33] L.-M. Lin, *Numerical study of nonlinear R-modes in neutron stars*, Ph.D. Thesis, Washington University in St. Louis, Saint Louis, MO, U.S.A. (2004).
- [34] L. Rezzolla, *The r-modes Oscillations and Instability: Surprises from Magnetized Neutron Stars*, in the proceedings of the *14th SIGRAV Congress on General Relativity and Gravitation (SIGRAV 2000)*, Genova, Italy, 18–22 September 2000, Springer, Milano, Italy (2002), pp. 235–248 [[DOI:10.1007/978-88-470-2101-3_16](#)] [[INSPIRE](#)].
- [35] M. Jasiulek and C. Chirenti, *R-mode frequencies of rapidly and differentially rotating relativistic neutron stars*, *Phys. Rev. D* **95** (2017) 064060 [[arXiv:1611.07924](#)] [[INSPIRE](#)].
- [36] O. Benhar, E. Berti and V. Ferrari, *The Imprint of the equation of state on the axial w modes of oscillating neutron stars*, *Mon. Not. Roy. Astron. Soc.* **310** (1999) 797 [[gr-qc/9901037](#)] [[INSPIRE](#)].
- [37] D. Bandyopadhyay and D. Chatterjee, *Axial W-modes of neutron stars with exotic matter*, in the proceedings of the *12th Marcel Grossmann Meeting on General Relativity*, Paris, France, 12–18 July 2009, World Scientific (2012), pp. 949–951 [[DOI:10.1142/9789814374552_0102](#)] [[INSPIRE](#)].
- [38] K.D. Kokkotas and B.G. Schmidt, *Quasinormal modes of stars and black holes*, *Living Rev. Rel.* **2** (1999) 2 [[gr-qc/9909058](#)] [[INSPIRE](#)].

- [39] H. Sotani, N. Yasutake, T. Maruyama and T. Tatsumi, *Signatures of hadron-quark mixed phase in gravitational waves*, *Phys. Rev. D* **83** (2011) 024014 [[arXiv:1012.4042](#)] [[INSPIRE](#)].
- [40] C.V. Flores and G. Lugones, *Discriminating hadronic and quark stars through gravitational waves of fluid pulsation modes*, *Class. Quant. Grav.* **31** (2014) 155002 [[arXiv:1310.0554](#)] [[INSPIRE](#)].
- [41] I.F. Ranea-Sandoval, O.M. Guilerá, M. Mariani and M.G. Orsaria, *Oscillation modes of hybrid stars within the relativistic Cowling approximation*, *JCAP* **12** (2018) 031 [[arXiv:1807.02166](#)] [[INSPIRE](#)].
- [42] S. Shibagaki, T. Kuroda, K. Kotake and T. Takiwaki, *A New Gravitational Wave Signature of Low- $T/|W|$ Instability in Rapidly Rotating Stellar Core Collapse*, *Mon. Not. Roy. Astron. Soc. Lett.* **493** (2020) L138 [[arXiv:1909.09730](#)] [[INSPIRE](#)].
- [43] N. Andersson and K.D. Kokkotas, *Towards gravitational wave asteroseismology*, *Mon. Not. Roy. Astron. Soc.* **299** (1998) 1059 [[gr-qc/9711088](#)] [[INSPIRE](#)].
- [44] H.K. Lau, P.T. Leung and L.M. Lin, *Inferring physical parameters of compact stars from their f -mode gravitational wave signals*, *Astrophys. J.* **714** (2010) 1234 [[arXiv:0911.0131](#)] [[INSPIRE](#)].
- [45] T.K. Chan, Y.-H. Sham, P.T. Leung and L.-M. Lin, *Multipolar universal relations between f -mode frequency and tidal deformability of compact stars*, *Phys. Rev. D* **90** (2014) 124023 [[arXiv:1408.3789](#)] [[INSPIRE](#)].
- [46] B.K. Pradhan, A. Vijaykumar and D. Chatterjee, *Impact of updated multipole Love numbers and f -Love universal relations in the context of binary neutron stars*, *Phys. Rev. D* **107** (2023) 023010 [[arXiv:2210.09425](#)] [[INSPIRE](#)].
- [47] K. Yagi and N. Yunes, *I -Love- Q Relations in Neutron Stars and their Applications to Astrophysics, Gravitational Waves and Fundamental Physics*, *Phys. Rev. D* **88** (2013) 023009 [[arXiv:1303.1528](#)] [[INSPIRE](#)].
- [48] K. Yagi and N. Yunes, *I -Love- Q anisotropically: Universal relations for compact stars with scalar pressure anisotropy*, *Phys. Rev. D* **91** (2015) 123008 [[arXiv:1503.02726](#)] [[INSPIRE](#)].
- [49] C. Breu and L. Rezzolla, *Maximum mass, moment of inertia and compactness of relativistic stars*, *Mon. Not. Roy. Astron. Soc.* **459** (2016) 646 [[arXiv:1601.06083](#)] [[INSPIRE](#)].
- [50] L. Rezzolla, P. Pizzochero, D. Jones, N. Rea and I. Vidaña, *The Physics and Astrophysics of Neutron Stars*, in *Astrophysics and Space Science Library*, Springer International Publishing (2019).
- [51] R. Riahi, S.Z. Kalantari and J.A. Rueda Hernandez, *Universal relations for the Keplerian sequence of rotating neutron stars*, *Phys. Rev. D* **99** (2019) 043004 [[arXiv:1902.00349](#)] [[INSPIRE](#)].
- [52] T. Gupta, B. Majumder, K. Yagi and N. Yunes, *I -Love- Q Relations for Neutron Stars in dynamical Chern Simons Gravity*, *Class. Quant. Grav.* **35** (2018) 025009 [[arXiv:1710.07862](#)] [[INSPIRE](#)].
- [53] J.-L. Jiang, S.-P. Tang, Y.-Z. Wang et al., *PSR J0030+0451, GW170817 and the nuclear data: joint constraints on equation of state and bulk properties of neutron stars*, *Astrophys. J.* **892** (2020) 55 [[arXiv:1912.07467](#)] [[INSPIRE](#)].
- [54] C.-H. Yeung, L.-M. Lin, N. Andersson and G. Comer, *The I -Love- Q Relations for Superfluid Neutron Stars*, *Universe* **7** (2021) 111 [[arXiv:2105.00798](#)] [[INSPIRE](#)].

- [55] S. Chakrabarti, T. Delsate, N. G rlebeck and J. Steinhoff, *I-Q relation for rapidly rotating neutron stars*, *Phys. Rev. Lett.* **112** (2014) 201102 [[arXiv:1311.6509](#)] [[INSPIRE](#)].
- [56] B. Haskell, R. Ciolfi, F. Pannarale and L. Rezzolla, *On the universality of I-Love-Q relations in magnetized neutron stars*, *Mon. Not. Roy. Astron. Soc. Lett.* **438** (2014) L71 [[arXiv:1309.3885](#)] [[INSPIRE](#)].
- [57] D. Bandyopadhyay, S.A. Bhat, P. Char and D. Chatterjee, *Moment of inertia, quadrupole moment, Love number of neutron star and their relations with strange matter equations of state*, *Eur. Phys. J. A* **54** (2018) 26 [[arXiv:1712.01715](#)] [[INSPIRE](#)].
- [58] S.S. Yazadjiev, *Relativistic models of magnetars: Nonperturbative analytical approach*, *Phys. Rev. D* **85** (2012) 044030 [[arXiv:1111.3536](#)] [[INSPIRE](#)].
- [59] C.Y. Cardall, M. Prakash and J.M. Lattimer, *Effects of strong magnetic fields on neutron star structure*, *Astrophys. J.* **554** (2001) 322 [[astro-ph/0011148](#)] [[INSPIRE](#)].
- [60] K. Ioka and M. Sasaki, *Relativistic stars with poloidal and toroidal magnetic fields and meridional flow*, *Astrophys. J.* **600** (2004) 296 [[astro-ph/0305352](#)] [[INSPIRE](#)].
- [61] R. Ciolfi and L. Rezzolla, *Twisted-torus configurations with large toroidal magnetic fields in relativistic stars*, *Mon. Not. Roy. Astron. Soc. Lett.* **435** (2013) L43 [[arXiv:1306.2803](#)] [[INSPIRE](#)].
- [62] R. Ciolfi, V. Ferrari and L. Gualtieri, *Structure and deformations of strongly magnetized neutron stars with twisted torus configurations*, *Mon. Not. Roy. Astron. Soc.* **406** (2010) 2540 [[arXiv:1003.2148](#)] [[INSPIRE](#)].
- [63] J. Friebe and L. Rezzolla, *Equilibrium models of relativistic stars with a toroidal magnetic field*, *Mon. Not. Roy. Astron. Soc.* **427** (2012) 3406 [[arXiv:1207.4035](#)] [[INSPIRE](#)].
- [64] A.G. Pili, N. Bucciantini and L. Del Zanna, *Axisymmetric equilibrium models for magnetized neutron stars in General Relativity under the Conformally Flat Condition*, *Mon. Not. Roy. Astron. Soc.* **439** (2014) 3541 [[arXiv:1401.4308](#)] [[INSPIRE](#)].
- [65] N. Bucciantini, A.G. Pili and L. Del Zanna, *The role of currents distribution in general relativistic equilibria of magnetized neutron stars*, *Mon. Not. Roy. Astron. Soc.* **447** (2015) 3278 [[arXiv:1412.5347](#)] [[INSPIRE](#)].
- [66] R.F. Sawyer, *Condensed π^- phase in neutron star matter*, *Phys. Rev. Lett.* **29** (1972) 382 [[INSPIRE](#)].
- [67] B. Carter and D. Langlois, *Relativistic models for superconducting superfluid mixtures*, *Nucl. Phys. B* **531** (1998) 478 [[gr-qc/9806024](#)] [[INSPIRE](#)].
- [68] V. Canuto, *Equation of State at Ultrahigh Densities. Part 1*, *Ann. Rev. Astron. Astrophys.* **12** (1974) 167 [[INSPIRE](#)].
- [69] M. Ruderman, *Pulsars: structure and dynamics*, *Ann. Rev. Astron. Astrophys.* **10** (1972) 427 [[INSPIRE](#)].
- [70] S.G. Nelmes and B.M.A.G. Piette, *Phase Transition and Anisotropic Deformations of Neutron Star Matter*, *Phys. Rev. D* **85** (2012) 123004 [[arXiv:1204.0910](#)] [[INSPIRE](#)].
- [71] R. Kippenhahn, A. Weigert and A. Weiss, *Stellar structure and evolution*, in *Astronomy and Astrophysics Library*, Springer (2012) [[DOI:10.1007/978-3-642-30304-3](#)] [[INSPIRE](#)].
- [72] N.K. Glendenning, *Compact Stars*, in *Astronomy and Astrophysics Library*, Springer, New York, NY, U.S.A. (1997) [[DOI:10.1007/978-1-4684-0491-3](#)].

- [73] H. Heiselberg and M. Hjorth-Jensen, *Phases of dense matter in neutron stars*, *Phys. Rep.* **328** (2000) 237 [[nucl-th/9902033](#)] [[INSPIRE](#)].
- [74] R.L. Bowers and E.P.T. Liang, *Anisotropic Spheres in General Relativity*, *Astrophys. J.* **188** (1974) 657 [[INSPIRE](#)].
- [75] D. Horvat, S. Ilić and A. Marunović, *Radial pulsations and stability of anisotropic stars with quasi-local equation of state*, *Class. Quant. Grav.* **28** (2011) 025009 [[arXiv:1010.0878](#)] [[INSPIRE](#)].
- [76] M. Cosenza, L. Herrera, M. Esculpi and L. Witten, *Some models of anisotropic spheres in general relativity*, *J. Math. Phys.* **22** (1981) 118 [[INSPIRE](#)].
- [77] H.O. Silva, C.F.B. Macedo, E. Berti and L.C.B. Crispino, *Slowly rotating anisotropic neutron stars in general relativity and scalar-tensor theory*, *Class. Quant. Grav.* **32** (2015) 145008 [[arXiv:1411.6286](#)] [[INSPIRE](#)].
- [78] W. Hillebrandt and K.O. Steinmetz, *Anisotropic neutron star models: stability against radial and nonradial pulsations*, *Astron. Astrophys.* **53** (1976) 283.
- [79] D.D. Doneva and S.S. Yazadjiev, *Gravitational wave spectrum of anisotropic neutron stars in Cowling approximation*, *Phys. Rev. D* **85** (2012) 124023 [[arXiv:1203.3963](#)] [[INSPIRE](#)].
- [80] S.S. Bayin, *Anisotropic Fluid Spheres in General Relativity*, *Phys. Rev. D* **26** (1982) 1262 [[INSPIRE](#)].
- [81] Z. Roupas, *Secondary component of gravitational-wave signal GW190814 as an anisotropic neutron star*, *Astrophys. Space Sci.* **366** (2021) 9 [[arXiv:2007.10679](#)] [[INSPIRE](#)].
- [82] D. Deb, B. Mukhopadhyay and F. Weber, *Effects of Anisotropy on Strongly Magnetized Neutron and Strange Quark Stars in General Relativity*, *Astrophys. J.* **922** (2021) 149 [[arXiv:2108.12436](#)] [[INSPIRE](#)].
- [83] G. Estevez-Delgado and J. Estevez-Delgado, *On the effect of anisotropy on stellar models*, *Eur. Phys. J. C* **78** (2018) 673 [[INSPIRE](#)].
- [84] M.L. Pattersons and A. Sulaksono, *Mass correction and deformation of slowly rotating anisotropic neutron stars based on Hartle-Thorne formalism*, *Eur. Phys. J. C* **81** (2021) 698 [[INSPIRE](#)].
- [85] R. Rizaldy, A.R. Alfarasyi, A. Sulaksono and T. Sumaryada, *Neutron-star deformation due to anisotropic momentum distribution of neutron-star matter*, *Phys. Rev. C* **100** (2019) 055804 [[INSPIRE](#)].
- [86] A. Rahmansyah, A. Sulaksono, A.B. Wahidin and A.M. Setiawan, *Anisotropic neutron stars with hyperons: implication of the recent nuclear matter data and observations of neutron stars*, *Eur. Phys. J. C* **80** (2020) 769 [[INSPIRE](#)].
- [87] A. Rahmansyah and A. Sulaksono, *Recent multimessenger constraints and the anisotropic neutron star*, *Phys. Rev. C* **104** (2021) 065805 [[INSPIRE](#)].
- [88] L. Herrera, J. Ospino and A. Di Prisco, *All static spherically symmetric anisotropic solutions of Einstein's equations*, *Phys. Rev. D* **77** (2008) 027502 [[arXiv:0712.0713](#)] [[INSPIRE](#)].
- [89] L. Herrera and W. Barreto, *General relativistic polytropes for anisotropic matter: The general formalism and applications*, *Phys. Rev. D* **88** (2013) 084022 [[arXiv:1310.1114](#)] [[INSPIRE](#)].
- [90] B. Biswas and S. Bose, *Tidal deformability of an anisotropic compact star: Implications of GW170817*, *Phys. Rev. D* **99** (2019) 104002 [[arXiv:1903.04956](#)] [[INSPIRE](#)].

- [91] S. Das, B. Parida, S. Ray and S. Pal, *Role of Anisotropy on the Tidal Deformability of Compact Stellar Objects*, in the proceedings of the *The 1st Electronic Conference on Universe*, online conference, 22–28 February 2021, *Phys. Sci. Forum* **2** (2021) 29.
- [92] Z. Roupas and G.G.L. Nashed, *Anisotropic neutron stars modelling: constraints in Krori-Barua spacetime*, *Eur. Phys. J. C* **80** (2020) 905 [[arXiv:2007.09797](#)] [[INSPIRE](#)].
- [93] A. Sulaksono, *Anisotropic pressure and hyperons in neutron stars*, *Int. J. Mod. Phys. E* **24** (2015) 1550007 [[arXiv:1412.7247](#)] [[INSPIRE](#)].
- [94] A.M. Setiawan and A. Sulaksono, *Anisotropic neutron stars and perfect fluid’s energy conditions*, *Eur. Phys. J. C* **79** (2019) 755 [[INSPIRE](#)].
- [95] H.C. Das, *I-Love-C relation for an anisotropic neutron star*, *Phys. Rev. D* **106** (2022) 103518 [[arXiv:2208.12566](#)] [[INSPIRE](#)].
- [96] M. Fortin, C. Providência, A.R. Raduta et al., *Neutron star radii and crusts: uncertainties and unified equations of state*, *Phys. Rev. C* **94** (2016) 035804 [[arXiv:1604.01944](#)] [[INSPIRE](#)].
- [97] B. Kumar and P. Landry, *Inferring neutron star properties from GW170817 with universal relations*, *Phys. Rev. D* **99** (2019) 123026 [[arXiv:1902.04557](#)] [[INSPIRE](#)].
- [98] P. Landry and B. Kumar, *Constraints on the moment of inertia of PSR J0737-3039A from GW170817*, *Astrophys. J. Lett.* **868** (2018) L22 [[arXiv:1807.04727](#)] [[INSPIRE](#)].
- [99] T. Malik, N. Alam, M. Fortin et al., *GW170817: constraining the nuclear matter equation of state from the neutron star tidal deformability*, *Phys. Rev. C* **98** (2018) 035804 [[arXiv:1805.11963](#)] [[INSPIRE](#)].
- [100] N. Alam, B.K. Agrawal, M. Fortin et al., *Strong correlations of neutron star radii with the slopes of nuclear matter incompressibility and symmetry energy at saturation*, *Phys. Rev. C* **94** (2016) 052801 [[arXiv:1610.06344](#)] [[INSPIRE](#)].
- [101] V. Parmar, H.C. Das, A. Kumar et al., *Crustal properties of a neutron star within an effective relativistic mean-field model*, *Phys. Rev. D* **105** (2022) 043017 [[arXiv:2111.07278](#)] [[INSPIRE](#)].
- [102] V. Parmar, H.C. Das, A. Kumar et al., *Pasta properties of the neutron star within effective relativistic mean-field model*, *Phys. Rev. D* **106** (2022) 023031 [[arXiv:2203.16827](#)] [[INSPIRE](#)].
- [103] R.M. Wald, *General Relativity*, Chicago University Press, Chicago, U.S.A. (1984) [[DOI:10.7208/chicago/9780226870373.001.0001](#)] [[INSPIRE](#)].
- [104] K. Schwarzschild, *On the gravitational field of a mass point according to Einstein’s theory*, *Sitzungsber. Preuss. Akad. Wiss. Berlin* **1916** (1916) 189 [[physics/9905030](#)] [[INSPIRE](#)].
- [105] A. Einstein, *The Field Equations of Gravitation*, *Sitzungsber. Preuss. Akad. Wiss. Berlin* **1915** (1915) 844.
- [106] P. Bhar and P. Rej, *Compact stellar model in the presence of pressure anisotropy in modified Finch Skea space-time*, *J. Astrophys. Astron.* **42** (2021) 74 [[arXiv:2106.01316](#)] [[INSPIRE](#)].
- [107] J.R. Oppenheimer and G.M. Volkoff, *On massive neutron cores*, *Phys. Rev.* **55** (1939) 374 [[INSPIRE](#)].
- [108] M.C. Miller et al., *PSR J0030+0451 Mass and Radius from NICER Data and Implications for the Properties of Neutron Star Matter*, *Astrophys. J. Lett.* **887** (2019) L24 [[arXiv:1912.05705](#)] [[INSPIRE](#)].
- [109] T.E. Riley et al., *A NICER View of PSR J0030+0451: Millisecond Pulsar Parameter Estimation*, *Astrophys. J. Lett.* **887** (2019) L21 [[arXiv:1912.05702](#)] [[INSPIRE](#)].

- [110] M.C. Miller et al., *The Radius of PSR J0740+6620 from NICER and XMM-Newton Data*, *Astrophys. J. Lett.* **918** (2021) L28 [[arXiv:2105.06979](#)] [[INSPIRE](#)].
- [111] LIGO SCIENTIFIC and VIRGO collaborations, *GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral*, *Phys. Rev. Lett.* **119** (2017) 161101 [[arXiv:1710.05832](#)] [[INSPIRE](#)].
- [112] LIGO SCIENTIFIC and VIRGO collaborations, *GW190814: Gravitational Waves from the Coalescence of a 23 Solar Mass Black Hole with a 2.6 Solar Mass Compact Object*, *Astrophys. J. Lett.* **896** (2020) L44 [[arXiv:2006.12611](#)] [[INSPIRE](#)].
- [113] A. Worley, P.G. Krastev and B.-A. Li, *Nuclear constraints on the momenta of inertia of neutron stars*, *Astrophys. J.* **685** (2008) 390 [[arXiv:0801.1653](#)] [[INSPIRE](#)].
- [114] T. Hinderer, *Tidal Love numbers of neutron stars*, *Astrophys. J.* **677** (2008) 1216 [[arXiv:0711.2420](#)] [[INSPIRE](#)].
- [115] T. Hinderer, *Erratum: Tidal love numbers of neutron stars*, *Astrophys. J.* **697** (2009) 964.
- [116] LIGO SCIENTIFIC and VIRGO collaborations, *GW170817: Measurements of neutron star radii and equation of state*, *Phys. Rev. Lett.* **121** (2018) 161101 [[arXiv:1805.11581](#)] [[INSPIRE](#)].
- [117] T.G. Cowling, *The Non-radial Oscillations of Polytropic Stars*, *Mon. Not. Roy. Astron. Soc.* **101** (1941) 367 [[INSPIRE](#)].
- [118] E.J.A. Curi, L.B. Castro, C.V. Flores and C.H. Lenzi, *Non-radial oscillations and global stellar properties of anisotropic compact stars using realistic equations of state*, *Eur. Phys. J. C* **82** (2022) 527 [[arXiv:2206.09260](#)] [[INSPIRE](#)].
- [119] N. Jiang and K. Yagi, *Analytic I-Love-C relations for realistic neutron stars*, *Phys. Rev. D* **101** (2020) 124006 [[arXiv:2003.10498](#)] [[INSPIRE](#)].
- [120] C. Chirenti, G.H. de Souza and W. Kastaun, *Fundamental oscillation modes of neutron stars: validity of universal relations*, *Phys. Rev. D* **91** (2015) 044034 [[arXiv:1501.02970](#)] [[INSPIRE](#)].
- [121] K. Yagi and N. Yunes, *I-Love-Q*, *Science* **341** (2013) 365 [[arXiv:1302.4499](#)] [[INSPIRE](#)].
- [122] K.V. Staykov, D.D. Doneva and S.S. Yazadjiev, *Moment-of-inertia-compactness universal relations in scalar-tensor theories and \mathcal{R}^2 gravity*, *Phys. Rev. D* **93** (2016) 084010 [[arXiv:1602.00504](#)] [[INSPIRE](#)].
- [123] S.R. Mohanty, S. Ghosh and B. Kumar, *Unstable Anisotropic Neutron Stars: Probing the Limits of Gravitational Collapse*, [arXiv:2304.02439](#) [[INSPIRE](#)].
- [124] D.G. Ravenhall and C.J. Pethick, *Neutron star moments of inertia*, *Astrophys. J.* **424** (1994) 846.
- [125] J.M. Lattimer and B.F. Schutz, *Constraining the equation of state with moment of inertia measurements*, *Astrophys. J.* **629** (2005) 979 [[astro-ph/0411470](#)] [[INSPIRE](#)].
- [126] D. Popchev, K.V. Staykov, D.D. Doneva and S.S. Yazadjiev, *Moment of inertia-mass universal relations for neutron stars in scalar-tensor theory with self-interacting massive scalar field*, *Eur. Phys. J. C* **79** (2019) 178 [[arXiv:1812.00347](#)] [[INSPIRE](#)].
- [127] M. Bejger and P. Haensel, *Moments of inertia for neutron and strange stars: Limits derived for the Crab pulsar*, *Astron. Astrophys.* **396** (2002) 917 [[astro-ph/0209151](#)] [[INSPIRE](#)].
- [128] H. Sotani, K. Tominaga and K.-i. Maeda, *Density discontinuity of a neutron star and gravitational waves*, *Phys. Rev. D* **65** (2001) 024010 [[gr-qc/0108060](#)] [[INSPIRE](#)].