

Epinephrine

1:1,000 · 1 mg/mL

SYMPATHOMIMETIC

CATECHOLAMINE

MECHANISM OF ACTION

a-1 -> Vasoconstriction

b-1 -> ^ HR & contractility

b-2 -> Bronchodilation

INDICATIONS

- Anaphylaxis
- Severe Allergic Rxn
- Refractory Bronchospasm
- Asthma (refractory)

CONTRAINDICATIONS

- ◆ Hypersensitivity
- ◆ Hypovolemic Shock
- ◆ Coronary insuff. **REL**
- ◆ Hypertension **REL**

DOSE & ROUTE

ADULT

0.3-0.5 mg IM

Lateral thigh

Repeat every 5-15 min

PEDIATRIC

0.01 mg/kg IM

Max 0.3 mg

Lateral thigh

ASTHMA 0.3-0.5 mg SQ

Onset 1-2 min · Duration 5-15 min

ADVERSE EFFECTS

Tachycardia

Hypertension

Dysrhythmias

Anxiety/Tremor

Pallor

Headache

Angina

PRECAUTIONS

Use caution in elderly, known cardiac disease, HTN, and diabetes. May precipitate angina or MI in susceptible patients.

Etomidate

2 mg/mL

ANESTHETIC

SEDATIVE

MECHANISM OF ACTION

Enhances **GABA-A** activity -> neuronal inhibition

-> **rapid sedation & unconsciousness**

INDICATIONS

- Procedural Sedation
- RSI Induction

CONTRAINDICATIONS

- ◆ Hypersensitivity
- ◆ Labor & Delivery

DOSE & ROUTE

ADULT

0.2-0.6 mg/kg IV/IO

Slow bolus + NS flush

Max 40 mg

PEDIATRIC

0.1-0.4 mg/kg IV/IO

Slow bolus + NS flush

Max 20 mg

Onset 30-60 sec · Duration 5-10 min

ADVERSE EFFECTS

Apnea

Myoclonus

Nausea/Vomiting

Adrenal Suppr.

Resp. Depression

BP Changes

PRECAUTIONS

Use caution in adrenal insufficiency - may decrease cortisol. May cause transient hypotension in unstable BP or cardiac patients. Lower dose in elderly.