

Epinephrine

1:1,000 · 1 mg/mL

SYMPATHOMIMETIC

CATECHOLAMINE

MECHANISM OF ACTION

α-1 → Vasoconstriction

β-1 → ↑ HR & contractility

β-2 → Bronchodilation

INDICATIONS

- Anaphylaxis
- Severe Allergic Reaction
- Refractory Bronchospasm
- Asthma (refractory)

CONTRAINDICATIONS

- ◆ Hypersensitivity
- ◆ Hypovolemic Shock
- ◆ Coronary insuff.
RELATIVE
- ◆ Hypertension **RELATIVE**

DOSE & ROUTE

ADULT

0.3–0.5 mg IM

Lateral thigh

Repeat every 5–15 min

PEDIATRIC

0.01 mg/kg IM

Max 0.3 mg

Lateral thigh

ASTHMA 0.3–0.5 mg SQ

Onset **1–2 min** · Duration **5–15 min**

ADVERSE EFFECTS

Tachycardia

Hypertension

Dysrhythmias

Anxiety / Tremor

Pallor / Headache

Angina

PRECAUTIONS

Use caution in **elderly**, known **cardiac disease**, **HTN**, and **diabetes**. May precipitate angina or MI in susceptible patients.

Etomidate

2 mg/mL

ANESTHETIC

SEDATIVE

MECHANISM OF ACTION

Enhances **GABA-A** activity → neuronal inhibition → **rapid sedation & unconsciousness**

INDICATIONS

- Procedural Sedation
- RSI Induction

CONTRAINDICATIONS

- ◆ Hypersensitivity
- ◆ Labor & Delivery

DOSE & ROUTE

ADULT

0.2–0.6 mg/kg IV/IO

Slow bolus + NS flush
Max 40 mg

PEDIATRIC

0.1–0.4 mg/kg IV/IO

Slow bolus + NS flush
Max 20 mg

Onset **30–60 sec** · Duration **5–10 min**

ADVERSE EFFECTS

- Apnea
- Myoclonus
- Nausea / Vomiting
- Adrenal Suppression
- Resp. Depression
- BP Changes

PRECAUTIONS

Use caution in **adrenal insufficiency** — may ↓ cortisol. May cause transient **hypotension** in unstable BP or **cardiac** patients. Lower dose in **elderly**.