

The Principles of Quantum Mechanics & Superposition States

Authors: P. A. M. Dirac, Niels Bohr

Published in: Oxford University Press (1930) | Archival ID: QV-QC-2

1. ABSTRACT & THEORETICAL FOUNDATION

We formulate the fundamental principle of linear quantum superposition of state vectors in Hilbert space, proving that physical observables represent Hermitian operators whose eigenvalues correspond to allowable measurement outcomes.

2. CORE CONCEPTUAL INTUITION (PLAIN ENGLISH)

"Imagine a musical chord: a single piano plays multiple notes at the same exact time. Superposition lets a quantum computer evaluate multiple possibilities concurrently."

3. Mathematical Formulation & Unitary Rule

$$|\psi\rangle = \sum_{x \in \{0,1\}^n} c_x |x\rangle, \quad \text{quad} \quad \sum_x |c_x|^2 = 1$$

For an n -qubit system, superposition allows simultaneous storage and linear manipulation of all 2^n configurations with unit Euclidean norm.

4. Step-by-Step Circuit Sequence

Applying $H^{\otimes n}$ across all n qubit lines initializes parallel state space.

5. Executable IBM Qiskit (Python) Code

```
from qiskit import QuantumCircuit

# 3 qubits in uniform superposition
qc = QuantumCircuit(3)
qc.h([0, 1, 2]) # Applies H gate to all 3 qubits

# State vector now contains 8 equal amplitudes: 1/sqrt(8)
print(qc.draw(output='text'))
```

6. Computational Complexity & Speedup

Classical Time:

Quantum Time:

Speedup Class:

$O(2^n)$ to evaluate 2^n distinct inputs sequentially $\rightarrow$ $O(1)$ gate depth $\rightarrow$ $O(2^n)$ to read out $\rightarrow$ Foundational

7. Key Applications & Future Research Directions

- Initial state preparation for quantum search and factoring
- Quantum Monte Carlo integration speedups
- Interferometric precision phase measurement

Future Work:

Investigating macroscopic superposition limits and continuous-variable state representations for optical quantum processors.