Alexey is playing with an array, A, of n integers. His friend, Ivan, asks him to calculate the sum of the maximum values for all subsegments of \boldsymbol{A} . More formally, he wants Alexey to find

$$F(A) = \sum_{l=1}^{n} \sum_{r=l}^{n} \max_{l \le x \le r} A[x].$$

Alexey solved Ivan's challenge faster than expected, so Ivan decides to add another layer of difficulty by having Alexey answer m queries. The i^{th} query contains subsegment $[L_i,R_i]$, and he must calculate the sum of maximum values on all subsegments inside subsegment $[L_i, R_i]$.

More formally, for each query *i*, Alexey must calculate the following function:

$$F(A,L_i,R_i) = \sum\limits_{l=L_i}^{R_i} \sum\limits_{r=l}^{R_i} \max\limits_{l \leq x \leq r} A[x].$$

Can you help Alexey solve this problem?

Input Format

The first line contains 2 space-separated positive integers, n (the length of array A) and m (number of queries), respectively.

The second line contains n space-separated integers, $a_0, a_1, \ldots, a_{n-1}$ describing each element a_i (where $0 \le j < n$) in array A.

Each of the m subsequent lines contains 2 space-separated positive integers describing the respective values for L_i and R_i in query i (where $0 \le i < m$).

Constraints

- $egin{array}{ll} \bullet & 1 \leq n, m \leq 135000 \ \bullet & -10^9 \leq a_i \leq 10^9 \ \bullet & 1 \leq L_i \leq R_i \leq n \end{array}$

Output Format

For each query i (where $0 \le i < m$), print its answer on a new line.

Sample Input

3 6

1 3 2

Sample Output

Explanation

The answer for the second query is shown below: $F(A,1,2) = \max_{1 \le x \le 1} A[x] + \max_{1 \le x \le 2} A[x] + \max_{1 \le x \le 2} A[x]$ =1+3+3=7

The answer for the third query is shown below:
$$F(A,1,3) = \max_{1 \leq x \leq 1} A[x] + \max_{1 \leq x \leq 2} A[x] + \max_{1 \leq x \leq 3} A[x] + \max_{2 \leq x \leq 2} A[x] + \max_{2 \leq x \leq 3} A[x] + \max_{3 \leq x \leq 3} A[x]$$