



3. Prevention or Delay of Diabetes and Associated Comorbidities: Standards of Care in Diabetes—2026

American Diabetes Association
Professional Practice Committee for
Diabetes*

Diabetes Care 2026;49(Suppl. 1):S50–S60 | <https://doi.org/10.2337/dc26-S003>

The American Diabetes Association (ADA) “Standards of Care in Diabetes” includes the ADA’s current clinical practice recommendations and is intended to provide the components of diabetes care, general treatment goals and guidelines, and tools to evaluate quality of care. Members of the ADA Professional Practice Committee for Diabetes, an interprofessional expert committee, are responsible for updating the Standards of Care annually, or more frequently as warranted. For a detailed description of ADA standards, statements, and reports, as well as the evidence-grading system for ADA’s clinical practice recommendations and a full list of Professional Practice Committee members, please refer to Introduction and Methodology. Readers who wish to comment on the Standards of Care are invited to do so at professional.diabetes.org/SOC.

For guidelines related to screening for increased risk for type 1 diabetes, prediabetes and type 2 diabetes, and other forms of diabetes, please refer to section 2, “Diagnosis and Classification of Diabetes.” For guidelines related to screening, diagnosis, and management of diabetes in children and adolescents, please refer to section 14, “Children and Adolescents.”

Recommendations

3.1 In people with prediabetes, monitor for the development of diabetes at least annually; modify frequency of testing based on individual risk assessment. **E**

3.2 In people with presymptomatic type 1 diabetes, monitor for disease progression using A1C approximately every 6 months and 75-g oral glucose tolerance test (i.e., fasting and 2-h plasma glucose) annually; modify frequency of monitoring and consider augmenting with other glycemic assessment tools such as continuous glucose monitoring metrics based on individual risk assessment incorporating age, number and type of autoantibodies, and glycemic metrics. **E**

Screening for prediabetes (which refers to hyperglycemia preceding the diagnosis of type 2 diabetes only) and type 2 diabetes risk through an assessment of risk factors (Table 2.5) or with an assessment tool, such as the American Diabetes Association risk test, which can be used by either a layperson or a health care professional (diabetes.org/diabetes-risk-test), is recommended to guide whether to perform a diagnostic test for prediabetes (Table 2.2) and type 2 diabetes (Table 2.1) (see section 2, “Diagnosis and Classification of Diabetes”). Testing high-risk individuals for prediabetes is warranted because the laboratory assessment is safe and reasonable

*A complete list of members of the American Diabetes Association Professional Practice Committee for Diabetes can be found at <https://doi.org/10.2337/dc26-SINT>.

Duality of interest information for each contributor is available at <https://doi.org/10.2337/dc26-SDIS>.

Suggested citation: American Diabetes Association Professional Practice Committee for Diabetes. 3. Prevention or delay of diabetes and associated comorbidities: Standards of Care in Diabetes—2026. *Diabetes Care* 2026;49(Suppl. 1):S50–S60

© 2025 by the American Diabetes Association. Readers may use this work for educational, noncommercial purposes if properly cited and unaltered. The version of record may be linked at <https://diabetesjournals.org/care>, but ADA permission is required to post this work on any third-party site or platform. This publication and its contents may not be reproduced, distributed, or used for text or data mining, machine learning, or similar technologies without prior written permission. Requests to reuse, adapt, or distribute this work may be sent to permissions@diabetes.org. More information is available at <https://diabetesjournals.org/journals/pages/license>.

in cost, and early detection of hyperglycemia affords substantial time to intervene and delay or prevent the onset of type 2 diabetes and/or its long-term complications. Indeed, once identified, several effective therapeutic approaches exist that can delay type 2 diabetes in those with prediabetes with an A1C 5.7–6.4% (39–47 mmol/mol), impaired glucose tolerance (IGT) on 75-g oral glucose tolerance test (OGTT), or impaired fasting glucose (IFG). The utility of screening with A1C for prediabetes and diabetes may be limited in the presence of certain hemoglobinopathies and conditions that affect red blood cell turnover (**Table 2.3**). See section 2, “Diagnosis and Classification of Diabetes,” and section 6, “Glycemic Goals, Hypoglycemia, and Hyperglycemic Crises,” for additional details on the appropriate use and limitations of A1C testing.

Three stages of type 1 diabetes have been defined, with symptomatic type 1 diabetes classified as stage 3 (**Table 2.4**). In individuals at risk for developing clinical type 1 diabetes, younger age of seroconversion (particularly under age 3 years), the total number of diabetes-related autoantibodies (1), and the development of autoantibodies against islet antigen 2 (IA-2) have all been associated with a more rapid progression to stage 3 type 1 diabetes. While continuous glucose monitoring can predict progression to stage 3 type 1 diabetes in children and adolescents with autoantibodies (2), OGTT-based metrics are more predictive of progression compared with continuous glucose monitoring (3). The decision to perform an OGTT may depend on such factors as eligibility and interest for stage-specific treatments, interest in participation in clinical research, OGTT availability, and the burden of testing. Consensus guidance provides expert recommendations on what should be monitored and how often in people with presymptomatic type 1 diabetes (4). Cost-effectiveness of population-based screening programs for type 1 diabetes has not been established (5,6), but such screening is being implemented outside the U.S. (7).

LIFESTYLE BEHAVIOR CHANGE FOR TYPE 2 DIABETES PREVENTION

Recommendations

3.3 Refer adults with overweight or obesity at high risk of type 2 diabetes to a diabetes prevention

program to achieve and maintain a weight reduction of at least 5–7% of initial body weight through a healthy reduced-calorie eating pattern and ≥ 150 min/week of moderate-intensity physical activity. **A**

3.4 Prescribe an evidence-based eating pattern (e.g., Mediterranean, low carbohydrate) to individuals with prediabetes to prevent type 2 diabetes. **B**

3.5 Offer diabetes prevention programs to adults at high risk for type 2 diabetes. **A** Diabetes prevention programs should be covered by third-party payors, and inconsistencies in access should be addressed. **E**

3.6 Based on individual preference, certified technology-assisted diabetes prevention programs through smartphones, web-based applications, and telehealth can be effective in preventing type 2 diabetes and should be considered. **B**

The Diabetes Prevention Program

Several major randomized controlled trials, including the Diabetes Prevention Program (DPP) (8), the Finnish Diabetes Prevention Study (DPS) (9), and the Da Qing Diabetes Prevention Study (Da Qing study) (10), demonstrated that lifestyle/behavioral intervention with an individualized reduced-calorie meal plan is highly effective in preventing or delaying type 2 diabetes and improving other cardiometabolic risk factors such as blood pressure, lipids, and markers of inflammation (11). The strongest evidence for diabetes prevention in the U.S. comes from the DPP trial (8). The DPP demonstrated that intensive lifestyle intervention could reduce the risk of incident type 2 diabetes by 58% over 3 years. Follow-up of three large trials of lifestyle intervention for diabetes prevention showed sustained reduction in the risk of progression to type 2 diabetes: 39% reduction at 30 years in the Da Qing study (12), 43% reduction at 7 years in the Finnish DPS (9), and 34% reduction at 10 years (13), 27% reduction at 15 years (14), and 24% reduction at 21 years in the U.S. Diabetes Prevention Program Outcomes Study (DPPOS) (15,16).

The DPP lifestyle intervention was a goal-based intervention. A complete description of the DPP and how to access it is available online at coveragetoolkit.org/

about-national-dpp/ndpp-overview/. Briefly, the DPP administered a 16-session structured core curriculum, completed within the first 6 months of the program, followed by an additional 6 months of a flexible maintenance program. The core curriculum included sessions on reducing calories, increasing physical activity, self-monitoring, maintaining healthy lifestyle behaviors (such as how to make healthy choices when eating out), and guidance on managing psychological, social, and motivational challenges (17).

Two major goals of the DPP were to achieve and maintain at least 7% weight loss and to obtain 150 min of moderate-intensity physical activity per week, such as brisk walking. Weight loss was the most important factor in reducing incident diabetes. Obtaining at least 150 min of physical activity per week, even without achieving the weight loss goal, reduced incidence of type 2 diabetes by 44% (18). Participants were encouraged to distribute physical activity throughout the week with a minimum frequency of three times per week and at least 10 min per session. A maximum of 75 min of strength training could be applied toward the total 150 min/week physical activity goal. The initial focus of the nutrition intervention was on reducing total fat intake rather than caloric restriction. However, the concepts of calorie balance and the need to restrict total calories and fat were also introduced. Overall, the DPP's lifestyle intervention was designed to produce a 700 kcal/week energy deficit (17).

The 7% weight loss goal was selected because it was feasible to achieve and maintain, hypothesized to reduce incident diabetes risk, and improved other cardiometabolic risk factors. Further analysis suggested greater benefit for diabetes prevention with at least 7–10% weight loss with lifestyle interventions (18). The recommended pace of weight loss was 1–2 lb/week.

While the DPP interventions were successful in preventing or delaying the onset of type 2 diabetes, major cardiovascular events were not significantly different 21 years after the DPP concluded, which may be due to the widespread use of statin, antihypertensive, and metformin therapy among all participants well after the conclusion of the study (19). However, the incidence of diabetes continued to be significantly

lower at 21 years of follow-up in both intensive lifestyle therapy and metformin groups, when compared with placebo, corresponding to increases in median diabetes-free survival of 3.5 years and 2.5 years, respectively (15). Importantly, the overall treatment effect was driven almost entirely by benefit seen during the initial DPP phase (15), underscoring the importance of early intervention and continued implementation of diabetes prevention efforts. Still, individuals can derive benefit from lifestyle interventions to prevent progression of prediabetes to diabetes irrespective of the duration of prediabetes prior to initiation of lifestyle intervention (20). Thus, there is clear potential benefit in diabetes prevention and minimal to no risk of harm with these interventions.

Nutrition

Nutrition counseling for weight loss in the DPP lifestyle intervention arm focused on reduction of total fat and calories (8,17,18). However, current evidence suggests there is no ideal percentage of calories from carbohydrate, protein, and fat to prevent diabetes; therefore, macronutrient distribution should be based on an individualized assessment of current eating patterns, preferences, and metabolic goals. A variety of eating patterns (21) can be appropriate for individuals with prediabetes or diabetes, including Mediterranean-style, plant-based (which may or may not be omnivorous), Dietary Approaches to Stop Hypertension (DASH), and low-carbohydrate (22–25). The overall quality of food consumed (as measured by the Healthy Eating Index, Alternative Healthy Eating Index, and DASH score), with an emphasis on whole grains, legumes, nuts, fruits, and vegetables and minimal refined and processed foods, is also associated with a lower risk of type 2 diabetes (26–28). For people at risk for or living with diabetes, individualized medical nutrition therapy (see section 5, “Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes,” for more detailed information) is effective in lowering A1C (29,30).

Physical Activity

Moderate-intensity physical activity, such as brisk walking for 150 min/week, has shown beneficial effects in those with prediabetes (8). Similarly, moderate-

intensity physical activity has been shown to improve insulin sensitivity and reduce abdominal fat in children, adolescents, and young adults (31,32). Health care professionals are encouraged to promote a DPP-style program to all individuals who have been identified to be at an increased risk of type 2 diabetes. In addition to aerobic activity, a physical activity plan designed to prevent diabetes can include resistance training (17,33). Breaking up prolonged sedentary time may also be encouraged, as it lowers postprandial glucose levels (34). The effects of regular physical activity participation appear to extend to the prevention of gestational diabetes mellitus (GDM) as well (35,36).

Delivery and Dissemination of Lifestyle Behavior Change for Diabetes Prevention

Because the intensive lifestyle intervention in the DPP was effective in preventing type 2 diabetes among those at high risk, and both DPP and adapted lifestyle behavior change programs for diabetes prevention were shown to be cost-effective across multiple applications, broader efforts to disseminate scalable DPP-like lifestyle behavior change programs for diabetes prevention with coverage by third-party payors ensued (37–42). Group delivery of DPP content in community or primary care settings has demonstrated the potential to reduce overall program costs while still producing weight loss and diabetes risk reduction (43,44). Interventions to prevent type 2 diabetes can also be delivered through workplace programming, with interventions that most closely adhere to DPP content shown to be most effective (45).

The Centers for Disease Control and Prevention (CDC) developed the National Diabetes Prevention Program (National DPP), a resource designed to bring such evidence-based lifestyle change programs for preventing type 2 diabetes to communities (cdc.gov/diabetes-prevention). To be eligible for this program, individuals must have a BMI in the overweight range and be at risk for diabetes based on laboratory testing, a previous diagnosis of GDM, or a positive risk test (cdc.gov/prediabetes/risktest/). The CDC has also developed the Diabetes Prevention Impact Tool Kit (nccd.cdc.gov/toolkit/diabetesimpact) to help organizations assess the economics of providing or

covering the National DPP (46). To expand preventive services using a cost-effective model, the Centers for Medicare & Medicaid Services expanded Medicare reimbursement coverage for the National DPP to organizations recognized by the CDC that become Medicare suppliers for this service (innovation.cms.gov/innovation-models/medicare-diabetes-prevention-program). The locations of Medicare DPPs are available online at innovation.cms.gov/innovation-models/medicare-diabetes-prevention-program/mdpp-map. To qualify for Medicare coverage, individuals must have BMI >25 kg/m² (or BMI >23 kg/m² if self-identified as Asian) and glycemic testing consistent with prediabetes in the last year. Medicaid DPP coverage is also expanding on a state-by-state basis.

While CDC-recognized behavioral counseling programs, including Medicare DPP services, have met minimum quality standards and are reimbursed by many payors, lower retention rates have been reported for younger adults and racial and ethnic minoritized populations (47). Therefore, other programs and modalities of behavioral counseling for diabetes prevention may also be appropriate and efficacious based on individual preferences and availability. Use of community health workers to support DPP-like interventions can be appropriate and cost-effective (48,49) (see section 1, “Improving Care and Promoting Health in Populations,” for more information). The use of community health workers may facilitate the adoption of behavior changes for diabetes prevention while bridging barriers related to social determinants of health. However, coverage by third-party payors remains limited. Counseling by a registered dietitian nutritionist has been shown to help individuals with prediabetes improve eating habits, increase physical activity, and achieve 7–10% weight loss (50–53). Individualized medical nutrition therapy (see section 5, “Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes,” for more detailed information) is also effective in improving glycemia in individuals diagnosed with prediabetes (51,54). Furthermore, trials involving medical nutrition therapy for adults with prediabetes found significant reductions in weight, waist circumference, and glycemia (54,55). Individuals with prediabetes can benefit

from referral to a registered dietitian nutritionist for individualized medical nutrition therapy upon diagnosis and at regular intervals throughout their treatment plan (52,56). Other health care professionals, such as pharmacists and diabetes care and education specialists, can also be considered for diabetes prevention efforts (56,57).

Technology-assisted programs may effectively deliver a DPP-like intervention (58–61). Such technology-assisted programs may deliver content through smartphones, web-based applications, and telehealth and may be an acceptable and efficacious option to bridge barriers, particularly for individuals with low income and/or in rural locations; however, not all technology-assisted programs are effective (58,62–64). The CDC Diabetes Prevention Recognition Program (DPRP) (<https://www.cdc.gov/diabetes-prevention/php/program-provider/program-requirements.html>) certifies technology-assisted modalities as effective vehicles for DPP-based interventions; such programs must use an approved curriculum, include interaction with a coach, and attain the DPP outcomes of participation, physical activity reporting, and weight loss. Health care professionals should consider referring adults with prediabetes to certified technology-assisted programs.

The CDC's Division of Diabetes Translation National DPP toolkit ([coveragetoolkit.org/recruitment-referral-for-the-national-dpp-lifestyle-change-program](https://www.cdc.gov/diabetes/prevention/toolkit/recruitment-referral-for-the-national-dpp-lifestyle-change-program)) provides diabetes care team members resources for how to increase referral and uptake of the DPP. Some ways to increase DPP utilization include direct mailings, phone scripts, and text-based outreach. Using multiple methods of outreach to increase awareness among those at risk, and among health care professionals, in addition offering the DPP virtually can improve participation and uptake (65,66). However, more work is needed to improve uptake among high-risk populations (67,68).

Sleep Characteristics Associated With Increased Risk of Type 2 Diabetes

Sleep occupies approximately one-third of the day for most people and modulates important metabolic, endocrine, and cardiovascular processes (69). Sleep can be characterized using three key

constructs: quantity, quality, and timing (i.e., chronotype). There is now established evidence for a U-shaped association between sleep duration and type 2 diabetes incidence, with the nadir typically occurring at 7 h per day, with short (typically defined as <6 h) and long (typically defined as >9 h) sleep duration having up to a 50% increase in the risk of type 2 diabetes, including progression from prediabetes (70). Sleep quality is defined as “an individual's self-satisfaction with all aspects of the sleep experience” (71,72). A meta-analysis reported that poor overall sleep quality was associated with a 40% increased risk of developing type 2 diabetes (73). Chronotype preference has been linked with many chronic diseases, including type 2 diabetes. For example, for those with a preference for evenings (i.e., going to bed late and getting up late), there was a 2.5-fold higher odds ratio for type 2 diabetes than for those with a preference for mornings (i.e., going to bed early and getting up early), independent of sleep duration and sleep sufficiency (74). More detailed information concerning sleep and the impact of sleep on prediabetes and diabetes risk and management can be found in section 5, “Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes.”

PHARMACOLOGIC INTERVENTIONS TO DELAY TYPE 2 DIABETES

Recommendations

3.7 Metformin for the prevention of type 2 diabetes should be considered in adults at high risk of type 2 diabetes, as typified by the Diabetes Prevention Program, especially those aged 25–59 years with BMI ≥ 35 kg/m², higher fasting plasma glucose (e.g., ≥ 110 mg/dL [≥ 6 mmol/L]), and higher A1C (e.g., $\geq 6.0\%$ [≥ 42 mmol/mol]), and in individuals with prior gestational diabetes mellitus. **A**

3.8 Consider using metformin to prevent hyperglycemia in high-risk individuals treated with a phosphatidylinositol 3-kinase α (PI3K α) inhibitor (e.g., alpelisib and inavolisib). **B**

3.9 Consider using metformin to prevent hyperglycemia in high-risk individuals treated with high-dose glucocorticoids. **B**

3.10 Consider periodic assessment of vitamin B12 levels in individuals receiving long-term metformin therapy, especially in those with anemia or peripheral neuropathy. **B**

Because weight loss and maintenance through lifestyle (e.g., nutrition and physical activity) behavior changes might not be sufficient on their own (13), some people at high risk for type 2 diabetes may benefit from additional pharmacotherapy support. Various pharmacologic agents used to treat diabetes have also been evaluated for diabetes prevention. Metformin, α -glucosidase inhibitors, incretin receptor agonists (e.g., liraglutide, semaglutide, and tirzepatide), thiazolidinediones, and insulin have all been shown to lower the incidence of diabetes in specific populations (75–83). However, whether initiating glucose-lowering treatment early—before hyperglycemia reaches the diagnostic threshold of diabetes—improves long-term health outcomes is unknown. Indeed, the benefits of glucose lowering to the normoglycemic range need to be balanced against the potential adverse effects of these therapies and the burden and cost of treatment. There are currently no long-term data to support the use of pharmacologic treatments other than metformin for the sole purpose of preventing type 2 diabetes. However, use of glucagon-like peptide 1–based therapies for weight management in individuals with overweight or obesity—an essential component of type 2 diabetes prevention—is highly beneficial and should be considered, as discussed in detail in section 8, “Obesity and Weight Management for the Prevention and Treatment of Diabetes.” Importantly, in the DPP, weight loss was key to reducing the risk of progression to diabetes, with every kilogram of weight loss conferring a 16% reduction in risk of progression over 3.2 years (18). In individuals with previous history of GDM, the risk of type 2 diabetes increased by 18% for every 1-unit increase in BMI above the preconception baseline (84). Several medications evaluated for obesity treatment (e.g., orlistat, phentermine and topiramate, liraglutide, semaglutide, and tirzepatide) have been shown to decrease the incidence of type 2 diabetes in those with prediabetes (80,85–87).

Other medications have also been assessed for their efficacy in preventing type 2 diabetes. The antihypertensive valsartan was found to modestly reduce the incidence of diabetes among individuals with prediabetes and cardiovascular disease or risk factors, but with no difference in cardiovascular outcomes (88). Thus, we do not recommend using valsartan for the prevention of diabetes. Testosterone therapy also reduced the risk of incident type 2 diabetes by 41% in men aged 50–74 years with central obesity and prediabetes (89). However, the benefit of testosterone was observed only when added to a structured lifestyle program. Indeed, the TRAVERSE Diabetes Study (a prespecified efficacy trial nested within the large multicenter TRAVERSE trial whose primary objective was to assess the cardiovascular outcomes of testosterone therapy in men aged 45–80 years with hypogonadism) found no benefit of testosterone as compared with placebo with respect to progression from prediabetes to diabetes or remission of diabetes among those who had prediabetes or diabetes at baseline, respectively (90). Testosterone therapy also did not alter glucose or A1C levels among participants with prediabetes or diabetes, nor were there any differences in outcomes by baseline testosterone level, preexisting cardiovascular disease, age, or race. Thus, we do not recommend testosterone therapy for the prevention of type 2 diabetes in men with hypogonadism; the decision to initiate testosterone replacement needs to be informed by individualized treatment considerations that weigh the expected benefits and harms of treatment.

There has been significant interest in understanding the potential of vitamin D therapy to prevent progression of high-risk prediabetes to type 2 diabetes in adults (91). Three randomized controlled trials tested whether vitamin D therapy in combination with lifestyle modification reduces the risk of developing diabetes in adults with high-risk prediabetes (i.e., IGT or meeting two or three criteria for the diagnosis of prediabetes [fasting glucose, A1C, 2-h glucose after a 75-g OGTT]): the Tromsø study in Norway, with 511 participants; the Vitamin D and Type 2 Diabetes (D2d) study in the U.S., with 2,423 participants; and the Diabetes Prevention with Active Vitamin D (DPVD) study

in Japan, with 1,256 participants (92–94). Although vitamin D therapy consistently appeared to modestly reduce the risk of developing diabetes in all three trials, none of the results were statistically significant, which was attributed to insufficient statistical power. Subsequently, several meta-analyses pooling the three randomized clinical trials along with other smaller studies suggested a modest (i.e., 10–15% risk reduction) potential benefit in specific populations, such as older adults, individuals with lower baseline vitamin D levels, and individuals without obesity (95,96). However, there are several concerns and uncertainties regarding recommending widespread vitamin D therapy for adults with high-risk prediabetes. First, these trials used varying dosages of vitamin D that were higher than the recommended daily allowance (i.e., 600 IU/day for those aged 18–70 years and 800 IU/day for those older than 70 years). Second, the benefit-to-risk ratio of vitamin D therapy for people with high-risk prediabetes remains uncertain. Although there were no safety concerns identified in these trials, the anticipated treatment of many millions of adults with prediabetes in the U.S. and globally could increase the risk of adverse events (such as hypercalcemia, hypercalciuria, nephrolithiasis, and kidney failure), especially treatment with unspecified doses of vitamin D without monitoring blood 25-hydroxy vitamin D levels.

Thus, the expected benefit, harm, treatment burden, and cost of using any medication to prevent type 2 diabetes must be considered for each individual as part of informed shared decision-making. This is particularly important because any pharmacologic intervention is likely to be needed to be used long-term because of waning effects after stopping the medication. At the present time, there are no pharmacologic agents approved by the U.S. Food and Drug Administration specifically for the prevention of type 2 diabetes.

Metformin

Metformin has the most robust efficacy and safety data as a pharmacologic therapy for diabetes prevention among people with prediabetes (82). Importantly, metformin was overall less effective than lifestyle modification in the DPP, although

group differences attenuated over time in the DPPOS (14) and metformin was as effective as lifestyle modification in participants with BMI ≥ 35 kg/m² and in younger participants aged 25–44 years (8). In individuals with a history of GDM, metformin and intensive lifestyle modification led to an equivalent 50% reduction in diabetes risk (97). Both interventions remained effective during a 10-year follow-up period (98), with lifestyle therapy being highly cost-effective (\$12,878 per quality-adjusted life-year [QALY]) and metformin marginally cost-saving compared with placebo (37). By the time of the 15-year follow-up (DPPOS), exploratory analyses demonstrated that participants with a higher baseline fasting glucose (≥ 110 mg/dL [≥ 6 mmol/L] vs. 95–109 mg/dL [5.3–5.9 mmol/L]), those with a higher A1C (6.0–6.4% [42–46 mmol/mol] vs. $<6.0\%$ [<42 mmol/mol]), and individuals with a history of GDM (vs. individuals without a history of GDM) experienced higher risk reductions with metformin, identifying subgroups of participants that may benefit the most from metformin (83). In the Indian Diabetes Prevention Program (IDPP-1), metformin and lifestyle intervention reduced diabetes risk similarly at 30 months; however, the lifestyle intervention in IDPP-1 was less intensive than that in the DPP (99). Based on findings from the DPP, metformin should be recommended as an option for high-risk individuals (e.g., younger individuals, those with history of GDM, or those with BMI ≥ 35 kg/m²).

Individuals receiving high-dose and/or long-duration glucocorticoid therapy are at risk for developing glucocorticoid-induced diabetes (see section 2, “Diagnosis and Classification of Diabetes”). Metformin can be considered to prevent the development of glucocorticoid-induced hyperglycemia in high-risk individuals, notably those receiving higher doses of glucocorticoids, with longer treatment duration, and in the presence of other risk factors for diabetes. The efficacy of metformin to prevent hyperglycemia—particularly insulin resistance and postprandial hyperglycemia—has been demonstrated in small randomized clinical trials both in healthy individuals (100) and in individuals receiving high-dose glucocorticoids for the treatment of cancer (101) and rheumatologic/autoimmune disorders (102,103). In one trial with 40 participants receiving high-

dose prednisolone for the treatment of inflammatory conditions, metformin also reduced rates of infection and all-cause hospital readmissions (103).

Proactive use of metformin may also be beneficial in preventing hyperglycemia in individuals receiving phosphatidylinositol 3-kinase α (PI3K α) inhibitors (e.g., alpelisib and inavolisib) for the treatment of cancer (see section 2, “Diagnosis and Classification of Diabetes,” for additional details on diabetes induced by systemic anticancer therapy). This is particularly important because hyperglycemia is the most frequent adverse event of PI3K α inhibitor therapy that leads to treatment discontinuation (104). In a phase 2 trial of metformin to prevent hyperglycemia in individuals receiving alpelisib for advanced breast cancer (metformin was started 1 week prior to starting alpelisib and continued for the duration of treatment), metformin prevented the development of grade 3–4 hyperglycemia in all but four participants, and none had to discontinue alpelisib therapy (105). Similar benefits were seen with implementation of a metformin hyperglycemia-prevention protocol in routine practice (106). Individuals at highest risk for developing hyperglycemia with PI3K α inhibitor therapy include those who are older (≥ 70 years old), with obesity (BMI ≥ 30 kg/m²), concurrently treated with glucocorticoids, and with hyperglycemia at baseline (e.g., A1C $\geq 5.7\%$ [39 mmol/mol] or fasting plasma glucose ≥ 100 mg/dL [5.6 mmol/L]) (107, 108). However, it is important to weigh the benefits of preventing hyperglycemia with the potential adverse effects of metformin therapy, particularly diarrhea, which is also a frequent adverse effect of PI3K α therapy (though there is no evidence that concurrent use of these two medications further exacerbates diarrhea risk) (108).

Decreased vitamin B12 levels are a known consequence of long-term treatment with metformin (109). Periodic assessment of vitamin B12 level in those taking metformin chronically should be considered to check for possible deficiency, especially in those receiving a higher dose (e.g., $\geq 1,500$ mg/day) (110), with longer treatment duration (4–5 years) (111), with other risk factors for vitamin B12 deficiency, and if a deficiency is suspected, such as in people with anemia,

peripheral neuropathy, or chronic kidney disease (109,112,113) (see section 9, “Pharmacologic Approaches to Glycemic Treatment,” for more details). A person who has been taking metformin for more than 4 years or is at risk for vitamin B12 deficiency for other reasons (e.g., vegan dietary pattern, previous gastric/small bowel surgery) should be monitored for vitamin B12 deficiency annually (114,115).

PREVENTION OF VASCULAR DISEASE AND MORTALITY

Recommendations

3.11 Prediabetes is associated with heightened cardiovascular risk; therefore, screening for and treatment of modifiable risk factors for cardiovascular disease are suggested. **B**

3.12 Statin therapy may increase the risk of type 2 diabetes in people at high risk of developing type 2 diabetes. In such individuals, glucose status should be monitored regularly and diabetes prevention approaches reinforced. It is not recommended that statins be avoided or discontinued for this adverse effect. **B**

3.13 In people with a history of stroke and evidence of insulin resistance and prediabetes, pioglitazone may be considered to lower the risk of stroke or myocardial infarction. However, this benefit needs to be balanced with the increased risk of weight gain, edema, and fractures. **A** Lower doses may mitigate the risk of adverse effects but may be less effective. **C**

People with prediabetes often have other cardiovascular risk factors, including hypertension and dyslipidemia (116), and are at increased risk for cardiovascular disease (117,118). Evaluation for tobacco use and referral for tobacco cessation should be part of routine care for those at risk for diabetes. Of note, the years immediately following smoking cessation may represent a time of increased risk for diabetes (119,120), and individuals should be monitored for diabetes development and receive evidence-based lifestyle behavior change for diabetes prevention as described in this section. See section 5, “Facilitating Positive Health Behaviors and Well-being to Improve Health Outcomes,” for more detailed information. The lifestyle interventions for weight loss in study populations at risk for

type 2 diabetes have shown a reduction in cardiovascular risk factors and the need for medications used to treat these cardiovascular risk factors (121,122). The lifestyle intervention in the Da Qing study was associated with lowering cardiovascular disease and mortality at 23 and 30 years of observational follow-up (10,12). Treatment goals and therapies for hypertension and dyslipidemia in the primary and secondary prevention of cardiovascular disease for people with prediabetes should be based on their level of cardiovascular risk. Increased vigilance is warranted to identify and treat these and other cardiovascular disease risk factors (123).

Statin use increases risk of diabetes (124,125). In the DPP, statin use was associated with greater diabetes risk irrespective of the treatment group (pooled hazard ratio [HR] [95% CI] for incident diabetes 1.36 [1.17–1.58]) (124). In trials of primary and secondary prevention of cardiovascular disease, cardiovascular and mortality benefits of statin therapy exceed the risk of diabetes (126,127), suggesting a highly favorable benefit-to-harm balance with statin therapy. Hence, discontinuation of statins due to concerns of diabetes risk is not recommended in this population.

Results of cardiovascular outcome trials in people without diabetes also provide important evidence for cardiovascular risk reduction in people with increased cardiometabolic risk, though not specifically prediabetes (see section 10, “Cardiovascular Disease and Risk Management,” for more details). The IRIS (Insulin Resistance Intervention after Stroke) trial of people with a recent (< 6 months) stroke or transient ischemic attack, without diabetes but with insulin resistance (as defined by a HOMA of insulin resistance index of ≥ 3.0), evaluated pioglitazone (goal dose of 45 mg daily) compared with placebo. At 4.8 years, the risk of stroke or myocardial infarction, as well as the risk of diabetes, was lower in the pioglitazone group than in the placebo group; weight gain, edema, and fractures were higher in the pioglitazone treatment group (128–130). Lower doses may mitigate the adverse effects but may also be less effective (131).

PERSON-CENTERED CARE GOALS

Recommendations

3.14 In adults with overweight or obesity at high risk of type 2 diabetes, care goals should include weight

loss and maintenance, minimizing the progression of hyperglycemia, and attention to cardiovascular risk. **B**

3.15 Pharmacotherapy (e.g., for weight management, minimizing the progression of hyperglycemia, and cardiovascular risk reduction) should be considered to support person-centered care goals. **A**

3.16 More intensive preventive approaches should be considered in individuals who are at particularly high risk of progression to diabetes, including individuals with BMI ≥ 35 kg/m², those with higher glucose levels (e.g., fasting plasma glucose 110–125 mg/dL [6.1–6.9 mmol/L], 2-h postchallenge glucose 173–199 mg/dL [9.6–11.0 mmol/L], and A1C $\geq 6.0\%$ [≥ 42 mmol/mol]), and individuals with a history of gestational diabetes mellitus. **A**

Individualized risk-to-benefit ratio should be considered in screening, intervention, and monitoring to lower the risk of type 2 diabetes and associated comorbidities. Multiple factors, including age, BMI, and other comorbidities, may influence the risk of progression to diabetes and lifetime risk of complications (132,133). Prediabetes is associated with increased cardiovascular disease and mortality (118), which emphasizes the importance of attending to cardiovascular risk in this population. However, the new diagnosis of prediabetes in older adults (aged >70 years) is less relevant for progression to diabetes; indeed, regression to normoglycemia or death was more frequent than progression to diabetes in the Atherosclerosis Risk in Communities (ARIC) study (132). Moreover, diabetes-related morbidity and mortality is lower and the benefits of treatment are not established in those who develop type 2 diabetes after age 70 years (134).

In the DPP, which enrolled high-risk individuals with IGT, elevated fasting glucose, and elevated BMI, the crude incidence of diabetes within the placebo group was 11 cases per 100 person-years, with a cumulative 3-year incidence of diabetes of 29% (8). Characteristics of individuals in the DPP and DPPOS who were at particularly high risk of progression to diabetes (crude incidence of diabetes 14–22 cases per 100 person-years)

included BMI ≥ 35 kg/m², higher glucose levels (e.g., fasting plasma glucose 110–125 mg/dL [6.0–6.9 mmol/L], 2-h postchallenge glucose 173–199 mg/dL [9.6–11.0 mmol/L], A1C $\geq 6.0\%$ [≥ 42 mmol/mol]), or a history of GDM (8,97,98). In contrast, in the community-based ARIC study, observational follow-up of adults with mean age 75 years with laboratory evidence of prediabetes (based on A1C 5.7–6.4% [39–47 mmol/mol] and/or fasting glucose 100–125 mg/dL [5.6–6.9 mmol/L]), but not meeting specific BMI criteria, found lower progression to diabetes over 6 years: 9% of those with A1C-defined prediabetes and 8% of those with IFG (133).

Thus, it is important to individualize the risk-to-benefit ratio of intervention and consider person-centered goals. Risk models have generally found a higher benefit of the intervention in those at highest risk (18). Diabetes prevention trials and observational studies highlight key principles that may guide person-centered goals. In the DPP, which enrolled a high-risk population meeting criteria for overweight or obesity, weight loss was an important mediator of diabetes prevention or delay, with greater metabolic benefit seen with greater weight loss (18,135). In the DPP and DPPOS, progression to diabetes, duration of diabetes, and mean level of glycemia were important determinants of the development of microvascular complications (14). Achieving normal glucose regulation, even once, during the DPP was associated with a lower risk of diabetes and lower risk of microvascular complications irrespective of the treatment arm (136). Observational follow-up of the Da Qing study also showed that regression from IGT to normal glucose tolerance or remaining with IGT rather than progressing to type 2 diabetes at the end of the 6-year intervention trial resulted in significantly lower risk of cardiovascular disease and microvascular disease over 30 years (137).

Pharmacotherapy for weight management and cardiovascular risk reduction (see section 8, “Obesity and Weight Management for the Prevention and Treatment of Diabetes,” and section 10, “Cardiovascular Disease and Risk Management,” for more details) can be considered to support individualized person-centered goals, with more intensive preventive approaches

considered in individuals at high risk of progression.

PREVENTION OR DELAY OF SYMPTOMATIC TYPE 1 DIABETES

Lifestyle and Type 1 Diabetes Progression

Observational studies have identified several factors that increase β -cell demand and risk of progression to clinical type 1 diabetes among individuals with islet autoantibodies, including lower levels of physical activity (138), higher glycemic index eating patterns (139), and total sugar intake (140). Similar associations have not been observed for the development of autoantibodies. In The Environmental Determinants of Diabetes in the Young (TEDDY) longitudinal study, daily minutes spent in moderate to vigorous physical activity were associated with a reduced risk of progression to type 1 diabetes in children 5–15 years of age with multiple islet autoantibodies (HR 0.92 [95% CI 0.86–0.99] per 10-min increase; $P = 0.02$) (138). In the Diabetes Autoimmunity Study in the Young (DAISY), also in children with islet autoantibodies, consumption of higher glycemic index foods (HR 2.20 [95% CI 1.17–4.15]) and total sugar intake (HR 1.75 [95% CI 1.07–2.85]) (139,140) were both associated with progression to type 1 diabetes. In nonobese diabetic mice, an animal model for the development of type 1 diabetes, sustained high-glucose drinking significantly aggravated islet inflammation and accelerated the onset of type 1 diabetes (141). However, efficacy of lifestyle interventions that modify these factors in individuals with stage 1 or stage 2 type 1 diabetes has not yet been reported.

Pharmacologic Interventions to Delay Symptomatic Type 1 Diabetes

Recommendation

3.17 Teplizumab-mzwv infusion to delay the onset of symptomatic type 1 diabetes (stage 3) should be discussed with selected individuals aged ≥ 8 years with stage 2 type 1 diabetes. Treatment should be in a setting with appropriately trained personnel. **B**

Teplizumab, a CD3-directed humanized monoclonal antibody engineered to have decreased Fc receptor binding, has been

approved to delay the onset of stage 3 type 1 diabetes in people 8 years and older with stage 2 type 1 diabetes based in part on the results of a single trial in relatives of people with type 1 diabetes (142). In this study, 44 individuals were randomized to a 14-day course of teplizumab and 32 to placebo. The median time to stage 3 type 1 diabetes diagnosis was 48.4 months in the teplizumab group and 24.4 months in the placebo group. Type 1 diabetes was diagnosed in 19 (43%) participants who received teplizumab and 23 (72%) participants who received placebo (HR 0.41 [95% CI 0.22–0.78]). In prespecified analyses, the presence of HLA-DR4, absence of HLA-DR3, and absence of antizinc transporter 8 antibody predicted response to teplizumab (HR 0.20 [95% CI 0.09–0.45], 0.18 [0.07–0.45], and 0.07 [0.02–0.26], respectively). The most common adverse reactions were transient lymphopenia (73%) followed by rash (36%).

Numerous clinical studies are being conducted to test methods for preventing or delaying the onset of stage 3 type 1 diabetes in those with evidence of autoimmunity without symptoms or for delaying loss of insulin secretory capacity after onset of stage 3, some with promising results (see ClinicalTrials.gov and TrialNet.org).

References

- Ziegler AG, Rewers M, Simell O, et al. Seroconversion to multiple islet autoantibodies and risk of progression to diabetes in children. *JAMA* 2013;309:2473–2479
- Steck AK, Dong F, Taki I, et al. Continuous glucose monitoring predicts progression to diabetes in autoantibody positive children. *J Clin Endocrinol Metab* 2019;104:3337–3344
- Ylescupidez A, Speake C, Pietropaolo SL, et al. OGTT metrics surpass continuous glucose monitoring data for T1D prediction in multiple-autoantibody-positive individuals. *J Clin Endocrinol Metab* 2023;109:57–67
- Phillip M, Achenbach P, Addala A, et al. Consensus guidance for monitoring individuals with islet autoantibody-positive pre-stage 3 type 1 diabetes. *Diabetes Care* 2024;47:1276–1298
- Sims EK, Besser REJ, Dayan C, et al.; NIDDK Type 1 Diabetes TrialNet Study Group. Screening for type 1 diabetes in the general population: a status report and perspective. *Diabetes* 2022;71:610–623
- Beran D, Bandini A, Bosi E, et al. Type 1 diabetes screening: need for ethical, equity, and health systems perspective. *Lancet Diabetes Endocrinol* 2025;13:175–176
- Wentworth JM, Sing ABE, Naselli G, et al. Islet autoantibody screening throughout Australia using in-home blood spot sampling: 2-year outcomes of Type1Screen. *Diabetes Care* 2025;48:556–563
- Knowler WC, Barrett-Connor E, Fowler SE, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393–403
- Lindström J, Ilanne-Parikka P, Peltonen M, et al.; Finnish Diabetes Prevention Study Group. Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study. *Lancet* 2006;368:1673–1679
- Li G, Zhang P, Wang J, et al. Cardiovascular mortality, all-cause mortality, and diabetes incidence after lifestyle intervention for people with impaired glucose tolerance in the Da Qing Diabetes Prevention Study: a 23-year follow-up study. *Lancet Diabetes Endocrinol* 2014;2:474–480
- Nathan DM, Bennett PH, Crandall JP, et al.; DPP Research Group. Does diabetes prevention translate into reduced long-term vascular complications of diabetes? *Diabetologia* 2019;62:1319–1328
- Gong Q, Zhang P, Wang J, et al.; Da Qing Diabetes Prevention Study Group. Morbidity and mortality after lifestyle intervention for people with impaired glucose tolerance: 30-year results of the Da Qing Diabetes Prevention Outcome Study. *Lancet Diabetes Endocrinol* 2019;7:452–461
- Knowler WC, Fowler SE, Hamman RF, et al. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet* 2009;374:1677–1686
- Diabetes Prevention Program Research Group. Long-term effects of lifestyle intervention or metformin on diabetes development and microvascular complications over 15-year follow-up: the Diabetes Prevention Program Outcomes Study. *Lancet Diabetes Endocrinol* 2015;3:866–875
- Knowler WC, Doherty L, Edelstein SL, et al.; DPP/DPPOS Research Group. Long-term effects and effect heterogeneity of lifestyle and metformin interventions on type 2 diabetes incidence over 21 years in the US Diabetes Prevention Program randomised clinical trial. *Lancet Diabetes Endocrinol* 2025;13:469–481
- Crandall JP, Dabelea D, Knowler WC, et al.; DPP Research Group. The Diabetes Prevention Program and its Outcomes Study: NIDDK's journey into the prevention of type 2 diabetes and its public health impact. *Diabetes Care* 2025;48:1101–1111
- Diabetes Prevention Program (DPP) Research Group. The Diabetes Prevention Program (DPP): description of lifestyle intervention. *Diabetes Care* 2002;25:2165–2171
- Hamman RF, Wing RR, Edelstein SL, et al. Effect of weight loss with lifestyle intervention on risk of diabetes. *Diabetes Care* 2006;29:2102–2107
- Goldberg RB, Orchard TJ, Crandall JP, et al.; Diabetes Prevention Program Research Group. Effects of long-term metformin and lifestyle interventions on cardiovascular events in the Diabetes Prevention Program and its Outcome Study. *Circulation* 2022;145:1632–1641
- Dagogo-Jack S, Umekwe N, Brewer AA, et al. Outcome of lifestyle intervention in relation to duration of pre-diabetes: the Pathobiology and Reversibility of Prediabetes in a Biracial Cohort (PROP-ABC) study. *BMJ Open Diabetes Res Care* 2022;10:e002748
- Dietary Guidelines Advisory Committee. *Scientific Report of the 2025 Dietary Guidelines Advisory Committee: Advisory Report to the Secretary of Health and Human Services and Secretary of Agriculture*. U.S. Department of Health and Human Services, 2024
- Jannasch F, Kröger J, Schulze MB. Dietary patterns and type 2 diabetes: a systematic literature review and meta-analysis of prospective studies. *J Nutr* 2017;147:1174–1182
- Salmela J, Kontinen H, Lappalainen R, et al. Eating behavior dimensions and 9-year weight loss maintenance: a sub-study of the Finnish Diabetes Prevention Study. *Int J Obes (Lond)* 2023;47:564–573
- Guest NS, Raj S, Landry MJ, et al. Vegetarian and vegan dietary patterns to treat adult type 2 diabetes: a systematic review and meta-analysis of randomized controlled trials. *Adv Nutr* 2024;15:100294
- Bensaoud A, Seery S, Gibson I, et al. Dietary approaches to stop hypertension (DASH) for the primary and secondary prevention of cardiovascular diseases. *Cochrane Database Syst Rev* 2025;5:CD013729
- Qian F, Liu G, Hu FB, Bhupathiraju SN, Sun Q. Association between plant-based dietary patterns and risk of type 2 diabetes: a systematic review and meta-analysis. *JAMA Intern Med* 2019;179:1335–1344
- Yau JW, Thor SM, Ramadas A. Nutritional strategies in prediabetes: a scoping review of recent evidence. *Nutrients* 2020;12:2990
- Ley SH, Hamdy O, Mohan V, Hu FB. Prevention and management of type 2 diabetes: dietary components and nutritional strategies. *Lancet* 2014;383:1999–2007
- Dudzík JM, Senkus KE, Evert AB, et al. The effectiveness of medical nutrition therapy provided by a dietitian in adults with prediabetes: a systematic review and meta-analysis. *Am J Clin Nutr* 2023;118:892–910
- Razaz JM, Rahmani J, Varkaneh HK, Thompson J, Clark C, Abdulazeem HM. The health effects of medical nutrition therapy by dietitians in patients with diabetes: a systematic review and meta-analysis: nutrition therapy and diabetes. *Prim Care Diabetes* 2019;13:399–408
- García-Hermoso A, López-Gil JF, Izquierdo M, Ramírez-Vélez R, Ezzatvar Y. Exercise and insulin resistance markers in children and adolescents with excess weight: a systematic review and network meta-analysis. *JAMA Pediatr* 2023;177:1276–1284
- Davis CL, Pollock NK, Waller JL, et al. Exercise dose and diabetes risk in overweight and obese children: a randomized controlled trial. *JAMA* 2012;308:1103–1112
- Dai X, Zhai L, Chen Q, et al. Two-year-supervised resistance training prevented diabetes incidence in people with prediabetes: a randomised control trial. *Diabetes Metab Res Rev* 2019;35:e3143
- Thorp AA, Kingwell BA, Sethi P, Hammond L, Owen N, Dunstan DW. Alternating bouts of sitting and standing attenuate postprandial glucose responses. *Med Sci Sports Exerc* 2014;46:2053–2061
- Rute-Larrieta C, Mota-Cátedra G, Carmona-Torres JM, et al. Physical activity during pregnancy and risk of gestational diabetes mellitus: a meta-review. *Life (Basel)* 2024;14:755
- Santa Cruz TE, Sarasqueta C, Muruzábal JC, Ansuategui E, Sanz O. A systematic review and meta-analysis of exercise-based intervention to prevent gestational diabetes in women with

- overweight or obesity. *BMC Pregnancy Childbirth* 2025;25:5
37. Herman WH, Hoerger TJ, Brandle M, et al.; Diabetes Prevention Program Research Group. The cost-effectiveness of lifestyle modification or metformin in preventing type 2 diabetes in adults with impaired glucose tolerance. *Ann Intern Med* 2005;142:323–332
 38. Chen F, Su W, Becker SH, et al. Clinical and economic impact of a digital, remotely-delivered intensive behavioral counseling program on medicare beneficiaries at risk for diabetes and cardiovascular disease. *PLoS One* 2016;11:e0163627
 39. Diabetes Prevention Program Research Group. The 10-year cost-effectiveness of lifestyle intervention or metformin for diabetes prevention: an intent-to-treat analysis of the DPP/DPPOS. *Diabetes Care* 2012;35:723–730
 40. Alva ML, Hoerger TJ, Jeyaraman R, Amico P, Rojas-Smith L. Impact of the YMCA of the USA diabetes prevention program on Medicare spending and utilization. *Health Aff (Millwood)* 2017;36:417–424
 41. Zhou X, Siegel KR, Ng BP, et al. Cost-effectiveness of diabetes prevention interventions targeting high-risk individuals and whole populations: a systematic review. *Diabetes Care* 2020;43:1593–1616
 42. Ackermann RT, O'Brien MJ. Evidence and challenges for translation and population impact of the Diabetes Prevention Program. *Curr Diab Rep* 2020;20:9
 43. Balk EM, Earley A, Raman G, Avendano EA, Pittas AG, Remington PL. Combined diet and physical activity promotion programs to prevent type 2 diabetes among persons at increased risk: a systematic review for the Community Preventive Services Task Force. *Ann Intern Med* 2015;163:437–451
 44. Ackermann RT, Kang R, Cooper AJ, et al. Effect on health care expenditures during nationwide implementation of the Diabetes Prevention Program as a health insurance benefit. *Diabetes Care* 2019;42:1776–1783
 45. Fitzpatrick-Lewis D, Ali MU, Horvath S, Nagpal S, Ghanem S, Sherifali D. Effectiveness of workplace interventions to reduce the risk for type 2 diabetes: a systematic review and meta-analysis. *Can J Diabetes* 2022;46:84–98
 46. Lanza A, Soler R, Smith B, Hoerger T, Neuwahl S, Zhang P. The Diabetes Prevention Impact Tool Kit: an online tool kit to assess the cost-effectiveness of preventing type 2 diabetes. *J Public Health Manag Pract* 2019;25:E1–E5
 47. Cannon MJ, Masalovich S, Ng BP, et al. Retention among participants in the National Diabetes Prevention Program lifestyle change program, 2012–2017. *Diabetes Care* 2020;43:2042–2049
 48. The Community Guide. Diabetes Prevention: interventions Engaging Community Health Workers, 2016. Accessed 19 August 2025. Available from <https://www.thecommunityguide.org/findings/diabetes-prevention-interventions-engaging-community-health-workers>
 49. Zare H, Delgado P, Spencer M, et al. Using community health workers to address barriers to participation and retention in diabetes prevention program: a concept paper. *J Prim Care Community Health* 2022;13:21501319221134563
 50. Evert AB, Dennison M, Gardner CD, et al. Nutrition therapy for adults with diabetes or prediabetes: a consensus report. *Diabetes Care* 2019;42:731–754
 51. Raynor HA, Davidson PG, Burns H, et al. Medical nutrition therapy and weight loss questions for the evidence analysis library prevention of type 2 diabetes project: systematic reviews. *J Acad Nutr Diet* 2017;117:1578–1611
 52. Khk L. Nutrition therapy for adults with diabetes or prediabetes. *ADCES Pract* 2022;10:34–38
 53. Briggs Early K, Stanley K. Position of the Academy of Nutrition and Dietetics: the role of medical nutrition therapy and registered dietitian nutritionists in the prevention and treatment of prediabetes and type 2 diabetes. *J Acad Nutr Diet* 2018;118:343–353
 54. Parker AR, Byham-Gray L, Denmark R, Winkle PJ. The effect of medical nutrition therapy by a registered dietitian nutritionist in patients with prediabetes participating in a randomized controlled clinical research trial. *J Acad Nutr Diet* 2014;114:1739–1748
 55. Esposito K, Chiodini P, Maiorino MI, Bellastella G, Panagiotakos D, Giugliano D. Which diet for prevention of type 2 diabetes? A meta-analysis of prospective studies. *Endocrine* 2014;47:107–116
 56. Powers MA, Bardsley JK, Cypress M, et al. Diabetes self-management education and support in adults with type 2 diabetes: a consensus report of the American Diabetes Association, the Association of Diabetes Care & Education Specialists, the Academy of Nutrition and Dietetics, the American Academy of Family Physicians, the American Academy of PAs, the American Association of Nurse Practitioners, and the American Pharmacists Association. *Diabetes Care* 2020;43:1636–1649
 57. Hudspeth BD. Power of prevention: the pharmacist's role in prediabetes management. *Diabetes Spectr* 2018;31:320–323
 58. Grock S, Ku J-h, Kim J, Moin T. A review of technology-assisted interventions for diabetes prevention. *Curr Diab Rep* 2017;17:107
 59. Bian RR, Piatt GA, Sen A, et al. The effect of technology-mediated diabetes prevention interventions on weight: a meta-analysis. *J Med Internet Res* 2017;19:e76
 60. Moin T, Damschroder LJ, AuYoung M, et al. Results from a trial of an online diabetes prevention program intervention. *Am J Prev Med* 2018;55:583–591
 61. Michaelides A, Major J, Pienkosz E, Wood M, Kim Y, Toro-Ramos T. Usefulness of a novel mobile diabetes prevention program delivery platform with human coaching: 65-week observational follow-up. *JMIR Mhealth Uhealth* 2018;6:e93
 62. Kim SE, Castro Sweet CM, Cho E, Tsai J, Cousineau MR. Evaluation of a digital diabetes prevention program adapted for low-income patients, 2016–2018. *Prev Chronic Dis* 2019;16:E155
 63. Vadheim LM, Patch K, Brokaw SM, et al. Telehealth delivery of the diabetes prevention program to rural communities. *Transl Behav Med* 2017;7:286–291
 64. Fischer HH, Durfee MJ, Raghunath SG, Ritchie ND. Short message service text message support for weight loss in patients with prediabetes: pragmatic trial. *JMIR Diabetes* 2019;4:e12985
 65. Formagini T, Teruel Camargo J, Perales-Puchalt J, Drees BM, Fracachan Cabrera M, Ramírez M. A culturally and linguistically adapted text-message diabetes prevention program for Latinos: feasibility, acceptability, and preliminary effectiveness. *Transl Behav Med* 2024;14:138–147
 66. Auster-Gussman LA, Lockwood KG, Graham SA, Stein N, Branch OH. Reach of a fully digital diabetes prevention program in health professional shortage areas. *Popul Health Manag* 2022;25:441–448
 67. Malone A, Clair K, Chanfreau C, et al. Predictors of enrollment in a virtual diabetes prevention program among women veterans: a retrospective analysis. *BMC Womens Health* 2024;24:465
 68. Wilson KE, Michaud TL, Almeida FA, et al. Using a population health management approach to enroll participants in a diabetes prevention trial: reach outcomes from the PREDICTS randomized clinical trial. *Transl Behav Med* 2021;11:1066–1077
 69. Henson J, Covenant A, Hall AP, et al. Waking up to the importance of sleep in type 2 diabetes management: a narrative review. *Diabetes Care* 2024;47:331–343
 70. Davies MJ, Aroda VR, Collins BS, et al. Management of hyperglycemia in type 2 diabetes, 2022. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* 2022;45:2753–2786
 71. Mostafa SA, Mena SC, Antza C, Balanos G, Nirantharakumar K, Tahrani AA. Sleep behaviours and associated habits and the progression of prediabetes to type 2 diabetes mellitus in adults: a systematic review and meta-analysis. *Diab Vasc Dis Res* 2022;19:14791641221088824
 72. Nelson KL, Davis JE, Corbett CF. Sleep quality: an evolutionary concept analysis. *Nurs Forum* 2022;57:144–151
 73. Anothaisintawee T, Reutrakul S, Van Cauter E, Thakkinian A. Sleep disturbances compared to traditional risk factors for diabetes development: systematic review and meta-analysis. *Sleep Med Rev* 2016;30:11–24
 74. Merikanto I, Lahti T, Puolijoki H, et al. Associations of chronotype and sleep with cardiovascular diseases and type 2 diabetes. *Chronobiol Int* 2013;30:470–477
 75. Gerstein HC, Bosch J, Dagenais GR, et al.; ORIGIN Trial Investigators. Basal insulin and cardiovascular and other outcomes in dysglycemia. *N Engl J Med* 2012;367:319–328
 76. DeFronzo RA, Tripathy D, Schwenke DC, et al.; ACT NOW Study. Pioglitazone for diabetes prevention in impaired glucose tolerance. *N Engl J Med* 2011;364:1104–1115
 77. Gerstein HC, Yusuf S, Bosch J, et al.; DREAM (Diabetes REduction Assessment with ramipril and rosiglitazone Medication) Trial Investigators. Effect of rosiglitazone on the frequency of diabetes in patients with impaired glucose tolerance or impaired fasting glucose: a randomised controlled trial. *Lancet* 2006;368:1096–1105
 78. Le Roux CW, Astrup A, Fujioka K, et al.; SCALE Obesity Prediabetes NN8022-1839 Study Group. 3 years of liraglutide versus placebo for type 2 diabetes risk reduction and weight management in individuals with prediabetes: a randomised, double-blind trial. *Lancet* 2017;389:1399–1409

79. Chiasson J-L, Josse RG, Gomis R, Hanefeld M, Karasik A, Laakso M, STOP-NIDDM Trial Research Group. Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomised trial. *Lancet* 2002;359:2072–2077
80. Wilding JPH, Batterham RL, Calanna S, et al.; STEP 1 Study Group. Once-weekly semaglutide in adults with overweight or obesity. *N Engl J Med* 2021;384:989–1002
81. Jastreboff AM, Aronne LJ, Ahmad NN, et al.; SURMOUNT-1 Investigators. Tirzepatide once weekly for the treatment of obesity. *N Engl J Med* 2022;387:205–216
82. Diabetes Prevention Program Research Group. Long-term safety, tolerability, and weight loss associated with metformin in the Diabetes Prevention Program Outcomes Study. *Diabetes Care* 2012;35:731–737
83. Diabetes Prevention Program Research Group. Long-term effects of metformin on diabetes prevention: identification of subgroups that benefited most in the Diabetes Prevention Program and Diabetes Prevention Program Outcomes Study. *Diabetes Care* 2019;42:601–608
84. Dennison RA, Chen ES, Green ME, et al. The absolute and relative risk of type 2 diabetes after gestational diabetes: a systematic review and meta-analysis of 129 studies. *Diabetes Res Clin Pract* 2021;171:108625
85. Torgerson JS, Hauptman J, Boldrin MN, Sjöström L. XENical in the prevention of diabetes in obese subjects (XENDOS) study: a randomized study of orlistat as an adjunct to lifestyle changes for the prevention of type 2 diabetes in obese patients. *Diabetes Care* 2004;27:155–161
86. Garvey WT, Ryan DH, Henry R, et al. Prevention of type 2 diabetes in subjects with prediabetes and metabolic syndrome treated with phentermine and topiramate extended release. *Diabetes Care* 2014;37:912–921
87. Jastreboff AM, Le Roux CW, Stefanski A, et al.; SURMOUNT-1 Investigators. Tirzepatide for obesity treatment and diabetes prevention. *N Engl J Med* 2025;392:958–971
88. McMurray JJ, Holman RR, Haffner SM, et al.; NAVIGATOR Study Group. Effect of valsartan on the incidence of diabetes and cardiovascular events. *N Engl J Med* 2010;362:1477–1490
89. Wittert G, Bracken K, Robledo KP, et al. Testosterone treatment to prevent or revert type 2 diabetes in men enrolled in a lifestyle programme (T4DM): a randomised, double-blind, placebo-controlled, 2-year, phase 3b trial. *Lancet Diabetes Endocrinol* 2021;9:32–45
90. Bhasin S, Lincoff AM, Nissen SE, et al. Effect of testosterone on progression from prediabetes to diabetes in men with hypogonadism: a substudy of the TRAVERSE randomized clinical trial. *JAMA Intern Med* 2024;184:353–362
91. Demay MB, Pittas AG, Bikle DD, et al. Vitamin D for the prevention of disease: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2024;109:1907–1947
92. Jorde R, Sollid ST, Svartberg J, et al. Vitamin D 20,000 IU per week for five years does not prevent progression from prediabetes to diabetes. *J Clin Endocrinol Metab* 2016;101:1647–1655
93. Pittas AG, Dawson-Hughes B, Sheehan P, et al.; D2d Research Group. Vitamin D supplementation and prevention of type 2 diabetes. *N Engl J Med* 2019;381:520–530
94. Kawahara T, Suzuki G, Mizuno S, et al. Effect of active vitamin D treatment on development of type 2 diabetes: DPVD randomised controlled trial in Japanese population. *BMJ* 2022;377:e066222
95. Pittas AG, Kawahara T, Jorde R, et al. Vitamin D and risk for type 2 diabetes in people with prediabetes: a systematic review and meta-analysis of individual participant data from 3 randomized clinical trials. *Ann Intern Med* 2023;176:355–363
96. Shah VP, Nayfeh T, Alsawaf Y, et al. A systematic review supporting the Endocrine Society clinical practice guidelines on vitamin D. *J Clin Endocrinol Metab* 2024;109:1961–1974
97. Ratner RE, Christophi CA, Metzger BE, et al.; Diabetes Prevention Program Research Group. Prevention of diabetes in women with a history of gestational diabetes: effects of metformin and lifestyle interventions. *J Clin Endocrinol Metab* 2008;93:4774–4779
98. Aroda VR, Christophi CA, Edelstein SL, et al.; Diabetes Prevention Program Research Group. The effect of lifestyle intervention and metformin on preventing or delaying diabetes among women with and without gestational diabetes: the Diabetes Prevention Program Outcomes Study 10-year follow-up. *J Clin Endocrinol Metab* 2015;100:1646–1653
99. Ramachandran A, Snehalatha C, Mary S, Mukesh B, Bhaskar AD, Vijay V, Indian Diabetes Prevention Programme (IDPP). The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia* 2006;49:289–297
100. Thierry S, Peterson CJ, Pfammatter S, et al. Short-term metformin protects against glucocorticoid-induced toxicity in healthy individuals: a randomized, double-blind, placebo-controlled trial. *Diabetes Care* 2025;48:719–727
101. Ochola LA, Nyamu DG, Guantai EM, Weru IW. Metformin's effectiveness in preventing prednisone-induced hyperglycemia in hematological cancers. *J Oncol Pharm Pract* 2020;26:823–834
102. Seelig E, Meyer S, Timper K, et al. Metformin prevents metabolic side effects during systemic glucocorticoid treatment. *Eur J Endocrinol* 2017;176:349–358
103. Pernicova I, Kelly S, Ajodha S, et al. Metformin to reduce metabolic complications and inflammation in patients on systemic glucocorticoid therapy: a randomised, double-blind, placebo-controlled, proof-of-concept, phase 2 trial. *Lancet Diabetes Endocrinol* 2020;8:278–291
104. Cheung Y-MM, Cromwell GE, Tolaney SM, Min L, McDonnell ME. Factors leading to alpelisib discontinuation in patients with hormone receptor positive, human epidermal growth factor receptor-2 negative breast cancer. *Breast Cancer Res Treat* 2022;192:303–311
105. Llombart-Cussac A, Pérez-García JM, Ruiz Borrego M, et al. Preventing alpelisib-related hyperglycaemia in HR+/HER2-/PIK3CA-mutated advanced breast cancer using metformin (METALLICA): a multicentre, open-label, single-arm, phase 2 trial. *EClinicalMedicine* 2024;71:102520
106. Burnette SE, Poehlein E, Lee H-J, Force J, Westbrook K, Moore HN. Evaluation of alpelisib-induced hyperglycemia prophylaxis and associated risk factors in PIK3CA-mutated hormone-receptor positive, human epidermal growth factor-2 negative advanced breast cancer. *Breast Cancer Res Treat* 2023;197:369–376
107. Moore HN, Goncalves MD, Johnston AM, et al. Effective strategies for the prevention and mitigation of phosphatidylinositol-3-kinase inhibitor-associated hyperglycemia: optimizing patient care. *Clin Breast Cancer* 2025;25:1–11
108. Rugo HS, André F, Yamashita T, et al. Time course and management of key adverse events during the randomized phase III SOLAR-1 study of PI3K inhibitor alpelisib plus fulvestrant in patients with HR-positive advanced breast cancer. *Ann Oncol* 2020;31:1001–1010
109. Aroda VR, Edelstein SL, Goldberg RB, et al.; Diabetes Prevention Program Research Group. Long-term metformin use and vitamin B12 deficiency in the Diabetes Prevention Program Outcomes Study. *J Clin Endocrinol Metab* 2016;101:1754–1761
110. Kim J, Ahn CW, Fang S, Lee HS, Park JS. Association between metformin dose and vitamin B12 deficiency in patients with type 2 diabetes. *Medicine (Baltimore)* 2019;98:e17918
111. de Jager J, Kooy A, Leher P, et al. Long term treatment with metformin in patients with type 2 diabetes and risk of vitamin B-12 deficiency: randomised placebo controlled trial. *BMJ* 2010;340:c2181
112. Griffin SJ, Bethel MA, Holman RR, et al. Metformin in non-diabetic hyperglycaemia: the GLINT feasibility RCT. *Health Technol Assess* 2018;22:1–64
113. Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2024 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int* 2024;105:S117–S314
114. Jung C, Park S, Kim H. Concomitant use of metformin and proton pump inhibitors increases vitamin B12 deficiency risk in type 2 diabetes. *J Diabetes Investig* 2025;16:1020–1027
115. Chapman LE, Darling AL, Brown JE. Association between metformin and vitamin B12 deficiency in patients with type 2 diabetes: a systematic review and meta-analysis. *Diabetes Metab* 2016;42:316–327
116. Ali MK, Bullard KM, Saydah S, Imperatore G, Gregg EW. Cardiovascular and renal burdens of prediabetes in the USA: analysis of data from serial cross-sectional surveys, 1988–2014. *Lancet Diabetes Endocrinol* 2018;6:392–403
117. Pan Y, Chen W, Wang Y. Prediabetes and outcome of ischemic stroke or transient ischemic attack: a systematic review and meta-analysis. *J Stroke Cerebrovasc Dis* 2019;28:683–692
118. Huang Y, Cai X, Mai W, Li M, Hu Y. Association between prediabetes and risk of cardiovascular disease and all cause mortality: systematic review and meta-analysis. *BMJ* 2016;355:i5953
119. Yeh H-C, Duncan BB, Schmidt MI, Wang N-Y, Brancati FL. Smoking, smoking cessation, and risk for type 2 diabetes mellitus: a cohort study. *Ann Intern Med* 2010;152:10–17
120. Hu Y, Zong G, Liu G, et al. Smoking cessation, weight change, type 2 diabetes, and mortality. *N Engl J Med* 2018;379:623–632
121. Orchard TJ, Tempresa M, Barrett-Connor E, et al. Long-term effects of the Diabetes Prevention Program interventions on cardiovascular risk factors:

a report from the DPP Outcomes Study. *Diabet Med* 2013;30:46–55

122. Salas-Salvadó J, Díaz-López A, Ruiz-Canela M, et al.; PREDIMED-Plus investigators. Effect of a lifestyle intervention program with energy-restricted Mediterranean diet and exercise on weight loss and cardiovascular risk factors: one-year results of the PREDIMED-Plus trial. *Diabetes Care* 2019;42:777–788

123. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA Guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019;140:e596–e646

124. Crandall JP, Mather K, Rajpathak SN, et al. Statin use and risk of developing diabetes: results from the Diabetes Prevention Program. *BMJ Open Diabetes Res Care* 2017;5:e000438

125. Cholesterol Treatment Trialists' (CTT) Collaboration. Effects of statin therapy on diagnoses of new-onset diabetes and worsening glycaemia in large-scale randomised blinded statin trials: an individual participant data meta-analysis. *Lancet Diabetes Endocrinol* 2024;12:306–319

126. Ridker PM, Pradhan A, MacFadyen JG, Libby P, Glynn RJ. Cardiovascular benefits and diabetes risks of statin therapy in primary prevention: an analysis from the JUPITER trial. *Lancet* 2012;380:565–571

127. Cai T, Abel L, Langford O, et al. Associations between statins and adverse events in primary prevention of cardiovascular disease: systematic review with pairwise, network, and dose-response meta-analyses. *BMJ* 2021;374:n1537

128. Kernan WN, Viscoli CM, Furie KL, et al.; IRIS Trial Investigators. Pioglitazone after ischemic

stroke or transient ischemic attack. *N Engl J Med* 2016;374:1321–1331

129. Inzucchi SE, Viscoli CM, Young LH, et al.; IRIS Trial Investigators. Pioglitazone prevents diabetes in patients with insulin resistance and cerebrovascular disease. *Diabetes Care* 2016;39:1684–1692

130. Spence JD, Viscoli CM, Inzucchi SE, et al.; IRIS Investigators. Pioglitazone therapy in patients with stroke and prediabetes: a post hoc analysis of the IRIS randomized clinical trial. *JAMA Neurol* 2019;76:526–535

131. Spence JD, Viscoli C, Kernan WN, et al. Efficacy of lower doses of pioglitazone after stroke or transient ischaemic attack in patients with insulin resistance. *Diabetes Obes Metab* 2022;24:1150–1158

132. Nadeau KJ, Anderson BJ, Berg EG, et al. Youth-onset type 2 diabetes consensus report: current status, challenges, and priorities. *Diabetes Care* 2016;39:1635–1642

133. Rooney MR, Rawlings AM, Pankow JS, et al. Risk of progression to diabetes among older adults with prediabetes. *JAMA Intern Med* 2021;181:511–519

134. Cigolle CT, Blaum CS, Lyu C, Ha J, Kabeto M, Zhong J. Associations of age at diagnosis and duration of diabetes with morbidity and mortality among older adults. *JAMA Netw Open* 2022;5:e2232766

135. Lachin JM, Christophi CA, Edelstein SL, et al.; DDK Research Group. Factors associated with diabetes onset during metformin versus placebo therapy in the diabetes prevention program. *Diabetes* 2007;56:1153–1159

136. Perreault L, Pan Q, Schroeder EB, et al.; Diabetes Prevention Program Research Group. Regression from prediabetes to normal glucose regulation and prevalence of microvascular

disease in the Diabetes Prevention Program Outcomes Study (DPPOS). *Diabetes Care* 2019;42:1809–1815

137. Chen Y, Zhang P, Wang J, et al. Associations of progression to diabetes and regression to normal glucose tolerance with development of cardiovascular and microvascular disease among people with impaired glucose tolerance: a secondary analysis of the 30 year Da Qing Diabetes Prevention Outcome Study. *Diabetologia* 2021;64:1279–1287

138. Liu X, Johnson SB, Lynch KF, et al.; TEDDY Study Group. Physical activity and the development of islet autoimmunity and type 1 diabetes in 5- to 15-year-old children followed in the TEDDY study. *Diabetes Care* 2023;46:1409–1416

139. Lamb MM, Yin X, Barriga K, et al. Dietary glycemic index, development of islet autoimmunity, and subsequent progression to type 1 diabetes in young children. *J Clin Endocrinol Metab* 2008;93:3936–3942

140. Lamb MM, Frederiksen B, Seifert JA, Kroehl M, Rewers M, Norris JM. Sugar intake is associated with progression from islet autoimmunity to type 1 diabetes: the Diabetes Autoimmunity Study in the Young. *Diabetologia* 2015;58:2027–2034

141. Li X, Wang L, Meng G, et al. Sustained high glucose intake accelerates type 1 diabetes in NOD mice. *Front Endocrinol (Lausanne)* 2022;13:1037822

142. Herold KC, Bundy BN, Long SA, et al.; Type 1 Diabetes TrialNet Study Group. An anti-CD3 antibody, teplizumab, in relatives at risk for type 1 diabetes. *N Engl J Med* 2019;381:603–613