NOVACAINE™

Anesthetic-Antacid Gel

COMPOSITION (Per 5ml) :
Oxetacaine IP 10mg + Aluminium Hydroxide IP 291mg + Magnesium Hydroxide IP 98 mg
Sugar Free Oral Gel (Contains Sorbitol)

An anesthetic–Antacid gel offering immediate symptomatic relief with sustained control of Gastric Acidity, Especially useful in Painful Acidity and Gastritis.

Mechanism of Action

Oxetacaine : Local anesthetic providing rapid relief from epigastric pain and burning; remains effective even in highly acidic pH.
Aluminium Hydroxide : Slow-acting antacid for sustained acid neutralization.
Magnesium Hydroxide : Fast-acting antacid; balances aluminium-induced constipation.
Overall Effect : Immediate pain relief with prolonged acid control.



200ml

SUGAR FREE



Mint FLAVOUR

DOSAGE & ADMINISTRATION :
5-10 ml, 3-4 times daily, Preferably 15 minutes before meals & at bedtime.
To be taken undiluted or as directed by the physician.

CLINICAL INDICATIONS

- Acid Peptic Disease & Gastritis
- GERD / Reflux Esophagitis
- Peptic Ulcer Disease
- Painful Hyperacidity & Heartburn
- Drug-Induced Gastric Irritation (NSAIDs, Steroids)

KEY ADVANTAGES

- Rapid relief from Burning and Pain
- Sustained Antacid Action
- Gel formulation for better Mucosal Coating
- Sugar-Free – Suitable for Diabetic Patients
- Good GI Tolerability