

Lecture 1: Basics of Graph Theory


*Instructor: Vinod Reddy**Scribe: GVN Sai Uday(11740320) & E Suyra Teja(11740310)*

In this lecture, we will be dealing with some basic terminology required to learn Graph Theory.

A graph G is a pair $G = (V, E)$, where V is a finite set and E is a set of 2-element subsets of V . The elements of V are called vertices of the graph G and the elements of E are called edges of G .

For a graph $G = (V, E)$, let $V(G)$ & $E(G)$ denote the vertex and edge set of G respectively.

Example:-



$$V = \{1, 2, 3, 4\}$$
$$E = \{\{1, 2\}, \{1, 4\}, \{2, 3\}, \{2, 4\}\}$$

The number of vertices of a graph G is its order(or size), denoted as $|G|$ or $|V|$.

Note:-

- From this lecture onwards n denotes the number of vertices in a graph.
- Also m denotes the number of edges in a graph.
- The graphs we consider in this course are finite unless otherwise stated.

An edge x, y is usually written as xy (or yx). Two vertices x, y of G are adjacent or neighbours if xy is an edge of G . Two edges e, f are adjacent if they have an end in common.

Definition 1.1 (Degree(or Valency)). *The degree(or valency) of a vertex v in a graph G is the number of edges incident with v , with loops counted twice. It is denoted by $\deg_G(v)$ or $d(v)$.*

Lemma 1.2 (Handshaking Lemma). *The sum of the degrees of the vertices of a graph equals twice the number of edges.*

Proof. Each edge contributes two to the sum of the degrees, one for each of its endpoints.

$$2|E| = \sum_{v \in V} d(v)$$

□

Corollary 1.3. *In any graph G , the number of vertices of odd degree is even.*

Proof. We know that

$$2|E| = \sum_{v \in V} d(v)$$

$$\underbrace{2|E|}_{\text{EVEN}} = \underbrace{\sum_{\substack{v \in V \\ d(v) \text{ is odd}}} d(v)}_{\mathbf{x}} + \underbrace{\sum_{\substack{v \in V \\ d(v) \text{ is even}}} d(v)}_{\text{EVEN}}$$

For the above equation to satisfy the value of \mathbf{x} should be even (i.e. for the LHS to be even sum of terms of RHS should be even, which is only possible when value of \mathbf{x} is even since the other term is even.) □

Definition 1.4 (Isolated vertex). *A vertex of degree zero is called isolated vertex.*


Definition 1.5 (Minimum degree). *The degree of the vertex with the least number of edges incident to it (i.e. $\min\{d(v) : v \in V\}$) is the minimum degree of G . It is represented by $\delta(G)$*



Definition 1.6 (Maximum degree). *The degree of the vertex with the most number of edges incident to it (i.e. $\max\{d(v) : v \in V\}$) is the maximum degree of G . It is represented by $\Delta(G)$*

Definition 1.7 (Regular Graph). *If all the vertices of G have the same degree k then the graph G is called k -regular.*

Example:-

(a) 2-Regular Graph



(b) Petersen's Graph(3-Regular Graph)**(c) 3-Regular Graph****1.1 Some theorems and stuff**

We now delve right into the proof.

Lemma 1.8. *This is the first lemma of the lecture.*

Proof. The proof is by induction on \dots . We also throw in a figure (which you might want to make larger).

This is the end of the proof, which is marked with a little box. □

1.1.1 A few items of note

Here is an itemized list:

- this is the first item;

- this is the second item.

Here is an enumerated list:

1. this is the first item;
2. this is the second item.

Here is an exercise:

Exercise: Find an efficient algorithm for triangulation.

Here is how to define things in the proper mathematical style. Let f_k be the *AND – OR* function, defined by

$$f_k(x_1, x_2, \dots, x_{2^k}) = \begin{cases} x_1 & \text{if } k = 0; \\ \text{AND}(f_{k-1}(x_1, \dots, x_{2^{k-1}}), f_{k-1}(x_{2^{k-1}+1}, \dots, x_{2^k})) & \text{if } k \text{ is even;} \\ \text{OR}(f_{k-1}(x_1, \dots, x_{2^{k-1}}), f_{k-1}(x_{2^{k-1}+1}, \dots, x_{2^k})) & \text{otherwise.} \end{cases}$$

Here is another equation that uses one of the AMS commands, align

$$\begin{aligned} p(x_{1:5}) &= \sum_{x_{2:5}} p(x_1) p(x_2|x_1) p(x_3|x_1) p(x_4|x_2) p(x_5|x_1) p(x_6|x_2, x_5) \\ &= p(x_1) \sum_{x_2} p(x_2|x_1) \sum_{x_3} p(x_3|x_1) \sum_{x_4} p(x_4|x_2) \sum_{x_5} p(x_5|x_1) p(x_6|x_2, x_5) \\ &= p(x_1) \sum_{x_2} p(x_2|x_1) \sum_{x_3} p(x_3|x_1) \sum_{x_4} p(x_4|x_2) \sum_{x_5} p(x_5|x_1) \phi_{X_6}(x_2, x_5) \end{aligned}$$

which assumes that $X_4 \perp\!\!\!\perp \{X_1, X_3\} | X_2$.

Theorem 1.9. *This is the first theorem.*

Proof. This is the proof of the first theorem. We show how to write pseudo-code now.

Consider a comparison between x and y :

```

if  $x$  or  $y$  or both are in  $S$  then
  answer accordingly
else
  Make the element with the larger score (say  $x$ ) win the comparison
  if  $F(x) + F(y) < \frac{n}{t-1}$  then
     $F(x) \leftarrow F(x) + F(y)$ 
     $F(y) \leftarrow 0$ 
  else
     $S \leftarrow S \cup \{x\}$ 
     $r \leftarrow r + 1$ 
  endif
endif

```

This concludes the proof. □

1.2 Next topic

Here is some citations [JB00] and [L2000]

References

[JB00] M.I. JORDAN and C. BISHOP, “An Introduction to Graphical Models,” *To be published*, 2000

[L2000] S.L. LAURITZEN “Graphical Models,” *Oxford Science Publications*, 1996