


Stirling numbers of the second kind

In [mathematics](#), particularly in [combinatorics](#), a **Stirling number of the second kind** (or Stirling partition number) is the number of ways to [partition](#) a [set](#) of n objects into k non-empty subsets and is denoted by $S(n, k)$ or $\{n \atop k\}$.^[1] Stirling numbers of the second kind occur in the field of [mathematics](#) called [combinatorics](#) and the study of [partitions](#).

Stirling numbers of the second kind are one of two kinds of [Stirling numbers](#), the other kind being called [Stirling numbers of the first kind](#) (or Stirling cycle numbers). Mutually inverse (finite or infinite) [triangular matrices](#) can be formed from the Stirling numbers of each kind according to the parameters n, k .



The 15 partitions of a 4-element set ordered in a Hasse diagram

There are $S(4,1), \dots, S(4,4) = 1, 7, 6, 1$ partitions containing 1, 2, 3, 4 sets.

Contents

Definition

Notation

Relation to Bell numbers

Table of values

Properties

- Recurrence relation
- Lower and upper bounds
- Maximum
- Parity
- Simple identities
- Explicit formula
- Generating functions
- Asymptotic approximation

Applications

- Moments of the Poisson distribution
- Moments of fixed points of random permutations
- Rhyming schemes

Variants

- Associated Stirling numbers of the second kind
- Reduced Stirling numbers of the second kind

See also

References

Definition

The Stirling numbers of the second kind, written $S(n, k)$ or $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ or with other notations, count the number of ways to partition a set of n labelled objects into k nonempty unlabelled subsets. Equivalently, they count the number of different equivalence relations with precisely k equivalence classes that can be defined on an n element set. In fact, there is a bijection between the set of partitions and the set of equivalence relations on a given set. Obviously,

$$\left\{ \begin{smallmatrix} n \\ n \end{smallmatrix} \right\} = 1 \text{ and for } n \geq 1, \left\{ \begin{smallmatrix} n \\ 1 \end{smallmatrix} \right\} = 1$$

as the only way to partition an n -element set into n parts is to put each element of the set into its own part, and the only way to partition a nonempty set into one part is to put all of the elements in the same part. They can be calculated using the following explicit formula:^[2]

$$\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n.$$

The Stirling numbers of the second kind may also be characterized as the numbers that arise when one expresses powers of an indeterminate x in terms of the falling factorials^[3]

$$(x)_n = x(x-1)(x-2) \cdots (x-n+1).$$

(In particular, $(x)_0 = 1$ because it is an empty product.) In particular, one has

$$\sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} (x)_k = x^n.$$

Notation

Various notations have been used for Stirling numbers of the second kind. The brace notation $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ was used by Imanuel Marx and Antonio Salmeri in 1962 for variants of these numbers.^{[4][5]} This led Knuth to use it, as shown here, in the first volume of *The Art of Computer Programming* (1968).^{[6][7]} However, according to the third edition of *The Art of Computer Programming*, this notation was also used earlier by Jovan Karamata in 1935.^{[8][9]} The notation $S(n, k)$ was used by Richard Stanley in his book *Enumerative Combinatorics*.^[6]

Relation to Bell numbers

Since the Stirling number $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ counts set partitions of an n -element set into k parts, the sum

$$B_n = \sum_{k=0}^n \left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$$

over all values of k is the total number of partitions of a set with n members. This number is known as the n th Bell number.

Analogously, the ordered Bell numbers can be computed from the Stirling numbers of the second kind via

$$a_n = \sum_{k=0}^n k! \left\{ \begin{matrix} n \\ k \end{matrix} \right\} .^{[10]}$$

Table of values

Below is a triangular array of values for the Stirling numbers of the second kind (sequence [A008277](#) in the [OEIS](#)):

$n \backslash k$	0	1	2	3	4	5	6	7	8	9	10
0	1										
1	0	1									
2	0	1	1								
3	0	1	3	1							
4	0	1	7	6	1						
5	0	1	15	25	10	1					
6	0	1	31	90	65	15	1				
7	0	1	63	301	350	140	21	1			
8	0	1	127	966	1701	1050	266	28	1		
9	0	1	255	3025	7770	6951	2646	462	36	1	
10	0	1	511	9330	34105	42525	22827	5880	750	45	1

As with the binomial coefficients, this table could be extended to $k > n$, but those entries would all be 0.

Properties

Recurrence relation

Stirling numbers of the second kind obey the recurrence relation

$$\left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\} = k \left\{ \begin{matrix} n \\ k \end{matrix} \right\} + \left\{ \begin{matrix} n \\ k-1 \end{matrix} \right\}$$

for $k > 0$ with initial conditions

$$\left\{ \begin{matrix} 0 \\ 0 \end{matrix} \right\} = 1 \quad \text{and} \quad \left\{ \begin{matrix} n \\ 0 \end{matrix} \right\} = \left\{ \begin{matrix} 0 \\ n \end{matrix} \right\} = 0$$

for $n > 0$.

For instance, the number 25 in column $k=3$ and row $n=5$ is given by $25=7+(3 \times 6)$, where 7 is the number above and to the left of 25, 6 is the number above 25 and 3 is the column containing the 6.

To understand this recurrence, observe that a partition of the $n+1$ objects into k nonempty subsets either contains the

$n+1$ -th object as a singleton or it does not. The number of ways that the singleton is one of the subsets is given by

$$\left\{ \begin{matrix} n \\ k-1 \end{matrix} \right\}$$

since we must partition the remaining n objects into the available $k-1$ subsets. In the other case the $n+1$ -th object belongs to a subset containing other objects. The number of ways is given by

$$k \left\{ \begin{matrix} n \\ k \end{matrix} \right\}$$

since we partition all objects other than the $n+1$ -th into k subsets, and then we are left with k choices for inserting object $n+1$. Summing these two values gives the desired result.

Some more recurrences are as follows:

$$\left\{ \begin{matrix} n+1 \\ k+1 \end{matrix} \right\} = \sum_{j=k}^n \binom{n}{j} \left\{ \begin{matrix} j \\ k \end{matrix} \right\},$$

$$\left\{ \begin{matrix} n+1 \\ k+1 \end{matrix} \right\} = \sum_{j=k}^n (k+1)^{n-j} \left\{ \begin{matrix} j \\ k \end{matrix} \right\},$$

$$\left\{ \begin{matrix} n+k+1 \\ k \end{matrix} \right\} = \sum_{j=0}^k j \left\{ \begin{matrix} n+j \\ j \end{matrix} \right\}.$$

Lower and upper bounds

If $n \geq 2$ and $1 \leq k \leq n-1$, then

$$L(n, k) \leq \left\{ \begin{matrix} n \\ k \end{matrix} \right\} \leq U(n, k)$$

where

$$L(n, k) = \frac{1}{2}(k^2 + k + 2)k^{n-k-1} - 1$$

and

$$U(n, k) = \frac{1}{2} \binom{n}{k} k^{n-k}. \text{ [11]}$$

Maximum

For fixed n , $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}$ has a single maximum, which is attained for at most two consecutive values of k . That is, there is an integer K_n such that

$$\left\{ \begin{matrix} n \\ 1 \end{matrix} \right\} < \left\{ \begin{matrix} n \\ 2 \end{matrix} \right\} < \cdots < \left\{ \begin{matrix} n \\ K_n \end{matrix} \right\},$$

$$\left\{ \begin{matrix} n \\ K_n \end{matrix} \right\} \geq \left\{ \begin{matrix} n \\ K_n + 1 \end{matrix} \right\} > \cdots > \left\{ \begin{matrix} n \\ n \end{matrix} \right\}.$$

When n is large

$$K_n \sim \frac{n}{\log n},$$

and the maximum value of the Stirling number of second kind is

$$\log \left\{ \begin{matrix} n \\ K_n \end{matrix} \right\} = n \log n - n \log \log n - n + O(n \log \log n / \log n). \quad [11]$$

Parity

The parity of a Stirling number of the second kind is equal to the parity of a related binomial coefficient:

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\} \equiv \binom{z}{w} \pmod{2}, \text{ where}$$

$$z = n - \left\lceil \frac{k+1}{2} \right\rceil, \quad w = \left\lfloor \frac{k-1}{2} \right\rfloor.$$

This relation is specified by mapping n and k coordinates onto the Sierpiński triangle.

More directly, let two sets contain positions of 1's in binary representations of results of respective expressions:

$$\mathbb{A} : \sum_{i \in \mathbb{A}} 2^i = n - k,$$


$$\mathbb{B} : \sum_{j \in \mathbb{B}} 2^j = \left\lfloor \frac{k-1}{2} \right\rfloor.$$

One can mimic a bitwise AND operation by intersecting these two sets:

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\} \bmod 2 = \begin{cases} 0, & \mathbb{A} \cap \mathbb{B} \neq \emptyset; \\ 1, & \mathbb{A} \cap \mathbb{B} = \emptyset; \end{cases}$$

to obtain the parity of a Stirling number of the second kind in $O(1)$ time. In pseudocode:

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\} \bmod 2 := [((n - k) \& ((k - 1) \operatorname{div} 2)) = 0];$$



Parity of Stirling numbers of the second kind.

where $[b]$ is the Iverson bracket.

Simple identities

Some simple identities include

$$\left\{ \begin{matrix} n \\ n-1 \end{matrix} \right\} = \binom{n}{2}.$$

This is because dividing n elements into $n - 1$ sets necessarily means dividing it into one set of size 2 and $n - 2$ sets of size 1. Therefore we need only pick those two elements;

and

$$\left\{ \begin{matrix} n \\ 2 \end{matrix} \right\} = 2^{n-1} - 1.$$

To see this, first note that there are 2^n *ordered* pairs of complementary subsets A and B . In one case, A is empty, and in another B is empty, so $2^n - 2$ ordered pairs of subsets remain. Finally, since we want *unordered* pairs rather than *ordered* pairs we divide this last number by 2, giving the result above.

Another explicit expansion of the recurrence-relation gives identities in the spirit of the above example.

$$\begin{aligned} \left\{ \begin{matrix} n \\ 2 \end{matrix} \right\} &= \frac{\frac{1}{1}(2^{n-1} - 1^{n-1})}{0!} \\ \left\{ \begin{matrix} n \\ 3 \end{matrix} \right\} &= \frac{\frac{1}{1}(3^{n-1} - 2^{n-1}) - \frac{1}{2}(3^{n-1} - 1^{n-1})}{1!} \\ \left\{ \begin{matrix} n \\ 4 \end{matrix} \right\} &= \frac{\frac{1}{1}(4^{n-1} - 3^{n-1}) - \frac{2}{2}(4^{n-1} - 2^{n-1}) + \frac{1}{3}(4^{n-1} - 1^{n-1})}{2!} \\ \left\{ \begin{matrix} n \\ 5 \end{matrix} \right\} &= \frac{\frac{1}{1}(5^{n-1} - 4^{n-1}) - \frac{3}{2}(5^{n-1} - 3^{n-1}) + \frac{3}{3}(5^{n-1} - 2^{n-1}) - \frac{1}{4}(5^{n-1} - 1^{n-1})}{3!} \\ &\vdots \end{aligned}$$

Explicit formula

The Stirling numbers of the second kind are given by the explicit formula:

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\} = \sum_{j=1}^k (-1)^{k-j} \frac{j^{n-1}}{(j-1)!(k-j)!} = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n.$$

This formula is a special case of the k th forward difference of the monomial x^n evaluated at $x = 0$:

$$\Delta^k x^n = \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} (x+j)^n.$$

Because the [Bernoulli polynomials](#) may be written in terms of these forward differences, one immediately obtains a relation in the [Bernoulli numbers](#):

$$B_m(0) = \sum_{k=0}^m \frac{(-1)^k k!}{k+1} \left\{ \begin{matrix} m \\ k \end{matrix} \right\}.$$

Generating functions

For a fixed integer n , the [ordinary generating function](#) for the Stirling numbers of the second kind $\left\{ \begin{matrix} n \\ 0 \end{matrix} \right\}, \left\{ \begin{matrix} n \\ 1 \end{matrix} \right\}, \dots$ is given by

$$\sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k = T_n(x),$$

where $T_n(x)$ are [Touchard polynomials](#). If one sums the Stirling numbers against the falling factorial instead, one can show the following identities, among others:

$$\sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (x)_k = x^n$$

and

$$\sum_{k=1}^n \left\{ \begin{matrix} n+1 \\ k \end{matrix} \right\} (x-1)_{k-1} = x^n.$$

For a fixed integer k , the Stirling numbers of the second kind $\left\{ \begin{matrix} 0 \\ k \end{matrix} \right\}, \left\{ \begin{matrix} 1 \\ k \end{matrix} \right\}, \dots$ have rational ordinary generating function

$$\sum_{n=k}^{\infty} \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^{n-k} = \prod_{r=1}^k \frac{1}{1-rx} = \frac{1}{(k+1)! x^{k+1} \left(\frac{1}{x} \right)_{k+1}}$$

and have [exponential generating function](#) given by

$$\sum_{n=k}^{\infty} \left\{ \begin{matrix} n \\ k \end{matrix} \right\} \frac{x^n}{n!} = \frac{(e^x - 1)^k}{k!}.$$

A mixed bivariate generating function for the Stirling numbers of the second kind is

$$\sum_{0 \leq k \leq n} \left\{ \begin{matrix} n \\ k \end{matrix} \right\} \frac{x^n}{n!} y^k = e^{y(e^x - 1)}.$$

Asymptotic approximation

For fixed value of k , the asymptotic value of the Stirling numbers of the second kind as $n \rightarrow \infty$ is given by

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\} \sim \frac{k^n}{k!}.$$

On the other side, if $k = o(\sqrt{n})$ (where o denotes the little o notation) then

$$\left\{ \begin{matrix} n \\ n - k \end{matrix} \right\} \sim \frac{(n - k)^{2k}}{2^k k!} \left(1 + \frac{1}{3} \frac{2k^2 + k}{n - k} + \frac{1}{18} \frac{4k^4 - k^2 - 3k}{(n - k)^2} + \dots \right).^{[12]}$$

Uniformly valid approximation also exist: for all k such that $1 < k < n$, one has

$$\left\{ \begin{matrix} n \\ k \end{matrix} \right\} \sim \frac{\sqrt{n - k}}{\sqrt{n(1 - G)} G^k (v - G)^{n - k}} \left(\frac{n - k}{e} \right)^{n - k} \binom{n}{k},$$

where $G = -W_0(-ve^{-v})$, $v = n/k$, $W_0(z)$ is main branch of Lambert W function.^{[13][14]} Relative error is bounded by about $0.066/n$.

Applications

Moments of the Poisson distribution

If X is a random variable with a Poisson distribution with expected value λ , then its n th moment is

$$E(X^n) = \sum_{k=1}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} \lambda^k.$$

In particular, the n th moment of the Poisson distribution with expected value 1 is precisely the number of partitions of a set of size n , i.e., it is the n th Bell number (this fact is Dobiński's formula).

Moments of fixed points of random permutations

Let the random variable X be the number of fixed points of a uniformly distributed random permutation of a finite set of size m . Then the n th moment of X is

$$E(X^n) = \sum_{k=1}^m \left\{ \begin{matrix} n \\ k \end{matrix} \right\}.$$

Note: The upper bound of summation is m , not n .

In other words, the n th moment of this [probability distribution](#) is the number of partitions of a set of size n into no more than m parts. This is proved in the article on [random permutation statistics](#), although the notation is a bit different.

Rhyming schemes

The Stirling numbers of the second kind can represent the total number of [rhyme schemes](#) for a poem of n lines. $S(n, k)$ gives the number of possible rhyming schemes for n lines using k unique rhyming syllables. As an example, for a poem of 3 lines, there is 1 rhyme scheme using just one rhyme (aaa), 3 rhyme schemes using two rhymes (aab, aba, abb), and 1 rhyme scheme using three rhymes (abc).

Variants

Associated Stirling numbers of the second kind

An r -associated Stirling number of the second kind is the number of ways to partition a set of n objects into k subsets, with each subset containing at least r elements.^[15] It is denoted by $S_r(n, k)$ and obeys the recurrence relation

$$S_r(n + 1, k) = k S_r(n, k) + \binom{n}{r-1} S_r(n - r + 1, k - 1)$$

The 2-associated numbers (sequence [A008299](#) in the [OEIS](#)) appear elsewhere as "Ward numbers" and as the magnitudes of the coefficients of [Mahler polynomials](#).


Reduced Stirling numbers of the second kind

Denote the n objects to partition by the integers 1, 2, ..., n . Define the reduced Stirling numbers of the second kind, denoted $S^d(n, k)$, to be the number of ways to partition the integers 1, 2, ..., n into k nonempty subsets such that all elements in each subset have pairwise distance at least d . That is, for any integers i and j in a given subset, it is required that $|i - j| \geq d$. It has been shown that these numbers satisfy

$$S^d(n, k) = S(n - d + 1, k - d + 1), n \geq k \geq d$$

(hence the name "reduced").^[16] Observe (both by definition and by the reduction formula), that $S^1(n, k) = S(n, k)$, the familiar Stirling numbers of the second kind.

See also

- [Bell number](#) – the number of partitions of a set with n members
- [Stirling numbers of the first kind](#)
- [Stirling polynomials](#)
- [Twelffold way](#)
-  [Partition related number triangles](#)

References

1. Ronald L. Graham, Donald E. Knuth, Oren Patashnik (1988) *Concrete Mathematics*, Addison–Wesley, Reading MA. ISBN 0-201-14236-8, p. 244.
2. "Stirling Number of the Second Kind" (<http://mathworld.wolfram.com/StirlingNumberoftheSecondKind.html>).
3. Confusingly, the notation that combinatorialists use for *falling* factorials coincides with the notation used in *special functions* for *rising* factorials; see Pochhammer symbol.
4. Transformation of Series by a Variant of Stirling's Numbers, Imanuel Marx, *The American Mathematical Monthly* **69**, #6 (June–July 1962), pp. 530–532, JSTOR 2311194 (<https://www.jstor.org/stable/2311194>).
5. Antonio Salmeri, Introduzione alla teoria dei coefficienti fattoriali, *Giornale di Matematiche di Battaglini* **90** (1962), pp. 44–54.
6. Knuth, D.E. (1992), "Two notes on notation", *Amer. Math. Monthly*, **99**: 403–422, arXiv:math/9205211 (<https://arxiv.org/abs/math/9205211>), doi:10.2307/2325085 (<https://doi.org/10.2307%2F2325085>), JSTOR 2325085 (<https://www.jstor.org/stable/2325085>)
7. Donald E. Knuth, *Fundamental Algorithms*, Reading, Mass.: Addison–Wesley, 1968.
8. p. 66, Donald E. Knuth, *Fundamental Algorithms*, 3rd ed., Reading, Mass.: Addison–Wesley, 1997.
9. Jovan Karamata, Théorèmes sur la sommabilité exponentielle et d'autres sommabilités s'y rattachant, *Mathematica* (Cluj) **9** (1935), pp. 164–178.
10. Sprugnoli, Renzo (1994), "Riordan arrays and combinatorial sums", *Discrete Mathematics*, **132** (1–3): 267–290, doi:10.1016/0012-365X(92)00570-H (<https://doi.org/10.1016%2F0012-365X%2892%2900570-H>), MR 1297386 (<https://www.ams.org/mathscinet-getitem?mr=1297386>)
11. B.C. Rennie, A.J. Dobson. "On Stirling Numbers of the Second Kind" (http://ac.els-cdn.com/S0021980069800451/1-s2.0-S0021980069800451-main.pdf?_tid=4e52b648-711a-11e2-a90d-00000aab0f26&acdnat=1360237038_c9d4f38f83a536ed624e29e3554016e2)
12. L. C. Hsu, Note on an Asymptotic Expansion of the nth Difference of Zero, AMS Vol.19 NO.2 1948, pp. 273--277
13. W. E. Bleick and Peter C. C. Wang, Asymptotics of Stirling Numbers of the Second Kind, Proceedings of the AMS Vol.42 No.2, 1974.
14. N. M. Temme, Asymptotic Estimates of Stirling Numbers, STUDIES IN APPLIED MATHEMATICS 89:233-243 (1993), Elsevier Science Publishing.
15. L. Comtet, *Advanced Combinatorics*, Reidel, 1974, p. 222.
16. A. Mohr and T.D. Porter, *Applications of Chromatic Polynomials Involving Stirling Numbers* (<http://www.austinmohr.com/work/files/stirling.pdf>), Journal of Combinatorial Mathematics and Combinatorial Computing **70** (2009), 57–64.
 - Khristo N. Boyadzhiev (2012). "Close encounters with the Stirling numbers of the second kind". *Mathematics Magazine*. **85** (4): 252–266.
 - "Stirling numbers of the second kind, S(n,k)" (<http://planetmath.org/?op=getobj&from=objects&id=2805>). *PlanetMath*.
 - Weisstein, Eric W. "Stirling Number of the Second Kind" (<http://mathworld.wolfram.com/StirlingNumberoftheSecondKind.html>). *MathWorld*.
 - Calculator for Stirling Numbers of the Second Kind (http://austinmohr.com/home/?page_id=431)
 - Set Partitions: Stirling Numbers (<http://dlmf.nist.gov/26.8#vii>)
 - Jack van der Elsen (2005). *Black and white transformations*. Maastricht. ISBN 90-423-0263-1.

Retrieved from "https://en.wikipedia.org/w/index.php?title=Stirling_numbers_of_the_second_kind&oldid=862710517"

This page was last edited on 6 October 2018, at 05:29 (UTC).

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia](#)

Foundation, Inc., a non-profit organization.