

Tamil Nadu - Key Indicators

Indicators	NFHS-6 (2023-24)			NFHS-5 (2019-21)
	Urban	Rural	Total	Total
Population and Household Profile				
1. Population below age 5 years (%)	5.8	6.5	6.2	6.7
2. Population below age 15 years (%)	19.1	21.0	20.2	21.0
3. Population age 60 years and above (%)	16.6	16.1	16.3	14.9
4. Population living in households with electricity (%)	99.6	99.1	99.3	99.3
5. Population living in households with an improved drinking-water source ¹ (%)	99.3	97.9	98.5	98.6
6. Households using iodized salt (%)	94.1	91.7	92.8	92.0
7. Households with any usual member covered under a health insurance/financing scheme (%)	56.7	64.7	61.1	66.5
8. Households with any usual member having a bank account/post office account (%)	96.4	98.0	97.3	96.6
9. Female population age 6 years and above who ever attended school (%)	86.1	75.8	80.4	80.4
10. Households with any usual female members owning a house and/or land (alone or jointly with others) (%)	14.3	19.5	17.2	17.2
11. Children age 2-4 years who attended pre-school (%)	63.7	66.5	65.3	46.1
Characteristics of Adults (age 15-49 years)				
12. Women with 10 or more years of schooling (%)	70.7	59.2	64.2	56.6
13. Men with 10 or more years of schooling (%)	72.8	66.2	69.2	59.1
14. Women who have ever used the internet (%)	67.9	52.8	59.5	46.9
15. Men who have ever used the internet (%)	84.0	80.0	81.8	70.2
Marriage and Fertility				
16. Women age 20-24 years married before age 18 years (%)	9.2	11.5	10.5	12.8
17. Men age 25-29 years married before age 21 years (%)	2.4	4.6	3.7	4.5
18. Total fertility rate (children per woman)	1.6	1.8	1.7	1.8
19. Women age 15-19 years who were already mothers or pregnant at the time of the survey (%)	2.4	5.4	4.1	6.3
Current Use of Family Planning Methods (currently married women age 15-49 years)				
20. Any method ² (%)	67.4	69.4	68.5	68.6
21. Any modern method ² (%)	63.7	66.2	65.1	65.5
22. Any traditional method (%)	3.7	3.2	3.4	3.1
23. Female sterilization (%)	54.5	58.2	56.6	57.8
24. Male sterilization (%)	0.1	0.1	0.1	0.1
Unmet Need for Family Planning (currently married women age 15-49 years)				
25. Total unmet need ³ (%)	6.4	6.4	6.4	7.5
26. Unmet need for spacing ³ (%)	3.2	3.5	3.4	3.0
27. Unmet need for limiting ³ (%)	3.2	2.9	3.0	4.5
Maternal and Child Health				
Antenatal Care (for last birth in the 5 years before the survey)				
28. Mothers who had an antenatal check-up in the first trimester (%)	70.6	71.7	71.2	77.4
29. Mothers who had any antenatal care visits (%)	91.7	92.5	92.2	95.8
30. Mothers who had at least 4 antenatal care visits (%)	86.2	88.6	87.6	90.6
31. Mothers whose last birth was protected against neonatal tetanus ⁴ (%)	86.8	89.3	88.2	89.7
32. Mothers who consumed iron folic acid for 100 days or more when they were pregnant (%)	82.4	82.0	82.1	82.5
33. Mothers who consumed iron folic acid for 180 days or more when they were pregnant (%)	65.7	63.7	64.5	63.1
34. Registered pregnancies for which the mother received a Mother and Child Protection (MCP) card (%)	98.2	99.7	99.1	98.8
Delivery Care (for births in the 5 years before the survey)				
35. Institutional births (%)	99.8	99.6	99.7	99.6
36. Institutional births in public facility (%)	54.2	70.2	63.6	66.9
37. Births attended by skilled health personnel ⁵ (%)	99.7	99.7	99.7	99.8
38. Births delivered by caesarean section (%)	47.8	46.3	46.9	44.9
39. Births in a private health facility that were delivered by caesarean section (%)	58.6	62.1	60.3	63.8
40. Births in a public health facility that were delivered by caesarean section (%)	38.9	40.0	39.6	36.0

Note: Major indicators are highlighted in grey.

*Percentage not shown; based on fewer than 25 unweighted cases

¹Piped water into dwelling/yard/plot, piped to neighbour, public tap/standpipe, tube well or borehole, protected dug well, protected spring, rain water, tanker truck, cart with small tank, bottled water, community RO plant. The main source used for drinking water at the time of interview is recorded.

²Any method includes other methods that are not shown separately; Any modern method includes other modern methods that are not shown separately.

³Unmet need for family planning refers to fecund women who are not using contraception but who wish to postpone the next birth (spacing) or stop childbearing altogether (limiting). Specifically, women are considered to have unmet need for spacing if they are:

- At risk of becoming pregnant, not using contraception, and either do not want to become pregnant within the next two years, or are unsure if or when they want to become pregnant.
- Pregnant with a mistimed pregnancy.
- Postpartum amenorrhoeic for up to two years following a mistimed birth and not using contraception.

Women are considered to have unmet need for limiting if they are:

- At risk of becoming pregnant, not using contraception, and want no (more) children.
- Pregnant with an unwanted pregnancy.
- Postpartum amenorrhoeic for up to two years following an unwanted birth and not using contraception.

⁴Includes mothers with two injections during the pregnancy for their last birth, or two or more injections (the last within 3 years of the last live birth), or three or more injections (the last within 5 years of the last birth), or four or more injections (the last within 10 years of the last live birth), or five or more injections at any time prior to the last birth.

⁵Doctor/nurse/LHV/ANM/midwife/other health personnel.