


Figure 2.1 Results of susceptibility of Enterobacteriaceae to various antibiotics tested.

Table 2.2 Susceptibility of *E. coli*, *Klebsiella* spp and *Enterobacter* spp from blood.

	Amikacin	Cefotax	Ceftazid	Cipro	Colistin	Ertapen	Imipen	Levoflox	Meropen	Pip-taz
	S	S	S	S	S	S	S	S	S	S
E. coli	82	19	24	25	100	69	82	10	77	60
Klebsiella	45	19	22	30	89	37	48	22	42	34
Enterobacter	79	23	33	62	100	68	78	53	74	64
Overall	66	20	23	30	94	54	67	20	62	49




Figure 2.2 Susceptibility of *E. coli*, *Klebsiella*spp and *Enterobacter*spp from blood.

Table 3.2. Comparison of susceptibility of *Escherichia coli* isolated from OPD, ward and ICU

	OPD		Ward		ICU		Total	
	n	%S	n	%S	n	%S	n	%S
Amikacin	1411	87	5311	81	1213	77	7935	81
Cefotaxime	1209	20	4555	15	1071	13	6835	16
Ceftazidime	859	26	3591	17	622	17	5072	19
Ciprofloxacin	1247	26	4838	22	1007	20	7092	22
Colistin	157	100	679	99	229	100	1065	99
Ertapenem	1054	80	3644	70	1031	63	5729	71
Imipenem	1279	79	4808	71	1104	68	7191	72
Levofloxacin	695	20	2424	19	643	17	3762	19
Meropenem	1267	85	5154	75	1078	70	7499	76
Pip-taz	1355	63	5325	52	1210	48	7890	53




Figure 3.3: Comparison of susceptibility of *Escherichia coli* isolated from OPD, ward and ICU.