

Substitution in FOL

Suppose $x \in \text{free}(\varphi)$ and t is any term.

We wish to replace every free occurrence of x in φ with t , such that free variables in t stay free in the resulting formula.

Term t is free for x in φ if no free occurrence of x in φ is in the scope of $\forall y$ or $\exists y$ for any variable y occurring in t .

- $\varphi \triangleq \exists y R(x, y) \vee \forall z R(z, x)$, and t is $f(z, x)$
- $f(z, x)$ is free for x in φ , but $f(y, x)$ is not

$\varphi[t/x]$: Formula obtained by replacing each free occurrence of x in φ by t , if t is free for x in φ

- For φ defined above,
 $\varphi[f(z, x)/x] \triangleq \exists y R(f(z, x), y) \vee \forall z R(z, x)$

$$t = \underline{f(z, z)}$$

$$\exists y \underline{R(x, y)} \vee \forall z \underline{R(z, x)}$$

x z
↓ ↓

bool $R(\text{int } a, \text{ int } b)$
{
}
}

$$x \mapsto \underline{z + x}$$

$$z \mapsto \underline{x * z}$$

result = false

{ for all int y .
 $\underline{z + x}$ is true
if $R(\underline{x}, y)$ then
result = true ; exit loop

result1 = true

for all int z
 $\underline{x * z}$ is false then
result1 = false;
exit loop

return result \vee result1

Semantics of FOL: Some Intuition

$$\varphi \triangleq \forall x \forall y (P(x, y) \rightarrow \exists z (\neg(z = x) \wedge \neg(z = y) \wedge P(x, z) \wedge P(z, y)))$$

Semantics of FOL: Some Intuition

$$\varphi \triangleq \forall x \forall y (\underline{P(x,y)} \rightarrow \exists z (\underline{\neg(z=x)} \wedge \underline{\neg(z=y)} \wedge \underline{P(x,z)} \wedge \underline{P(z,y)}))$$

English reading: For every x and y , if $P(x,y)$ holds, we can find z distinct from x and y such that both $P(x,z)$ and $P(z,y)$ hold.

Semantics of FOL: Some Intuition

$$\varphi \triangleq \forall \underline{x} \forall \underline{y} (P(x, y) \rightarrow \exists z (\neg(z = x) \wedge \neg(z = y) \wedge P(x, z) \wedge P(z, y)))$$

English reading: For every x and y , if $P(x, y)$ holds, we can find z distinct from x and y such that both $P(x, z)$ and $P(z, y)$ hold.

Case 1:

- Variables take values from real numbers
- $P(x, y)$ represents $x < y$
- English reading simply states “real numbers are dense”
- φ is true

Semantics of FOL: Some Intuition

$$\varphi \triangleq \forall x \forall y (P(x, y) \rightarrow \exists z (\neg(z = x) \wedge \neg(z = y) \wedge P(x, z) \wedge P(z, y)))$$

$$x \leq y \rightarrow z \neq x \quad z \neq y \quad \underline{x \leq z \wedge z \leq y}$$

Case 2:

- Variables take values from **real numbers**
- $P(x, y)$ represents $x \leq y$
- English reading requires the following to be true
 - If $x = y$, there is a z such that $z \neq x$, $x \leq z$ and $z \leq x$
 - Thus, $z \neq x$ and $z = x$
- φ is **false**

Semantics of FOL: Some Intuition

$$\varphi \triangleq \forall x \forall y (P(x, y) \rightarrow \exists z (\neg(z = x) \wedge \neg(z = y) \wedge P(x, z) \wedge P(z, y)))$$

Case 3:

- Variables take values from natural numbers
- $P(x, y)$ represents $x < y$
- English reading states that “natural numbers are dense”
- φ is false

Semantics of FOL: Some Intuition

$$\varphi \triangleq \forall x \forall y (P(x, y) \rightarrow \exists z (\neg(z = x) \wedge \neg(z = y) \wedge P(x, z) \wedge P(z, y)))$$

Case 3:

- Variables take values from **natural numbers**
- $P(x, y)$ represents $x < y$
- English reading states that “natural numbers are dense”
- φ is **false**

Truth of φ depends on the underlying set from which variables take values, and on how constants, functions, predicates are interpreted

Semantics of FOL: Formalizing the intuition

Vocabulary \mathcal{V} : E.g. $\mathcal{V} : \{a, f, =, R\}$



Semantics of FOL: Formalizing the intuition

Vocabulary \mathcal{V} : E.g. $\mathcal{V} : \{a, f, =, R\}$

\mathcal{V} -formula: E.g. $\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$
 $\underline{\text{free}(\varphi) = \{y\}}$

Semantics of FOL: Formalizing the intuition

Vocabulary \mathcal{V} : E.g. $\mathcal{V} : \{a, f, =, R\}$

\mathcal{V} -formula: E.g. $\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$
 $\text{free}(\varphi) = \{y\}$

Truth of φ depends on

Semantics of FOL: Formalizing the intuition

Vocabulary \mathcal{V} : E.g. $\mathcal{V} : \{a, f, =, R\}$

\mathcal{V} -formula: E.g. $\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$
 $\text{free}(\varphi) = \{y\}$

Truth of φ depends on

① Universe U from which variables take values, e.g. \mathbb{N}

Semantics of FOL: Formalizing the intuition

Vocabulary \mathcal{V} : E.g. $\mathcal{V} : \{a, f, \dot{=}, \overline{R}\}$

\mathcal{V} -formula: E.g. $\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$
 $\text{free}(\varphi) = \{y\}$

Truth of φ depends on

- ① Universe U from which variables take values, e.g. \mathbb{N}
- ② Interpretation of vocabulary \mathcal{V} on U

Semantics of FOL: Formalizing the intuition

Vocabulary \mathcal{V} : E.g. $\mathcal{V} : \{a, f, =, R\}$

\mathcal{V} -formula: E.g. $\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$
 $\text{free}(\varphi) = \{y\}$

Truth of φ depends on

- ① Universe U from which variables take values, e.g. \mathbb{N}
- ② Interpretation of vocabulary \mathcal{V} on U
 - Map each constant symbol to an element of U , e.g. $a \mapsto 0$

Semantics of FOL: Formalizing the intuition

Vocabulary \mathcal{V} : E.g. $\mathcal{V} : \{a, f, =, R\}$

\mathcal{V} -formula: E.g. $\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$
 $\text{free}(\varphi) = \{y\}$

Truth of φ depends on

① Universe U from which variables take values, e.g. \mathbb{N}

② Interpretation of vocabulary \mathcal{V} on U

- Map each constant symbol to an element of U , e.g. $a \mapsto 0$
- Map each n -ary function symbol to a function from U^n to U ,
e.g. $f(u, v) = u + v$

Semantics of FOL: Formalizing the intuition

Vocabulary \mathcal{V} : E.g. $\mathcal{V} : \{a, f, =, R\}$

\mathcal{V} -formula: E.g. $\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$
 $\text{free}(\varphi) = \{y\}$

Truth of φ depends on

① Universe U from which variables take values, e.g. \mathbb{N}

② Interpretation of vocabulary \mathcal{V} on U

- Map each constant symbol to an element of U , e.g. $a \mapsto 0$
- Map each n -ary function symbol to a function from U^n to U ,
e.g. $f(u, v) = u + v$
- Map each m -ary predicate symbol to a subset of U^m
e.g. Interp. for $=$: $\{(c, c) \mid c \in \mathbb{N}\}$ – fixed interpretation
Interp. for R : $\{(c, d) \mid c, d \in U, c < d\}$

Semantics of FOL: Formalizing the intuition

Vocabulary \mathcal{V} : E.g. $\mathcal{V} : \{a, f, =, R\}$

\mathcal{V} -formula: E.g. $\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$
 $\text{free}(\varphi) = \{y\}$

Truth of φ depends on

① Universe U from which variables take values, e.g. \mathbb{N}

② Interpretation of vocabulary \mathcal{V} on U

- Map each constant symbol to an element of U , e.g. $a \mapsto 0$
- Map each n -ary function symbol to a function from U^n to U ,
e.g. $f(u, v) = u + v$
- Map each m -ary predicate symbol to a subset of U^m
e.g. Interp. for $=$: $\{(c, c) \mid c \in \mathbb{N}\}$ – fixed interpretation
Interp. for R : $\{(c, d) \mid c, d \in U, c < d\}$

1 and 2 define a **\mathcal{V} -structure** $M = (U^M, (a^M, f^M, R^M))$

Semantics of FOL: Formalizing the intuition

Vocabulary \mathcal{V} : E.g. $\mathcal{V} : \{a, f, =, R\}$

\mathcal{V} -formula: E.g. $\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$
 $\text{free}(\varphi) = \{y\}$

Truth of φ depends on

① Universe U from which variables take values, e.g. \mathbb{N}

② Interpretation of vocabulary \mathcal{V} on U

- Map each constant symbol to an element of U , e.g. $a \mapsto 0$
- Map each n -ary function symbol to a function from U^n to U ,
e.g. $f(u, v) = u + v$
- Map each m -ary predicate symbol to a subset of U^m
e.g. Interp. for $=$: $\{(c, c) \mid c \in \mathbb{N}\}$ – fixed interpretation
Interp. for R : $\{(c, d) \mid c, d \in U, c < d\}$

1 and 2 define a \mathcal{V} -structure $M = (U^M, (a^M, f^M, R^M))$

③ Binding (aka environment) $\alpha : \text{free}(\varphi) \rightarrow U$

e.g. $\alpha(y) = 2$

$$M \rightarrow (\mathbb{N}, (a, f, R))$$

Semantics of FOL: Formalizing the intuition

Given structure M and binding α , does φ evaluate to **true**?

Notationally, does $\underline{M}, \underline{\alpha} \models \underline{\varphi}$?

Semantics of FOL: Formalizing the intuition

Given structure M and binding α , does φ evaluate to **true**?

Notationally, does $\mathbf{M}, \alpha \models \varphi$?

- Extend $\underline{\alpha} : \text{free}(\varphi) \rightarrow U^M$ to $\bar{\alpha} : \text{Terms}(\varphi) \rightarrow U^M$
 - If t is a variable x , $\bar{\alpha}(t) = \alpha(x)$
 - If t is $f(t_1, \dots, t_m)$, $\bar{\alpha}(t) = f^M(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m))$



Semantics of FOL: Formalizing the intuition

Given structure M and binding α , does φ evaluate to **true**?

Notationally, does $\mathbf{M}, \alpha \models \varphi$?

- Extend $\alpha : \text{free}(\varphi) \rightarrow U^M$ to $\bar{\alpha} : \text{Terms}(\varphi) \rightarrow U^M$
 - If t is a variable x , $\bar{\alpha}(t) = \alpha(x)$
 - If t is $f(t_1, \dots, t_m)$, $\bar{\alpha}(t) = f^M(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m))$
 - In prev. example, $\bar{\alpha}(f(y, a)) = f^M(\alpha(y), a^M) = 2 + 0 = 2$

Semantics of FOL: Formalizing the intuition

Given structure M and binding α , does φ evaluate to **true**?

Notationally, does $M, \alpha \models \varphi$?

- Extend $\alpha : \text{free}(\varphi) \rightarrow U^M$ to $\bar{\alpha} : \text{Terms}(\varphi) \rightarrow U^M$
 - If t is a variable x , $\bar{\alpha}(t) = \alpha(x)$
 - If t is $f(t_1, \dots, t_m)$, $\bar{\alpha}(t) = f^M(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m))$
 - In prev. example, $\bar{\alpha}(f(y, a)) = f^M(\alpha(y), a^M) = 2 + 0 = 2$
- If φ is an atomic formula
 - $M, \alpha \models (t_1 = t_2)$ iff $\bar{\alpha}(t_1)$ and $\bar{\alpha}(t_2)$ coincide
 - $M, \alpha \models P(t_1, \dots, t_m)$ iff $(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m)) \in P^M$
 $\exists \quad \sim \quad \neg$

Semantics of FOL: Formalizing the intuition

Given structure M and binding α , does φ evaluate to **true**?

Notationally, does $\mathbf{M}, \alpha \models \varphi$?

- Extend $\alpha : \text{free}(\varphi) \rightarrow U^M$ to $\bar{\alpha} : \text{Terms}(\varphi) \rightarrow U^M$
 - If t is a variable x , $\bar{\alpha}(t) = \alpha(x)$
 - If t is $f(t_1, \dots, t_m)$, $\bar{\alpha}(t) = f^M(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m))$
 - In prev. example, $\bar{\alpha}(f(y, a)) = f^M(\alpha(y), a^M) = 2 + 0 = 2$
- If φ is an atomic formula
 - $M, \alpha \models (t_1 = t_2)$ iff $\bar{\alpha}(t_1)$ and $\bar{\alpha}(t_2)$ coincide
 - $M, \alpha \models P(t_1, \dots, t_m)$ iff $(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m)) \in P^M$
 - In prev. example, suppose $\alpha'(x) = 1, \alpha'(y) = 2$. Then $M, \alpha' \models R(x, f(y, a))$ as $(\bar{\alpha}'(x), \bar{\alpha}'(f(y, a))) = (1, 2) \in R^M$.

Semantics of FOL: Formalizing the intuition

Given structure M and binding α , does φ evaluate to **true**?

Notationally, does $\mathbf{M}, \alpha \models \varphi$?

- Extend $\alpha : \text{free}(\varphi) \rightarrow U^M$ to $\bar{\alpha} : \text{Terms}(\varphi) \rightarrow U^M$
 - If t is a variable x , $\bar{\alpha}(t) = \alpha(x)$
 - If t is $f(t_1, \dots, t_m)$, $\bar{\alpha}(t) = f^M(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m))$
 - In prev. example, $\bar{\alpha}(f(y, a)) = f^M(\alpha(y), a^M) = 2 + 0 = 2$
- If φ is an atomic formula
 - $M, \alpha \models (t_1 = t_2)$ iff $\bar{\alpha}(t_1)$ and $\bar{\alpha}(t_2)$ coincide
 - $M, \alpha \models P(t_1, \dots, t_m)$ iff $(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m)) \in P^M$
 - In prev. example, suppose $\alpha'(x) = 1, \alpha'(y) = 2$. Then $M, \alpha' \models R(x, f(y, a))$ as $(\bar{\alpha}'(x), \bar{\alpha}'(f(y, a))) = (1, 2) \in R^M$.

• $M, \alpha \models \neg\varphi$; iff $M, \alpha \not\models \varphi_1$

Didn't understand

Semantics of FOL: Formalizing the intuition

Given structure M and binding α , does φ evaluate to **true**?

Notationally, does $\mathbf{M}, \alpha \models \varphi$?

- Extend $\alpha : \text{free}(\varphi) \rightarrow U^M$ to $\bar{\alpha} : \text{Terms}(\varphi) \rightarrow U^M$
 - If t is a variable x , $\bar{\alpha}(t) = \alpha(x)$
 - If t is $f(t_1, \dots, t_m)$, $\bar{\alpha}(t) = f^M(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m))$
 - In prev. example, $\bar{\alpha}(f(y, a)) = f^M(\alpha(y), a^M) = 2 + 0 = 2$
- If φ is an atomic formula
 - $M, \alpha \models (t_1 = t_2)$ iff $\bar{\alpha}(t_1)$ and $\bar{\alpha}(t_2)$ coincide
 - $M, \alpha \models P(t_1, \dots, t_m)$ iff $(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m)) \in P^M$
 - In prev. example, suppose $\alpha'(x) = 1, \alpha'(y) = 2$. Then $M, \alpha' \models R(x, f(y, a))$ as $(\bar{\alpha}'(x), \bar{\alpha}'(f(y, a))) = (1, 2) \in R^M$.
- $M, \alpha \models \neg\varphi$; iff $M, \alpha \not\models \varphi$
- $\underline{M, \alpha \models \varphi_1 \wedge \varphi_2}$ iff $\underline{M, \alpha \models \varphi_1}$ and $\underline{M, \alpha \models \varphi_2}$

Semantics of FOL: Formalizing the intuition

Given structure M and binding α , does φ evaluate to **true**?

Notationally, does $M, \alpha \models \varphi$?

- Extend $\alpha : \text{free}(\varphi) \rightarrow U^M$ to $\bar{\alpha} : \text{Terms}(\varphi) \rightarrow U^M$
 - If t is a variable x , $\bar{\alpha}(t) = \alpha(x)$
 - If t is $f(t_1, \dots, t_m)$, $\bar{\alpha}(t) = f^M(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m))$
 - In prev. example, $\bar{\alpha}(f(y, a)) = f^M(\alpha(y), a^M) = 2 + 0 = 2$
- If φ is an atomic formula
 - $M, \alpha \models (t_1 = t_2)$ iff $\bar{\alpha}(t_1)$ and $\bar{\alpha}(t_2)$ coincide
 - $M, \alpha \models P(t_1, \dots, t_m)$ iff $(\bar{\alpha}(t_1), \dots, \bar{\alpha}(t_m)) \in P^M$
 - In prev. example, suppose $\alpha'(x) = 1, \alpha'(y) = 2$. Then $M, \alpha' \models R(x, f(y, a))$ as $(\bar{\alpha}'(x), \bar{\alpha}'(f(y, a))) = (1, 2) \in R^M$.
- $M, \alpha \models \neg\varphi$; iff $M, \alpha \not\models \varphi$
- $M, \alpha \models \varphi_1 \wedge \varphi_2$ iff $M, \alpha \models \varphi_1$ and $M, \alpha \models \varphi_2$
- $M, \alpha \models \exists x \varphi$ iff there is some $c \in U^M$ such that $M, \alpha[x \mapsto c] \models \varphi$, where
 - $\alpha[x \mapsto c](v) = \underline{\alpha(v)}$, if variable v is different from \underline{x}
 - $\underline{\alpha[x \mapsto c](x)} = c$

Semantics of FOL: Illustration

$$\varphi \triangleq \underbrace{\exists x R(x, f(y, a))}_{\not x, y} \rightarrow \underbrace{\exists z (\neg(z = a) \wedge R(z, y))}_{\not z, y}$$

Semantics of FOL: Illustration

$$\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$$

- Vocabulary $\mathcal{V} : \{a, f, =, R\}$

Semantics of FOL: Illustration

$$\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$$

- Vocabulary $\mathcal{V} : \{a, f, =, R\}$
- $M = (\underbrace{\mathbb{N}}_{=}, (\underbrace{0}_{=}, \underbrace{u+v}_{=}, \underbrace{u < v}_{=}))$

Semantics of FOL: Illustration

$$\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$$

- Vocabulary $\mathcal{V} : \{a, f, =, R\}$
- $M = (\overbrace{\mathbb{N}}^{U^M}, (\overbrace{0}^{a^M}, \overbrace{u+v}^{f^M(u,v)}, \overbrace{u < v}^{R^M(u,v)}))$
- $\text{free}(\varphi) = \{y\}$, and $\alpha(y) = 2$

Semantics of FOL: Illustration

$$\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$$

- Vocabulary $\mathcal{V} : \{a, f, =, R\}$

- $M = (\overbrace{\mathbb{N}}^{U^M}, \underbrace{0}_{a^M}, \underbrace{u+v}_{f^M(u,v)}, \underbrace{u < v}_{R^M(u,v)})$
 - $\text{free}(\varphi) = \{y\}$, and $\alpha(y) = 2$
 - $M, \underline{\alpha[z \mapsto 1]} \models (\underbrace{\neg(z = a)}_{1 \neq 0} \wedge R(z, \underbrace{y}_{?}))$
 - Therefore, $\underline{M, \alpha \models \exists z (\neg(z = a) \wedge R(z, y))}$
- 1 < 2 holds
y ↦ 2, z ↦ 1
z ↦ 1*

Semantics of FOL: Illustration

$$\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$$

- Vocabulary $\mathcal{V} : \{a, f, =, R\}$
- $M = (\overbrace{\mathbb{N}}^{U^M}, (\overbrace{0}^{a^M}, \overbrace{u+v}^{f^M(u,v)}, \overbrace{u < v}^{R^M(u,v)}))$
- $\text{free}(\varphi) = \{y\}$, and $\alpha(y) = 2$
- $M, \alpha[z \mapsto 1] \models (\neg(z = a) \wedge R(z, y))$
- Therefore, $M, \alpha \models \exists z (\neg(z = a) \wedge R(z, y))$
- Similarly, $M, \alpha[x \mapsto 0] \models R(x, f(y, a))$
- Therefore, $M, \alpha \models \exists x R(x, f(y, a))$

Semantics of FOL: Illustration

$$\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$$

- Vocabulary $\mathcal{V} : \{a, f, =, R\}$
- $M = (\overbrace{\mathbb{N}}^{U^M}, (\overbrace{0}^{a^M}, \overbrace{u+v}^{f^M(u,v)}, \overbrace{u < v}^{R^M(u,v)}))$
- $\text{free}(\varphi) = \{y\}$, and $\alpha(y) = 2$
- $M, \alpha[z \mapsto 1] \models (\neg(z = a) \wedge R(z, y))$
- Therefore, $M, \alpha \models \exists z (\neg(z = a) \wedge R(z, y))$
- Similarly, $M, \alpha[x \mapsto 0] \models R(x, f(y, a))$
- Therefore, $M, \alpha \models \exists x R(x, f(y, a))$
- Finally, $M, \alpha \models \varphi$

Semantics of FOL: Illustration

$$\varphi \triangleq \exists x R(x, f(y, a)) \rightarrow \exists z (\neg(z = a) \wedge R(z, y))$$

- Vocabulary $\mathcal{V} : \{a, f, =, R\}$

$$\bullet M = (\underbrace{U^M}_{\mathbb{N}}, \underbrace{a^M}_0, \underbrace{f^M(u, v)}_{u + v}, \underbrace{R^M(u, v)}_{u < v})$$

- $\text{free}(\varphi) = \{y\}$, and $\alpha(y) = 2$

- $M, \alpha[z \mapsto 1] \models (\neg(z = a) \wedge R(z, y))$

- Therefore, $M, \alpha \models \exists z (\neg(z = a) \wedge R(z, y))$

- Similarly, $M, \alpha[x \mapsto 0] \models R(x, f(y, a))$

- Therefore, $M, \alpha \models \exists x R(x, f(y, a))$

- Finally, $M, \alpha \models \varphi$

- Note that if $\alpha'(y) = 1$, $M, \alpha' \not\models \varphi$

won't work

holds