# Section 1

Atlas 12-Week General Fitness Program is a fictional synthetic training guide created for dRAGn retrieval demonstrations. It targets general strength, cardiovascular capacity, mobility, and consistency. It is not medical advice.

## Program design

- Three strength sessions per week.
- Two cardio sessions per week.
- One active recovery session.
- One full rest day.
- Most strength sets finish with approximately 2–3 repetitions in reserve.

The program prioritizes consistency and controlled progression. A participant should not force a planned progression when technique, recovery, or symptoms indicate that the current workload is inappropriate.

## Operational scenario

The weekly structure is intentionally distributed across the week. Monday, Wednesday, and Friday are strength days; Tuesday and Saturday are cardio; Thursday is active recovery; Sunday is the full rest day. The schedule is not a requirement to perform seven hard workouts. Adding another intense session on Thursday can change the recovery balance and make the following strength session less productive.

## Interpretation and operational notes

The weekly schedule should be interpreted as a workload distribution rather than seven mandatory hard sessions. Monday, Wednesday, and Friday are strength days. Tuesday and Saturday are cardio days. Thursday is intentionally recovery-oriented, and Sunday is the full rest day. Treating Thursday as another strength session or adding unscheduled high-intensity work can change the intended recovery balance. This distinction makes the guide useful for testing retrieval of schedule-specific conditions rather than simple exercise-name matching.

Strength progression is conditional. The plan does not instruct the participant to add weight every week regardless of performance. Load should increase only after the top of the prescribed repetition range is completed with good technique for two sessions. If form deteriorates, recovery is poor, or symptoms occur, maintaining or reducing the workload is more appropriate. This is a useful example of why an AI answer should retrieve both the progression rule and the condition attached to it.

Cardio progression has its own conditions. Tuesday uses steady moderate aerobic work for 25–40 minutes. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good. The plan therefore contains a time-based gate, a session structure, and a progression ceiling that should not be conflated.

The guide treats recovery as part of training rather than as an optional afterthought. Sleep, hydration, food intake, and manageable training volume affect readiness. Persistent decline in performance, unusually poor sleep, or prolonged soreness are signals to reduce training volume rather than force progression. The safety section also distinguishes ordinary fatigue from concerning symptoms such as sharp pain, fainting, unusual shortness of breath, or sudden dizziness. These distinctions are important when generating grounded answers from the document.

# 1. Weekly Structure

| Day | Session | Duration | Focus |
|---|---|---|---|
| Monday | Strength A | 45–60 min | Squat, push, core |
| Tuesday | Cardio 1 | 25–40 min | Steady aerobic |
| Wednesday | Strength B | 45–60 min | Hinge, pull, core |
| Thursday | Recovery | 20–30 min | Mobility / walking |
| Friday | Strength C | 45–60 min | Full body |
| Saturday | Cardio 2 | 25–45 min | Intervals / steady |
| Sunday | Rest | — | Recovery |

## Weekly interpretation

Thursday is intentionally low intensity and should not be treated as a fourth strength workout. Sunday is the full rest day. Saturday intervals are not recommended immediately for an inconsistent beginner.

## Operational scenario

A participant who can complete all prescribed repetitions with good technique should progress gradually, but the plan does not require an increase every week. Warm up for 5–8 minutes, use controlled range of motion, and rest 90–150 seconds between demanding compound sets. Accessory work generally uses 45–90 seconds of rest. If recovery or technique deteriorates, repeating the previous workload is preferable to forcing progression.

## Interpretation and operational notes

Strength progression is conditional. The plan does not instruct the participant to add weight every week regardless of performance. Load should increase only after the top of the prescribed repetition range is completed with good technique for two sessions. If form deteriorates, recovery is poor, or symptoms occur, maintaining or reducing the workload is more appropriate. This is a useful example of why an AI answer should retrieve both the progression rule and the condition attached to it.

Cardio progression has its own conditions. Tuesday uses steady moderate aerobic work for 25–40 minutes. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good. The plan therefore contains a time-based gate, a session structure, and a progression ceiling that should not be conflated.

The guide treats recovery as part of training rather than as an optional afterthought. Sleep, hydration, food intake, and manageable training volume affect readiness. Persistent decline in performance, unusually poor sleep, or prolonged soreness are signals to reduce training volume rather than force progression. The safety section also distinguishes ordinary fatigue from concerning symptoms such as sharp pain, fainting, unusual shortness of breath, or sudden dizziness. These distinctions are important when generating grounded answers from the document.

The weekly schedule should be interpreted as a workload distribution rather than seven mandatory hard sessions. Monday, Wednesday, and Friday are strength days. Tuesday and Saturday are cardio days. Thursday is intentionally recovery-oriented, and Sunday is the full rest day. Treating Thursday as another strength session or adding unscheduled high-intensity work can change the intended recovery balance. This distinction makes the guide useful for testing retrieval of schedule-specific conditions rather than simple exercise-name matching.

# 2. Strength Training Rules

- Begin with 5–8 minutes of general warm-up followed by movement-specific preparation.
- Use controlled movement and consistent range of motion.
- Rest 90–150 seconds between demanding compound sets and 45–90 seconds for accessory movements.
- Increase load by the smallest practical increment after completing the prescribed range with good technique for two sessions.
- Do not increase load when technique deteriorates or recovery is poor.

## Readiness

Readiness can vary from session to session. The plan is a framework rather than a command to increase load every week. A participant can repeat a workload when recovery is inadequate.

## Operational scenario

Strength A emphasizes squat, push, row, hinge, and core patterns. A participant who cannot perform a full push-up can use an incline variation without changing the purpose of the session. The Romanian deadlift should remain a controlled hip hinge rather than a maximal lift. The dead bug is performed with controlled trunk position. The goal is repeatable technique across sessions, not simply completing the largest possible number of repetitions.

## Interpretation and operational notes

Cardio progression has its own conditions. Tuesday uses steady moderate aerobic work for 25–40 minutes. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good. The plan therefore contains a time-based gate, a session structure, and a progression ceiling that should not be conflated.

The guide treats recovery as part of training rather than as an optional afterthought. Sleep, hydration, food intake, and manageable training volume affect readiness. Persistent decline in performance, unusually poor sleep, or prolonged soreness are signals to reduce training volume rather than force progression. The safety section also distinguishes ordinary fatigue from concerning symptoms such as sharp pain, fainting, unusual shortness of breath, or sudden dizziness. These distinctions are important when generating grounded answers from the document.

The weekly schedule should be interpreted as a workload distribution rather than seven mandatory hard sessions. Monday, Wednesday, and Friday are strength days. Tuesday and Saturday are cardio days. Thursday is intentionally recovery-oriented, and Sunday is the full rest day. Treating Thursday as another strength session or adding unscheduled high-intensity work can change the intended recovery balance. This distinction makes the guide useful for testing retrieval of schedule-specific conditions rather than simple exercise-name matching.

Strength progression is conditional. The plan does not instruct the participant to add weight every week regardless of performance. Load should increase only after the top of the prescribed repetition range is completed with good technique for two sessions. If form deteriorates, recovery is poor, or symptoms occur, maintaining or reducing the workload is more appropriate. This is a useful example of why an AI answer should retrieve both the progression rule and the condition attached to it.

# 3. Strength A

| Exercise | Sets | Repetitions |
|---|---|---|
| Goblet squat | 3 | 8–12 |
| Push-up / incline push-up | 3 | 8–15 |
| Dumbbell row | 3 | 8–12 / side |
| Romanian deadlift | 2 | 8–12 |
| Dead bug | 3 | 8–12 / side |

## Technique notes

- Keep the squat controlled and maintain stable foot pressure.
- Use an incline push-up when full push-ups compromise technique.
- Avoid momentum during rows.
- Keep the Romanian deadlift as a controlled hip hinge rather than a maximal lift.

## Operational scenario

Strength B emphasizes the hinge, vertical pulling, unilateral leg work, shoulder pressing, and lateral core stability. A supported split squat is an appropriate substitution when balance is the limiting factor. The shoulder press should stop when the participant cannot maintain controlled shoulder position. The session complements Strength A by changing movement emphasis while keeping weekly volume manageable.

## Interpretation and operational notes

The guide treats recovery as part of training rather than as an optional afterthought. Sleep, hydration, food intake, and manageable training volume affect readiness. Persistent decline in performance, unusually poor sleep, or prolonged soreness are signals to reduce training volume rather than force progression. The safety section also distinguishes ordinary fatigue from concerning symptoms such as sharp pain, fainting, unusual shortness of breath, or sudden dizziness. These distinctions are important when generating grounded answers from the document.

The weekly schedule should be interpreted as a workload distribution rather than seven mandatory hard sessions. Monday, Wednesday, and Friday are strength days. Tuesday and Saturday are cardio days. Thursday is intentionally recovery-oriented, and Sunday is the full rest day. Treating Thursday as another strength session or adding unscheduled high-intensity work can change the intended recovery balance. This distinction makes the guide useful for testing retrieval of schedule-specific conditions rather than simple exercise-name matching.

Strength progression is conditional. The plan does not instruct the participant to add weight every week regardless of performance. Load should increase only after the top of the prescribed repetition range is completed with good technique for two sessions. If form deteriorates, recovery is poor, or symptoms occur, maintaining or reducing the workload is more appropriate. This is a useful example of why an AI answer should retrieve both the progression rule and the condition attached to it.

Cardio progression has its own conditions. Tuesday uses steady moderate aerobic work for 25–40 minutes. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good. The plan therefore contains a time-based gate, a session structure, and a progression ceiling that should not be conflated.

# 4. Strength B

| Exercise | Sets | Repetitions |
|---|---|---|
| Romanian deadlift | 3 | 8–12 |
| Lat pulldown / band pulldown | 3 | 8–12 |
| Split squat | 3 | 8–12 / side |
| Dumbbell shoulder press | 2 | 8–12 |
| Side plank | 3 | 20–40 sec / side |

## Technique notes

- Maintain a neutral spine during the hinge.
- Use a supported split squat if balance is limiting.
- Stop a set if shoulder position cannot be controlled.

The purpose of Strength B is to expose the participant to a different movement emphasis from Strength A while preserving manageable weekly volume.

## Operational scenario

Strength C provides a full-body session using a squat or leg-press variation, horizontal press, row, hip thrust, and loaded carry. When the top of the repetition range is achieved with good technique for two sessions, the load can increase slightly and repetitions can return toward the lower end. A push-up can substitute for dumbbell bench press, and a loaded backpack carry can substitute for a farmer carry when equipment is limited.

## Interpretation and operational notes

The weekly schedule should be interpreted as a workload distribution rather than seven mandatory hard sessions. Monday, Wednesday, and Friday are strength days. Tuesday and Saturday are cardio days. Thursday is intentionally recovery-oriented, and Sunday is the full rest day. Treating Thursday as another strength session or adding unscheduled high-intensity work can change the intended recovery balance. This distinction makes the guide useful for testing retrieval of schedule-specific conditions rather than simple exercise-name matching.

Strength progression is conditional. The plan does not instruct the participant to add weight every week regardless of performance. Load should increase only after the top of the prescribed repetition range is completed with good technique for two sessions. If form deteriorates, recovery is poor, or symptoms occur, maintaining or reducing the workload is more appropriate. This is a useful example of why an AI answer should retrieve both the progression rule and the condition attached to it.

Cardio progression has its own conditions. Tuesday uses steady moderate aerobic work for 25–40 minutes. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good. The plan therefore contains a time-based gate, a session structure, and a progression ceiling that should not be conflated.

The guide treats recovery as part of training rather than as an optional afterthought. Sleep, hydration, food intake, and manageable training volume affect readiness. Persistent decline in performance, unusually poor sleep, or prolonged soreness are signals to reduce training volume rather than force progression. The safety section also distinguishes ordinary fatigue from concerning symptoms such as sharp pain, fainting, unusual shortness of breath, or sudden dizziness. These distinctions are important when generating grounded answers from the document.

# 5. Strength C

| Exercise | Sets | Repetitions |
|---|---|---|
| Leg press / squat variation | 3 | 8–12 |
| Dumbbell bench press | 3 | 8–12 |
| Seated row | 3 | 8–12 |
| Hip thrust | 2 | 10–15 |
| Farmer carry | 3 | 30–45 sec |

## Progression

When the top of the repetition range is achieved with good technique for two sessions, increase load slightly and return toward the lower end of the range. If technique deteriorates, keep the previous load rather than forcing progression.

## Substitution

- Leg press may substitute for a squat variation; a push-up may substitute for dumbbell bench press; a loaded backpack carry may substitute for a farmer carry.

## Operational scenario

Steady cardio should feel sustainable enough that the participant can speak in short sentences without gasping. Tuesday uses 25–40 minutes of moderate steady work. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good.

## Interpretation and operational notes

Strength progression is conditional. The plan does not instruct the participant to add weight every week regardless of performance. Load should increase only after the top of the prescribed repetition range is completed with good technique for two sessions. If form deteriorates, recovery is poor, or symptoms occur, maintaining or reducing the workload is more appropriate. This is a useful example of why an AI answer should retrieve both the progression rule and the condition attached to it.

Cardio progression has its own conditions. Tuesday uses steady moderate aerobic work for 25–40 minutes. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good. The plan therefore contains a time-based gate, a session structure, and a progression ceiling that should not be conflated.

The guide treats recovery as part of training rather than as an optional afterthought. Sleep, hydration, food intake, and manageable training volume affect readiness. Persistent decline in performance, unusually poor sleep, or prolonged soreness are signals to reduce training volume rather than force progression. The safety section also distinguishes ordinary fatigue from concerning symptoms such as sharp pain, fainting, unusual shortness of breath, or sudden dizziness. These distinctions are important when generating grounded answers from the document.

The weekly schedule should be interpreted as a workload distribution rather than seven mandatory hard sessions. Monday, Wednesday, and Friday are strength days. Tuesday and Saturday are cardio days. Thursday is intentionally recovery-oriented, and Sunday is the full rest day. Treating Thursday as another strength session or adding unscheduled high-intensity work can change the intended recovery balance. This distinction makes the guide useful for testing retrieval of schedule-specific conditions rather than simple exercise-name matching.

# 6. Cardio

## Steady aerobic work

- Zone 2 sessions should feel sustainable enough to speak in short sentences without gasping.
- Tuesday cardio lasts 25–40 minutes at a steady moderate effort.

## Intervals

- After at least four weeks of consistent training, Saturday may use 6 rounds of 1 minute faster effort followed by 2 minutes easy.
- Beginners should establish consistent steady cardio before adding intervals.

## Progression

- Increase weekly cardio duration by no more than approximately 10% when recovery is good.

Cardio progression should consider total training load. A participant adding intervals should not simultaneously make large increases to strength volume and weekly duration.

## Operational scenario

Thursday recovery should leave the participant feeling better rather than exhausted. Useful mobility work includes ankle mobility, hip-flexor stretching, thoracic rotation, hamstring mobility, and shoulder range-of-motion drills. Sleep, hydration, and adequate food intake affect readiness. Persistent performance decline, unusually poor sleep, or prolonged soreness is a reason to reduce volume rather than add more training.

## Interpretation and operational notes

Cardio progression has its own conditions. Tuesday uses steady moderate aerobic work for 25–40 minutes. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good. The plan therefore contains a time-based gate, a session structure, and a progression ceiling that should not be conflated.

The guide treats recovery as part of training rather than as an optional afterthought. Sleep, hydration, food intake, and manageable training volume affect readiness. Persistent decline in performance, unusually poor sleep, or prolonged soreness are signals to reduce training volume rather than force progression. The safety section also distinguishes ordinary fatigue from concerning symptoms such as sharp pain, fainting, unusual shortness of breath, or sudden dizziness. These distinctions are important when generating grounded answers from the document.

The weekly schedule should be interpreted as a workload distribution rather than seven mandatory hard sessions. Monday, Wednesday, and Friday are strength days. Tuesday and Saturday are cardio days. Thursday is intentionally recovery-oriented, and Sunday is the full rest day. Treating Thursday as another strength session or adding unscheduled high-intensity work can change the intended recovery balance. This distinction makes the guide useful for testing retrieval of schedule-specific conditions rather than simple exercise-name matching.

Strength progression is conditional. The plan does not instruct the participant to add weight every week regardless of performance. Load should increase only after the top of the prescribed repetition range is completed with good technique for two sessions. If form deteriorates, recovery is poor, or symptoms occur, maintaining or reducing the workload is more appropriate. This is a useful example of why an AI answer should retrieve both the progression rule and the condition attached to it.

# 7. Mobility and Recovery

- Thursday should leave the participant feeling better rather than exhausted.
- Useful movements include ankle mobility, hip-flexor stretch, thoracic rotation, hamstring mobility, and shoulder range-of-motion drills.
- Sleep, hydration, and adequate food intake influence recovery and training performance.
- Persistent decline in performance, unusually poor sleep, or prolonged soreness is a reason to reduce training volume.

## Recovery checklist

- Keep a consistent sleep schedule.
- Hydrate according to climate, activity, body size, and sweat rate.
- Avoid adding unplanned high-intensity sessions when recovery is poor.

## Operational scenario

The 12-week structure starts with technique, moves into gradual overload, consolidates progress, and finishes with a deload. During weeks 1–4 the emphasis is consistency and 2–3 repetitions in reserve. Weeks 5–8 emphasize gradual increases, weeks 9–11 consolidate, and week 12 reduces strength volume by approximately 30–40%. Progression should follow performance rather than the calendar alone.

## Interpretation and operational notes

The guide treats recovery as part of training rather than as an optional afterthought. Sleep, hydration, food intake, and manageable training volume affect readiness. Persistent decline in performance, unusually poor sleep, or prolonged soreness are signals to reduce training volume rather than force progression. The safety section also distinguishes ordinary fatigue from concerning symptoms such as sharp pain, fainting, unusual shortness of breath, or sudden dizziness. These distinctions are important when generating grounded answers from the document.

The weekly schedule should be interpreted as a workload distribution rather than seven mandatory hard sessions. Monday, Wednesday, and Friday are strength days. Tuesday and Saturday are cardio days. Thursday is intentionally recovery-oriented, and Sunday is the full rest day. Treating Thursday as another strength session or adding unscheduled high-intensity work can change the intended recovery balance. This distinction makes the guide useful for testing retrieval of schedule-specific conditions rather than simple exercise-name matching.

Strength progression is conditional. The plan does not instruct the participant to add weight every week regardless of performance. Load should increase only after the top of the prescribed repetition range is completed with good technique for two sessions. If form deteriorates, recovery is poor, or symptoms occur, maintaining or reducing the workload is more appropriate. This is a useful example of why an AI answer should retrieve both the progression rule and the condition attached to it.

Cardio progression has its own conditions. Tuesday uses steady moderate aerobic work for 25–40 minutes. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good. The plan therefore contains a time-based gate, a session structure, and a progression ceiling that should not be conflated.

# 8. 12-Week Progression

| Weeks | Focus | Approach |
|---|---|---|
| 1–4 | Technique | Consistency; 2–3 reps in reserve |
| 5–8 | Overload | Gradual reps or load increase |
| 9–11 | Consolidation | Progress only with adequate recovery |
| 12 | Deload | Reduce strength volume 30–40% |

## Example

If an exercise target is 8–12 repetitions and a participant completes 12, 12, and 12 with good form for two sessions, increase the load slightly and return toward 8–10 repetitions. If the participant instead completes 12, 10, and 8 with deteriorating technique, the load should not be increased.

## Operational scenario

Nutrition should support training rather than undermine it. Protein is best distributed across meals, while fruit or vegetables, minimally processed carbohydrate sources, and adequate overall food intake support recovery. Aggressive calorie restriction can reduce training performance. Substitutions such as bodyweight squat for goblet squat or resistance-band row for dumbbell row preserve movement patterns when equipment is unavailable.

## Interpretation and operational notes

The weekly schedule should be interpreted as a workload distribution rather than seven mandatory hard sessions. Monday, Wednesday, and Friday are strength days. Tuesday and Saturday are cardio days. Thursday is intentionally recovery-oriented, and Sunday is the full rest day. Treating Thursday as another strength session or adding unscheduled high-intensity work can change the intended recovery balance. This distinction makes the guide useful for testing retrieval of schedule-specific conditions rather than simple exercise-name matching.

Strength progression is conditional. The plan does not instruct the participant to add weight every week regardless of performance. Load should increase only after the top of the prescribed repetition range is completed with good technique for two sessions. If form deteriorates, recovery is poor, or symptoms occur, maintaining or reducing the workload is more appropriate. This is a useful example of why an AI answer should retrieve both the progression rule and the condition attached to it.

Cardio progression has its own conditions. Tuesday uses steady moderate aerobic work for 25–40 minutes. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good. The plan therefore contains a time-based gate, a session structure, and a progression ceiling that should not be conflated.

The guide treats recovery as part of training rather than as an optional afterthought. Sleep, hydration, food intake, and manageable training volume affect readiness. Persistent decline in performance, unusually poor sleep, or prolonged soreness are signals to reduce training volume rather than force progression. The safety section also distinguishes ordinary fatigue from concerning symptoms such as sharp pain, fainting, unusual shortness of breath, or sudden dizziness. These distinctions are important when generating grounded answers from the document.

# 9. Nutrition, Substitutions and Safety

## Nutrition

- Distribute protein across meals rather than concentrating it in one meal.
- Choose minimally processed foods, vegetables or fruit, whole-food carbohydrate sources, and protein-rich foods.
- Aggressive calorie restriction can reduce training performance and recovery.

## Substitutions

- Goblet squat → bodyweight squat or leg press.
- Dumbbell row → resistance-band row.
- Lat pulldown → resistance-band pulldown.
- Dumbbell bench press → push-up or incline push-up.

## Safety

- Stop for sharp pain, sudden dizziness, fainting, unusual shortness of breath, or another concerning symptom.
- People with medical conditions, recent injuries, pregnancy, or significant limitations should obtain appropriate professional guidance before starting.

## Operational scenario

The program is a general fitness guide, not medical advice. Sharp pain, sudden dizziness, fainting, unusual shortness of breath, or another concerning symptom is a reason to stop and seek appropriate professional guidance. People with medical conditions, recent injuries, pregnancy, or significant limitations should obtain suitable advice before starting. Consistency, controlled progression, and recovery are more important than forcing a planned workload.

## Interpretation and operational notes

Strength progression is conditional. The plan does not instruct the participant to add weight every week regardless of performance. Load should increase only after the top of the prescribed repetition range is completed with good technique for two sessions. If form deteriorates, recovery is poor, or symptoms occur, maintaining or reducing the workload is more appropriate. This is a useful example of why an AI answer should retrieve both the progression rule and the condition attached to it.

Cardio progression has its own conditions. Tuesday uses steady moderate aerobic work for 25–40 minutes. Saturday intervals are introduced only after at least four weeks of consistent training, using six rounds of one minute faster effort followed by two minutes easy. Weekly cardio duration should increase by no more than approximately 10% when recovery is good. The plan therefore contains a time-based gate, a session structure, and a progression ceiling that should not be conflated.

The guide treats recovery as part of training rather than as an optional afterthought. Sleep, hydration, food intake, and manageable training volume affect readiness. Persistent decline in performance, unusually poor sleep, or prolonged soreness are signals to reduce training volume rather than force progression. The safety section also distinguishes ordinary fatigue from concerning symptoms such as sharp pain, fainting, unusual shortness of breath, or sudden dizziness. These distinctions are important when generating grounded answers from the document.

The weekly schedule should be interpreted as a workload distribution rather than seven mandatory hard sessions. Monday, Wednesday, and Friday are strength days. Tuesday and Saturday are cardio days. Thursday is intentionally recovery-oriented, and Sunday is the full rest day. Treating Thursday as another strength session or adding unscheduled high-intensity work can change the intended recovery balance. This distinction makes the guide useful for testing retrieval of schedule-specific conditions rather than simple exercise-name matching.