

CareTrust Medical Centre

Dr. Sunitha Rao — General Physician & Cardiologist

MBBS, MD (General Medicine) | Reg: TN-MC-29341 | Care Hospital, Chennai

Patient Name	Rahul Kumar	Date	28 May 2026
Age / Gender	58 Yrs / Male	Room No	402
BP at Visit	172/102 mmHg	Weight	88 kg
Diagnosis	Hypertension Stage 2 — persistent elevated BP above 160 systolic for 3 days		

PRESCRIPTION

#	Medicine	Dose	Frequency	Duration	Instructions
1	Amlodipine	10 mg	Once daily morning	30 days	After breakfast with water
2	Telmisartan	80 mg	Once daily morning	30 days	Monitor BP weekly. Target below 130/80
3	Atorvastatin	40 mg	Once daily night	30 days	For high cholesterol — take at night
4	Aspirin (EC)	75 mg	Once daily morning	30 days	After food — do NOT crush or chew
5	Metoprolol	25 mg	Twice daily	30 days	Morning and evening after food

IMPORTANT INSTRUCTIONS

- Measure BP every morning before breakfast and record in the care log.
- Target BP: below 130/80 mmHg. Escalate immediately if BP exceeds 180 systolic.
- Amlodipine 10mg: take once daily in the morning after breakfast.
- Telmisartan 80mg: take once daily in the morning. Do NOT stop without doctor advice.
- Metoprolol 25mg: do NOT skip — sudden stopping can cause rebound hypertension.
- Avoid salty food, pickles, papad, processed food — all raise BP.
- Adequate rest — minimum 7 hours sleep. Stress management important.
- Atorvastatin is for cholesterol — take at night. Avoid grapefruit juice.
- Aspirin EC 75mg is a blood thinner — do NOT crush, chew, or break the tablet.
- Notify doctor immediately if patient reports chest pain, vision blurring, or severe headache.

FOLLOW-UP

Next review: 15 June 2026 | Lab: Lipid profile, RFT, Electrolytes — due 10 June 2026

Emergency contact: 044-6788-1234

Dr. Sunitha Rao | Signature: _____ | CareTrust Medical Centre, Chennai