

CareTrust Cardiology Department

Dr. Ramesh Nair — Interventional Cardiologist

MBBS, DM (Cardiology) | Reg: TN-MC-38821 | Care Hospital, Chennai

Patient Name	James Carter	Date	27 May 2026
Age / Gender	71 Yrs / Male	Room No	404
Diagnosis	Cardiac Monitoring — Oxygen saturation below 95% for 2 consecutive days		
Current SpO2	92–93%	Target SpO2	Above 95%

CARDIAC INVESTIGATIONS

Investigation	Result	Normal	Status
ECG	Sinus rhythm	Normal sinus	Normal
Echocardiogram (EF)	48%	55–70%	Mildly Reduced
Chest X-Ray	Mild cardiomegaly	Normal	Abnormal
SpO2 (2-day average)	92.5%	Above 95%	LOW — Alert Active
Heart Rate	88 bpm	60–100 bpm	Normal
BNP (Heart failure)	280 pg/mL	Below 100	Elevated
Troponin I	0.02 ng/mL	Below 0.04	Normal
Serum Potassium	3.8 mEq/L	3.5–5.0	Normal

PRESCRIPTION

#	Medicine	Dose	Frequency	Duration	Instructions
1	Bisoprolol	5 mg	Once daily morning	30 days	After breakfast — do NOT stop suddenly
2	Furosemide	40 mg	Once daily morning	30 days	Diuretic — monitor urine output and weight
3	Ramipril	5 mg	Once daily morning	30 days	ACE inhibitor — monitor BP and kidney function
4	Spironolactone	25 mg	Once daily	30 days	Monitor potassium — avoid potassium supplements
5	Aspirin (EC)	75 mg	Once daily morning	30 days	After food — do NOT crush
6	Atorvastatin	40 mg	Once daily night	30 days	Cholesterol management

IMPORTANT INSTRUCTIONS

- Record oxygen saturation (SpO2) every morning — alert threshold: below 92%. Call doctor immediately.
- Target SpO2: above 95%. If SpO2 remains below 93% for 2+ days — escalate to respiratory support.
- Furosemide is a diuretic — expect increased urine output. Monitor daily weight — report gain above 2kg/day.
- Bisoprolol 5mg: never stop suddenly — can cause rebound heart rate spike. Taper only under medical supervision.
- Ramipril: monitor BP after starting. First dose may cause dizziness — patient must not stand up suddenly.
- Spironolactone: avoid potassium-rich foods (banana, orange juice) — can raise potassium to dangerous levels.
- Salt restriction: maximum 2g sodium daily. No added salt. Avoid processed food.
- Elevate head of bed by 30 degrees — reduces breathing difficulty at night.
- Daily weight monitoring: same time each morning before breakfast. Record in care log.
- Signs of worsening heart failure: increasing breathlessness, ankle swelling, sudden weight gain — alert immediately.

ALERT — CURRENT STATUS

SpO2 has been below 95% for 2 consecutive days (93% on May 26, 92% on May 27). An AI alert has been triggered in the CareTrust system. Nurse must record SpO2 daily. If oxygen falls below 92% again — supplemental oxygen and immediate physician review required.

FOLLOW-UP

Next cardiology review: 10 June 2026 | Repeat Echo, BNP, Renal function — due 7 June 2026.

Dr. Ramesh Nair | Signature: _____ | CareTrust Cardiology, Chennai