

# MSMS 408 : Practical 01

Ananda Biswas

Exam Roll No. : 24419STC053

February 9, 2026

## ➔ Question

Use Monte Carlo Integration to approximate the following integrals

$$(i) \int_0^1 e^{e^x} dx,$$

$$(ii) \int_{-2}^2 e^{x+x^2} dx$$

and check how the estimates converge to the true value as the sample size increases.

## ➔ Theory

Suppose we have to compute

$$I = \int_a^b f(x) dx. \quad (1)$$

We rewrite

$$I = (b - a)E[f(X)] \quad (2)$$

where  $X \sim U(a, b)$ .

To approximate  $I$ , we generate random numbers  $X_1, X_2, \dots, X_n$  from  $U(a, b)$  and calculate

$$I \approx (b - a) \cdot \frac{1}{n} \sum_{i=1}^n f(X_i). \quad (3)$$

By law of large numbers,  $\frac{1}{n} \sum_{i=1}^n f(X_i) \rightarrow E[f(X)]$  as  $n \rightarrow \infty$ .

## ➔ R Program

```
set.seed(22)
```

```
Monte.Carlo.Integration <- function(integrand, lower.limit, upper.limit, sample.size){  
  x <- runif(n = sample.size, min = lower.limit, max = upper.limit)  
  I <- (upper.limit - lower.limit) * mean(integrand(x))  
  return(I)  
}
```

**1**  $\int_0^1 e^{e^x} dx$

```
func_1 <- function(x) exp(exp(x))
```

```
sample_sizes <- c(100, 500, 700, 1000, 3000, 5000)
```

```
results.1 <- c()

for (size in sample_sizes) {
  results.1 <- append(results.1,
                      Monte.Carlo.Integration(integrand = func_1, sample.size = size,
                                              lower.limit = 0, upper.limit = 1))
}
```

```
df1 <- data.frame(sample.size = sample_sizes,
                  approximate.integral.value = results.1)
```

```
stargazer(df1, summary = FALSE, rownames = FALSE, label = "Table 1")
```

Table 1:

sample.size	approximate.integral.value
100	6.006
500	6.015
700	6.322
1,000	6.410
3,000	6.246
5,000	6.316

$$2 \int_{-2}^2 e^{x+x^2} dx$$

```
func_2 <- function(x) exp(x + x^2)
```

```
results.2 <- c()
```

```
for (size in sample_sizes) {
  results.2 <- append(results.2,
    Monte.Carlo.Integration(integrand = func_2, sample.size = size,
      lower.limit = -2, upper.limit = 2))
}
```


```
df2 <- data.frame(sample.size = sample_sizes,
  approximate.integral.value = results.2)
```

```
stargazer(df2, summary = FALSE, rownames = FALSE, label = "Table 2")
```

Table 2:

sample.size	approximate.integral.value
100	107.312
500	103.273
700	93.557
1,000	98.880
3,000	95.832
5,000	95.987

## ➔ Conclusion

 As sample size increases, the approximates gradually converge to the true integral value.