

21-259 Calculus in 3D

Compiled by Salman Hajizada

First Edition

Chapter 0

Disclaimer

This is not a textbook. Do not use this text to learn the concepts. This is a revision guide or notes used to refresh your memory of the theorems. The best way to learn the content is to read the book, work through the guided exercises and do some practice problems.

Good luck,


SH

Chapter 12

Vectors and the Geometry of Space

12.1 Three-Dimensional Coordinate System

Right-hand rule for determining the coordinate axis:



Projection of a point (x, y, z) onto

- $x - y$ axis: $(x, y, 0)$
- $x - z$ axis: $(x, 0, z)$
- $y - z$ axis: $(0, y, z)$

Distance formula in 3 dimensions: Between two points (x_1, y_1, z_1) and (x_2, y_2, z_2) , the distance is given by:

$$\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

Equation of a Sphere with center at (h, k, l) and radius r is:

$$(x - h)^2 + (y - k)^2 + (z - l)^2 = r^2$$

12.2 Vectors

I will skip the basics, like vector addition definition, etc.

Components of a vector (for notation):

$$\mathbf{a} = \langle a_1, a_2, a_3 \rangle$$

Given points $A(x_1, y_1, z_1)$ and $B(x_2, y_2, z_2)$,

$$\overrightarrow{AB} = \mathbf{a} = \langle x_2 - x_1, y_2 - y_1, z_2 - z_1 \rangle$$

Magnitude of a vector $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$ is

$$|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

Unit vectors:

$$\mathbf{i} = \langle 1, 0, 0 \rangle, \quad \mathbf{j} = \langle 0, 1, 0 \rangle, \quad \mathbf{k} = \langle 0, 0, 1 \rangle$$

Algebraic Properties of Vectors in \mathbb{R}^n

$\forall \mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^n$ and scalars c and d :

1. **Commutativity of addition:** $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
2. **Associativity of addition:** $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
3. **Additive identity:** $\mathbf{u} + \mathbf{0} = \mathbf{u}$
4. **Additive inverse:** $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
5. **Distributivity of scalar multiplication over addition (vectors):** $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
6. **Distributivity of scalar multiplication over addition (scalars):** $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
7. **Associativity of scalar multiplication:** $(cd)\mathbf{u} = c(d\mathbf{u})$
8. **Scalar identity:** $1\mathbf{u} = \mathbf{u}$

12.3 The Dot Product

Definition 1. If $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$ and $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$ then the **dot product** of \mathbf{a} and \mathbf{b} is

$$\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$$

Properties of the Dot Product

\forall vectors $\mathbf{a}, \mathbf{b}, \mathbf{c}$ and scalar c :

1. $\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$
2. $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$
3. $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c}) = \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$
4. $(c\mathbf{a}) \cdot \mathbf{b} = c(\mathbf{a} \cdot \mathbf{b}) = \mathbf{a} \cdot (c\mathbf{b})$
5. $\mathbf{0} \cdot \mathbf{a} = 0$

Theorem 1. If θ is the angle between two vectors \mathbf{a}, \mathbf{b} then

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$$

Corollary 1. Two vectors \mathbf{a}, \mathbf{b} are orthogonal iff $\mathbf{a} \cdot \mathbf{b} = 0$

The **direction angles** of a non-zero vector \mathbf{a} are the angles α, β, γ that \mathbf{a} makes with the positive x -, y -, and z -axes, respectively.

The cosines of these angles are called **direction cosines**:

$$\cos \alpha = \frac{\mathbf{a} \cdot \mathbf{i}}{|\mathbf{a}||\mathbf{i}|} = \frac{a_1}{|\mathbf{a}|}, \quad \cos \beta = \frac{a_2}{|\mathbf{a}|}, \quad \cos \gamma = \frac{a_3}{|\mathbf{a}|}$$

Some nice properties of direction cosines:

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$$

$$\frac{\mathbf{a}}{|\mathbf{a}|} = \langle \cos \alpha, \cos \beta, \cos \gamma \rangle$$

Projections:

The **projection of \mathbf{b} onto \mathbf{a}** means we drop a perpendicular from the tip of \mathbf{b} onto the line spanned by \mathbf{a} . The resulting vector is called the vector projection of \mathbf{b} onto \mathbf{a} .

The **scalar projection of \mathbf{b} onto \mathbf{a}** is the length of this projection, given by

$$\text{comp}_{\mathbf{a}} \mathbf{b} = |\mathbf{b}| \cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|},$$

where θ is the angle between \mathbf{a} and \mathbf{b} .

The **vector projection** of \mathbf{b} onto \mathbf{a} is obtained by multiplying the unit vector in the direction of \mathbf{a} by the scalar projection:

$$\text{proj}_{\mathbf{a}} \mathbf{b} = (\text{comp}_{\mathbf{a}} \mathbf{b}) \frac{\mathbf{a}}{|\mathbf{a}|} = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|^2} \mathbf{a}$$

12.4 The Cross Product