

# 21-259 Calculus in 3D

Compiled by Salman Hajizada

First Edition

# Chapter 0

## Disclaimer

This is not a textbook. Do not use this text to learn the concepts. This is a revision guide or notes used to refresh your memory of the theorems. The best way to learn the content is to read the book, work through the guided exercises and do some practice problems.

Good luck,


SH

# Chapter 12

## Vectors and the Geometry of Space

### 12.1 Three-Dimensional Coordinate System

**Right-hand rule** for determining the coordinate axis:



Projection of a point  $(x, y, z)$  onto

- $x - y$  axis:  $(x, y, 0)$
- $x - z$  axis:  $(x, 0, z)$
- $y - z$  axis:  $(0, y, z)$

**Distance formula in 3 dimensions:** Between two points  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$ , the distance is given by:

$$\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

**Equation of a Sphere** with center at  $(h, k, l)$  and radius  $r$  is:

$$(x - h)^2 + (y - k)^2 + (z - l)^2 = r^2$$

### 12.2 Vectors

I will skip the basics, like vector addition definition, etc.

Components of a vector (for notation):

$$\mathbf{a} = \langle a_1, a_2, a_3 \rangle$$

Given points  $A(x_1, y_1, z_1)$  and  $B(x_2, y_2, z_2)$ ,

$$\overrightarrow{AB} = \mathbf{a} = \langle x_2 - x_1, y_2 - y_1, z_2 - z_1 \rangle$$

**Magnitude of a vector**  $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$  is

$$|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

**Unit vectors:**

$$\mathbf{i} = \langle 1, 0, 0 \rangle, \quad \mathbf{j} = \langle 0, 1, 0 \rangle, \quad \mathbf{k} = \langle 0, 0, 1 \rangle$$

#### Algebraic Properties of Vectors in $\mathbb{R}^n$

$\forall \mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^n$  and scalars  $c$  and  $d$ :

1. **Commutativity of addition:**  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
2. **Associativity of addition:**  $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
3. **Additive identity:**  $\mathbf{u} + \mathbf{0} = \mathbf{u}$
4. **Additive inverse:**  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$
5. **Distributivity of scalar multiplication over addition (vectors):**  $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
6. **Distributivity of scalar multiplication over addition (scalars):**  $(c + d)\mathbf{u} = c\mathbf{u} + d\mathbf{u}$
7. **Associativity of scalar multiplication:**  $(cd)\mathbf{u} = c(d\mathbf{u})$
8. **Scalar identity:**  $1\mathbf{u} = \mathbf{u}$

## 12.3 The Dot Product

**Definition 1.** If  $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$  and  $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$  then the **dot product** of  $\mathbf{a}$  and  $\mathbf{b}$  is

$$\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$$

#### Properties of the Dot Product

$\forall$  vectors  $\mathbf{a}, \mathbf{b}, \mathbf{c}$  and scalar  $c$ :

1.  $\mathbf{a} \cdot \mathbf{a} = |\mathbf{a}|^2$
2.  $\mathbf{a} \cdot \mathbf{b} = \mathbf{b} \cdot \mathbf{a}$
3.  $\mathbf{a} \cdot (\mathbf{b} + \mathbf{c}) = \mathbf{a} \cdot \mathbf{b} + \mathbf{a} \cdot \mathbf{c}$
4.  $(c\mathbf{a}) \cdot \mathbf{b} = c(\mathbf{a} \cdot \mathbf{b}) = \mathbf{a} \cdot (c\mathbf{b})$
5.  $\mathbf{0} \cdot \mathbf{a} = 0$

**Theorem 1.** If  $\theta$  is the angle between two vectors  $\mathbf{a}, \mathbf{b}$  then

$$\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}| \cos \theta$$

**Corollary 1.** Two vectors  $\mathbf{a}, \mathbf{b}$  are orthogonal iff  $\mathbf{a} \cdot \mathbf{b} = 0$

The **direction angles** of a non-zero vector  $\mathbf{a}$  are the angles  $\alpha, \beta, \gamma$  that  $\mathbf{a}$  makes with the positive  $x$ -,  $y$ -, and  $z$ -axes, respectively.

The cosines of these angles are called **direction cosines**:

$$\cos \alpha = \frac{\mathbf{a} \cdot \mathbf{i}}{|\mathbf{a}||\mathbf{i}|} = \frac{a_1}{|\mathbf{a}|}, \quad \cos \beta = \frac{a_2}{|\mathbf{a}|}, \quad \cos \gamma = \frac{a_3}{|\mathbf{a}|}$$

Some nice properties of direction cosines:

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$$

$$\frac{\mathbf{a}}{|\mathbf{a}|} = \langle \cos \alpha, \cos \beta, \cos \gamma \rangle$$

### Projections:

The **projection of  $\mathbf{b}$  onto  $\mathbf{a}$**  means we drop a perpendicular from the tip of  $\mathbf{b}$  onto the line spanned by  $\mathbf{a}$ . The resulting vector is called the vector projection of  $\mathbf{b}$  onto  $\mathbf{a}$ .

The **scalar projection of  $\mathbf{b}$  onto  $\mathbf{a}$**  is the length of this projection, given by

$$\text{comp}_{\mathbf{a}} \mathbf{b} = |\mathbf{b}| \cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|},$$

where  $\theta$  is the angle between  $\mathbf{a}$  and  $\mathbf{b}$ .

The **vector projection** of  $\mathbf{b}$  onto  $\mathbf{a}$  is obtained by multiplying the unit vector in the direction of  $\mathbf{a}$  by the scalar projection:

$$\text{proj}_{\mathbf{a}} \mathbf{b} = (\text{comp}_{\mathbf{a}} \mathbf{b}) \frac{\mathbf{a}}{|\mathbf{a}|} = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{a}|^2} \mathbf{a}$$

## 12.4 The Cross Product

**Definition 2.** For  $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$  and  $\mathbf{b} = \langle b_1, b_2, b_3 \rangle$ , the **cross product** of  $\mathbf{a}$  and  $\mathbf{b}$  is:

$$\mathbf{a} \times \mathbf{b} = \langle a_2 b_3 - a_3 b_2, a_3 b_1 - a_1 b_3, a_1 b_2 - a_2 b_1 \rangle = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

**Theorem 2.** The vector  $\mathbf{a} \times \mathbf{b}$  is orthogonal to both  $\mathbf{a}$  and  $\mathbf{b}$

**Theorem 3.** The length of the cross product is given by:

$$|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}| |\mathbf{b}| \sin \theta$$

where  $\theta$  is the angle between  $\mathbf{a}$  and  $\mathbf{b}$ .

**Corollary 2.** Two nonzero vectors  $\mathbf{a}$  and  $\mathbf{b}$  are parallel iff

$$\mathbf{a} \times \mathbf{b} = \mathbf{0}$$

**Area of parallelogram** determined by  $\mathbf{a}$  and  $\mathbf{b}$  is  $|\mathbf{a} \times \mathbf{b}|$

**Area of parallelogram** determined by  $\mathbf{a}$  and  $\mathbf{b}$  is  $\frac{1}{2} |\mathbf{a} \times \mathbf{b}|$

**IMPORTANT:** cross product is NOT commutative, and associate law for multiplication does not usually hold.

### Properties of the Cross Product

$\forall$  vectors  $\mathbf{a}, \mathbf{b}, \mathbf{c}$  and scalar  $c$ :

1.  $\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a}$
2.  $(c\mathbf{a}) \times \mathbf{b} = c(\mathbf{a} \times \mathbf{b}) = \mathbf{a} \times (c\mathbf{b})$
3.  $\mathbf{a} \times (\mathbf{b} + \mathbf{c}) = \mathbf{a} \times \mathbf{b} + \mathbf{a} \times \mathbf{c}$
4.  $(\mathbf{a} + \mathbf{b}) \times \mathbf{c} = \mathbf{a} \times \mathbf{c} + \mathbf{b} \times \mathbf{c}$
5.  $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \times \mathbf{b}) \cdot \mathbf{c}$
6.  $\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{a} \cdot \mathbf{c})\mathbf{b} - (\mathbf{a} \cdot \mathbf{b})\mathbf{c}$

### Triple products:

$$\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c}) = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

**Volume of the parallelepiped** determined by the vectors  $\mathbf{a}, \mathbf{b}$  and  $\mathbf{c}$  is:

$$V = |\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})|$$

Useful: if the volume is 0, then the vectors must lie in the same plane, i.e. **coplanar**

## 12.5 Equations of Lines and Planes

### 12.5.1 Equation of Lines

The equation of a line that goes through point with position vector  $\mathbf{r}_0$ , and is parallel to vector  $\mathbf{v}$  is

$$\mathbf{r} = \mathbf{r}_0 + t\mathbf{v}$$

If  $\mathbf{v} = \langle a, b, c \rangle$ ,  $\mathbf{r}_0 = \langle x_0, y_0, z_0 \rangle$  then

$$\langle x, y, z \rangle = \langle x_0 + ta, y_0 + tb, z_0 + tc \rangle$$

**Parametric form:**

$$x = x_0 + ta \quad y = y_0 + tb \quad z = z_0 + tc$$

**Direction numbers** of a line  $L$  are the numbers  $a, b, c$  from above.

**Symmetric form:** (eliminating  $t$ )

$$\frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}$$

IMPORTANT: If  $a, b$  or  $c$  is zero, then we cannot divide by them. For example, if  $a = 0$  then the equation becomes:

$$x = x_0, \quad \frac{y - y_0}{b} = \frac{z - z_0}{c}$$

In general, the symmetric equations between two points  $P_0(x_0, y_0, z_0), P_1(x_1, y_1, z_1)$  are

$$\frac{x - x_0}{x_1 - x_0} = \frac{y - y_0}{y_1 - y_0} = \frac{z - z_0}{z_1 - z_0}$$

The line segment from  $\mathbf{r}_0$  to  $\mathbf{r}_1$  is given by the vector equation:

$$\mathbf{r}(t) = (1 - t)\mathbf{r}_0 + t\mathbf{r}_1 \quad 0 \leq t \leq 1$$

### 12.5.2 Equation of Planes

Let:

- $\mathbf{n} = \langle a, b, c \rangle$ : vector orthogonal to plane
- $\mathbf{r}_0 = \langle x_0, y_0, z_0 \rangle$ : a position vector for a point in the plane
- $\mathbf{r} = \langle x, y, z \rangle$ : a position vector for an arbitrary point in the plane.

Then the **vector equation of the plane** is:

$$\mathbf{n} \cdot (\mathbf{r} - \mathbf{r}_0) = 0 \quad \text{or} \quad \mathbf{n} \cdot \mathbf{r} = \mathbf{n} \cdot \mathbf{r}_0$$

The **scalar equation** is:

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0$$

The **linear equation** is:

$$ax + by + cz - (ax_0 + by_0 + cz_0) = 0$$

Two planes are parallel if their normal vectors are parallel.

If two planes are not parallel, then they intersect in a straight line and the angle between the two planes is defined as the acute angle between their normal vectors.

The **distance**  $D$  from point  $(x_1, y_1, z_1)$  to the plane  $ax + by + cz + d = 0$  is

$$D = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$$