

Chapter 4 - Elementary Hilbert Space Theory

Salvador Castagnino
scastagnino@itba.edu.ar

Exercise Solutions

Exercise 4. We start by assuming that $\{u_n\}_{n \in \mathbb{N}}$ is a countable maximal orthonormal system in H , we want to see that H is separable. Observe that it suffices to find a countable set A such that $P \subset \overline{A}$, with P the set of all finite linear combinations of elements in $\{u_n\}_{n \in \mathbb{N}}$. Define the set A to be,

$$A = \left\{ \sum_{n=1}^N (q_n + ip_n) u_n : q_n, p_n \in \mathbb{Q}, N \geq 1 \right\}$$

which is clearly countable. Take $x \in P$, we have $x = \sum_{n=1}^N c_n u_n$ for some c_n complex numbers and some $N \geq 1$. Observe that for every $1 \leq n \leq N$ there exist $\{q_{nk}\}_{k \in \mathbb{N}}$ and $\{p_{nk}\}_{k \in \mathbb{N}}$ sequences of rational numbers such that $(q_{nk} + ip_{nk}) \rightarrow c_n$ as k goes to infinity. Then let $\epsilon > 0$ we can ask for a k large enough such that $|(q_{nk} + ip_{nk}) - c_n| < \frac{\epsilon}{N}$ for all $1 \leq n \leq N$, we in turn have

$$\left\| \sum_{n=1}^N (q_{nk} + ip_{nk}) u_n - \sum_{n=1}^N c_n u_n \right\| \leq \sum_{n=1}^N |(q_{nk} + ip_{nk}) - c_n| < \sum_{n=1}^N \frac{\epsilon}{N} < \epsilon$$

Given that every element in P can be approximated by elements in A we have that, $P \subset \overline{A}$, which concludes the proof.

Now suppose that $\{x_n\}_{n \in \mathbb{N}}$ is countable and dense in H , we are going to build a countable maximal orthonormal system. Let A be the set such that

$$\begin{cases} x_1 \in A \\ x_{n+1} \in A \text{ iff } x_{n+1} \notin [x_1, \dots, x_n] \end{cases}$$

where $[x_1, \dots, x_n]$ denotes the span of $\{x_1, \dots, x_n\}$, let's see that A is linearly independent. Suppose that for some v_1, \dots, v_{m+1} in A and $\alpha_1, \dots, \alpha_m$ nonzero complex numbers we have that

$$\sum_{j=1}^m \alpha_j v_j = v_{m+1}$$

Every v_j can be expressed as x_{n_j} , so let v_k be such that n_k is the greatest index between all n_j , we then have

$$v_k = \frac{v_{m+1}}{\alpha_k} - \sum_{1 \leq n \neq k \leq m} \frac{\alpha_n v_n}{\alpha_k}$$

which clashes with the construction of A , thus A is linearly independent. By **Ex4.3** we can build from A a countable orthonormal set $\{u_n\}$ such that $[v_1, \dots, v_n] = [u_1, \dots, u_n]$ for all $n \geq 1$. Given that every x_n can be expressed as a linear combination of elements in A , one can see that every x_n will be in $[u_1, \dots, u_m]$ for some m . Then we can see that every x_n is an element of \overline{P} , with P the set of finite linear combinations of elements in $\{u_n\}$, which in turn implies that P is dense in H and concludes the proof.

Exercise 6. We start by supposing that $\sum_{n=1}^{\infty} \delta_n^2 < \infty$, let's prove that S is compact. In order to do this, let $\{x_n\}_{n \in \mathbb{N}}$ be a sequence in S , we will construct a convergent subsequence.

- Let $y_{n1} = \hat{x}_n(1)$, notation as in the book. Given that the closed disk $\overline{B}(0, \delta_1)$ is compact and $y_{n1} \in \overline{B}(0, \delta_1)$, we can find y_{n_k1} a convergent subsequence of y_{n1} which converges to a y_1 in that interval. We replace our original sequence x_n for the new subsequence x_{n_k} , define $z_1 = x_{n_1}$ and proceed by repeating what we just did but for the second terms (we also rename the new sequence x_{n_k} as x_n for the sake of simplicity).
- Notation as above we can find a convergent subsequence y_{n_k2} of the sequence of second terms which converges in $\overline{B}(0, \delta_2)$ to a y_2 in that interval. We again define $z_2 = x_{n_2}$ and replace x_n by its subsequence x_{n_k} . Observe that the sequence of first terms still converges as it's a subsequence of a convergent sequence.

Repeating this process an arbitrary number of times we build a sequence z_n in S and a sequence of numbers y_n in $\overline{B}(0, \delta_n)$ for each n . Given that $0 \leq |y_n| \leq \delta_n$ for all $n \geq 1$ we have that $\sum_{n=1}^{\infty} |y_n|^2 < \infty$ and thus there exists a $y \in S$ such that $\hat{y}(n) = y_n$ for all $n \geq 1$. By the definition of the sequences z_n and y_n , given $\epsilon > 0$ we can find $N, M \in \mathbb{N}$ large enough such that,

$$\sum_{n=N}^{\infty} |\hat{z}_m(n) - y_n|^2 \leq \sum_{n=N}^{\infty} 4\delta_n^2 < \frac{\epsilon}{2} \quad (1)$$

$$|\hat{z}_m(n) - y_n|^2 < \frac{\epsilon}{2^{n+1}} \quad \forall 1 \leq n < N \quad (2)$$

for all $m \geq M$. With this in mind we can see that $\sum_{n=1}^{\infty} |\hat{z}_m(n) - y_n|^2 = \|z_m - y\|^2 < \epsilon$ for all $m \geq M$ which shows that z_n converges to y in S and concludes the proof.

Now suppose that S is compact and that $\sum_{n=1}^{\infty} \delta_n^2 = \infty$, we will get to a contradiction. Define $x_k = \sum_{n=1}^{\infty} c_{kn} u_n$ with

$$c_{kn} = \begin{cases} \delta_n & \text{if } n \leq k \\ 0 & \text{else} \end{cases}$$

Clearly $x_k \in S$ for all $k \geq 1$ and $\|x_k\| \rightarrow \sum_{n=1}^{\infty} \delta_n^2 = \infty$ but given that $\|x_k\| \leq \|x_k - x\| + \|x\|$ for every $x \in S$ the sequence x_n cannot converge which implies that S is not compact, contradicting our assumption and concluding the proof.

Useful Properties