

(5'-GTGGAGCAAATGCAGGACAACCGAATGCAG-3'; 5'-AGCAATGCAAGGACAACCGAATGCAGAAGG-3') in the *csp* gene of Pkpah-1. Furthermore, the alignment showed 21 variable sites between the genes. Phylogenetic analysis based on SSU rRNA sequences indicated that Pkpah-1 and Pkprk-1 formed a cluster, which was more related to *P. knowlesi* isolates from Thailand than to isolates from Sarawak, a Malaysian state in Borneo Island (Figure). Another isolate from Peninsular Malaysia, KAL-1, was also grouped with Pkpah-1 and Pkprk-1. The KAL-1 isolate was detected in a traveler from Finland returning from Peninsular Malaysia in 2007 (7).

*P. knowlesi* infection does not relapse because the parasite has no liver hypnozoite stage (8). Reinfection and recrudescence are not uncommon in malaria. The finding in the patient reported here indicates reinfection rather than recrudescence as confirmed by PCR genotyping. He acquired his infection while jungle trekking in Raub in October 2009 and was reinfected in Tanjung Malim in June 2010 (the great-circle distance between the 2 locations is 38.61 km). This case shows that immunity toward *P. knowlesi* infection is strain specific as has been observed in other malaria species (9). *P. knowlesi* is a nonhuman primate malaria species, and humans are accidental hosts when they go into areas where macaques dwell. *P. knowlesi* can cause reinfection and can potentially be severe in areas where it is endemic. Travelers to forested areas of endemicity should be advised to take strict antimosquito measures and prophylaxis. Physicians should be aware that *P. knowlesi* infection is a vital differential diagnosis in febrile travelers with a recent travel history to forested areas in Southeast Asia.

This study was supported by High Impact Research grant no. J-20011-73592 from the University of Malaya.

**Yee Ling Lau, Lian Huat Tan,  
Lit Chein Chin, Mun Yik Fong,  
Mydin Abdul-Aziz Noraishah,  
and Mahmud Rohela**

Author affiliations: University of Malaya, Kuala Lumpur, Malaysia (Y.L. Lau, L.C. Chin, M.Y. Fong, M.A.-A. Noraishah, M. Rohela); and Sunway Medical Centre, Selangor, Malaysia (L.H. Tan)

DOI: 10.3201/eid1707.101295

## References

1. Singh B, Lee KS, Matusop A, Radhakrishnan A, Shamsul SSG, Cox-Singh J, et al. A large focus of naturally acquired *Plasmodium knowlesi* infections in human beings. *Lancet*. 2004;363:1017–24. doi:10.1016/S0140-6736(04)15836-4
2. Vythilingam I, Noorazian YM, Huat TC, Jiram AI, Yusri YM, Azahari AH, et al. *Plasmodium knowlesi* in humans, macaques and mosquitoes in Peninsular Malaysia. *Parasit Vectors*. 2008;1:26. doi:10.1186/1756-3305-1-26
3. Singh B, Bobogare A, Cox-Singh J, Snounou G, Abdullah MS, Rahman HA. A genus- and species-specific nested polymerase chain reaction malaria detection assay for epidemiologic studies. *Am J Trop Med Hyg*. 1999;60:687–92.
4. Daneshvar C, Davis C, Cox-Singh J, Rafa'ee M, Zakaria S, Divis P, et al. Clinical and laboratory features of human *Plasmodium falciparum* infections. *Clin Infect Dis*. 2009;49:852–60. doi:10.1086/605439
5. McCutchan TF, Kissinger JC, Touray MG, Rogers MJ, Li J, Sullivan M, et al. Comparison of circumsporozoite proteins from avian and mammalian malarias: biological and phylogenetic implications. *Proc Natl Acad Sci U S A*. 1996;93:11889–94. doi:10.1073/pnas.93.21.11889
6. Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol*. 2007;24:1596–9. doi:10.1093/molbev/msm092
7. Kantele A, Marti H, Felger I, Müller D, Jokiranta TS. Monkey malaria in a European traveler returning from Malaysia. *Emerg Infect Dis*. 2008;14:1434–6. doi:10.3201/eid1409.080170
8. Krotoski WA, Collins WE. Failure to detect hypnozoites in hepatic tissue containing exoerythrocytic schizonts of *Plasmodium knowlesi*. *Am J Trop Med Hyg*. 1982;31:854–6.
9. Cheesman SO, Mahony E, Pattaradilokrat S, Degnan K, Knott S, Carter R. A single parasite gene determines strain-specific protective immunity against malaria: the role of the merozoite surface pro-

tein I. *Int J Parasitol*. 2010;40:951–61. doi:10.1016/j.ijpara.2010.02.003

Address for correspondence: Yee Ling Lau, Department of Parasitology, Faculty of Medicine, University of Malaya, 50603 Kuala Lumpur, Malaysia; email: lauyeeling@um.edu.my

## Antibody to Arenaviruses in Rodents, Caribbean Colombia

**To the Editor:** The ≈20 recognized arenaviruses in the Americas are hosted by rodents of the family *Cricetidae*; 1 exception may be hosted by a bat (genus *Artibeus*, family *Phyllostomidae*) (1). Pichindé virus, hosted by *Oryzomys albigularis*, was described from animals in the Pichindé Valley near Cali, Colombia (2), and antibody reactive to Pichindé virus was found in 2 of 82 serum samples from humans in the same area. No studies of arenavirus infection in rodents or humans have been conducted in Colombia since 1971. Although Pichindé virus is not associated with human disease, Guanarito virus, which is hosted by *Zygodontomys brevicauda*, the short-tailed cane mouse (3,4), causes Venezuelan hemorrhagic fever in the Venezuelan state of Portuguesa (5). This state borders on Colombia, and *Z. brevicauda* is a common species in Caribbean Colombia. Our aim was to determine the prevalence of antibody to arenaviruses among wild rodents in this region.

During November 1, 2008–June 10, 2009, we trapped 322 rodents in 3 rural localities in the Department of Córdoba, Colombia (Montería, Vereda El Escondido, 8°34.183'N,

75°42.776'W; Sahagun, Vereda Las Llanadas, 8°56.533'N, 75°20.909'W; and Lorica, Colegio Instituto Técnico Agrícola, 9°24.067'N, 075°75.707'W). The landscape at the study sites is dominated by tropical savanna, small patches of forest, and cultivated land. Live-capture traps were set in a variety of habitats at each locality, and captured rodents were processed according to methods by Mills et al. (6). Rodents were anesthetized, and blood (by cardiac puncture) and tissue (liver, lung, spleen, heart, kidney) samples were collected into individual cryovials, placed in liquid nitrogen in the field, and transferred to freezers and stored at -80°C at the Instituto de Investigaciones Biológicas del Trópico, Universidad de Córdoba, Montería, Colombia. Rodent species were identified on the basis of morphologic analyses of formalin-fixed carcasses; chromosomal data and mitochondrial DNA sequencing of the cytochrome b gene were used to confirm identification of most antibody-positive animals. Alcohol-preserved voucher specimens are archived at the Museum of Texas Tech University (Lubbock, TX, USA).

Blood samples were tested for arenavirus immunoglobulin (Ig) G by indirect immunofluorescent antibody assays. Guanarito and Pichindé virus-infected Vero E6 cells were used as antigens on spot slides. The secondary antibody was a fluorescein-conjugated goat,

antimouse IgG. Serum samples were screened at a dilution of 1:10 and endpoint titers were measured by using serial 2-fold dilutions (1:10–1:320) (7). Attempts to amplify viral RNA in tissues by reverse transcription PCR were unsuccessful.

We collected 210 sigmodontine rodents of 3 species: 181 *Z. brevicauda*, 28 *Oligoryzomys fulvescens*, and 1 *Oecomys concolor*. Eleven serum samples, 10 from *Z. brevicauda* and 1 from *O. fulvescens* rodents, had detectable arenavirus antibody. Three *Z. brevicauda* rodent samples had antibody reactive to both Pichindé and Guanarito virus, and 7 more were positive for either Pichindé or Guanarito arenaviruses (Table).

We used only 2 viral antigens in our screening belonging to the 2 viruses that are either known to occur in Colombia (Pichindé virus) or known to be hosted by species that we captured (Guanarito virus). Among the 10 *Z. brevicauda* samples with detectable antibody, 5 reacted only to Pichindé virus antigen or their antibody titer to Pichindé virus was at least 4-fold higher than their titer to Guanarito virus (Table), suggesting those rodents were infected with Pichindé or a closely related virus. Additional studies, including isolation and sequencing are needed to definitively identify this virus.

Surprisingly, only 2 *Z. brevicauda* rodent (1.1%) had antibody only to Guanarito virus or had a 4-fold greater

titer to Guanarito virus, much lower than the 15% antibody prevalence in the same species in the Venezuelan hemorrhagic fever-endemic area, Portuguesa State, Venezuela (5). Our testing protocols differed from the earlier study, and we have not definitively identified Guanarito virus in those 3 rodents; nevertheless, this low prevalence might help explain the absence of Venezuelan hemorrhagic fever in Colombia, although inadequate surveillance is a second possible explanation.

The single antibody-positive *O. fulvescens* rodent had a low antibody titer only to Pichindé virus. This apparent 4% antibody prevalence is based on only 28 mice. The significance of this finding is not clear but may represent spillover or an undescribed arenavirus specific to the species *O. fulvescens*. Again, additional studies are needed.

Our results demonstrate the presence of  $\geq 1$  arenaviruses circulating among common rodent hosts in Caribbean Colombia. We emphasize that many New World arenaviruses are likely cross-reactive to the antigens we used; recovery and sequencing of viral RNA will be essential to fully characterize these viruses. Hemorrhagic fever of arenaviral origin should be included in the differential diagnosis of tropical fevers, at least in our study region. As the human population of the rural Department of Córdoba and adjacent areas of

Table. Comparison of PICHV and GTOV antibody titers in rodents after serologic screening of 210 sigmodontine rodents, Córdoba, Colombia, November 1, 2008–June 10, 2009\*

Specimen no.	Species	PICHV titer	GTOV titer	Rural locality
49	<i>Zygodontomys brevicauda</i>	<b>40</b>	<10	Monteria
70	<i>Z. brevicauda</i>	<b>80</b>	10	Monteria
105	<i>Z. brevicauda</i>	<b>40</b>	<10	Monteria
107	<i>Oligoryzomys fulvescens</i>	20	<10	Monteria
272	<i>Z. brevicauda</i>	<b>80</b>	10	Monteria
289	<i>Z. brevicauda</i>	20	<10	Monteria
317	<i>Z. brevicauda</i>	<b>40</b>	<10	Monteria
344	<i>Z. brevicauda</i>	20	<10	Monteria
345	<i>Z. brevicauda</i>	<10	20	Monteria
209	<i>Z. brevicauda</i>	10	<b>80</b>	Lorica
211	<i>Z. brevicauda</i>	<10	<b>40</b>	Lorica

\*PICHV, Pichindé arenavirus; GTOV, Guanarito arenavirus. Titers in **boldface** are  $\geq 4$ -fold higher than titers for the other antigen.

the Caribbean coast of Colombia continues to increase, the potential for arenavirus-related disease could become a public health concern.

This study was supported by grant CIUC FMV0406-1120222 from the Universidad de Córdoba.

**Salim Mattar, Camilo Guzmán,  
Justiniano Arrazola, Ella Soto,  
José Barrios, Noemí Pini,  
Silvana Levis,  
Jorge Salazar-Bravo,  
and James N. Mills**

Author affiliations: Universidad de Córdoba, Montería, Colombia (S. Mattar, C. Guzmán, J. Arrazola, E. Soto, J. Barrios); Instituto Nacional de Enfermedades Virales Humanas "Dr. Julio I Maiztegui," Pergamino, Argentina (N. Pini, S. Levis); Texas Tech University, Lubbock, Texas, USA (J. Salazar-Bravo); and Centers for Disease Control and Prevention, Atlanta, Georgia, USA (J. Mills)

DOI: 10.3201/eid1707.101961

## References

- Salazar-Bravo J, Ruedas LA, Yates TL. Mammalian reservoirs of arenaviruses. *Curr Top Microbiol Immunol*. 2002;262:25–63.
- Trapido H, SanMartín C. Pichindé virus a new virus of the Tacaribe group from Colombia. *Am J Trop Med Hyg*. 1971;20:631–41.
- Fulhorst CF, Bowen MD, Salas RA, Duno G, Utrera A, Ksiazek TG, et al. Natural rodent host associations of Guanarito and Pirital viruses (family *Arenaviridae*) in central Venezuela. *Am J Trop Med Hyg*. 1999;61:325–30.
- Fulhorst CF, Cajimat MNB, Milazzo ML, Paredes H, de Manzione N, Salas RA, et al. Genetic diversity between and within the arenavirus species indigenous to western Venezuela. *Virology*. 2008;378:205–13. doi:10.1016/j.virol.2008.05.014
- Salas R, de Manzione N, Tesh RB, Rico-Hesse R, Shope RE, Betancourt A, et al. Venezuelan haemorrhagic fever. *Lancet*. 1991;338:1033–6. doi:10.1016/0140-6736(91)91899-6
- Mills JN, Yates T, Childs JE, Parmenter RR, Ksiazek TG, Rollin PE, et al. Guidelines for working with rodents potentially infected with hantavirus. *J Mammal*. 1995;76:716–22. doi:10.2307/1382742

- Bennett SG, Milazzo ML, Webb JP Jr, Fulhorst CF. Arenavirus antibody in rodents indigenous to coastal southern California. *Am J Trop Med Hyg*. 2000;62:626–30.

Address for correspondence: Jorge Salazar-Bravo, Department of Biological Sciences, Texas Tech University, Lubbock, TX 79409, USA; email: j.salazar-bravo@ttu.edu

## High Incidence of Guillain-Barré Syndrome in Children, Bangladesh

**To the Editor:** Bangladesh has achieved remarkable success in its drive to eliminate poliomyelitis; no case has been reported from that country since 2000. Still, the nonpolio incidence rate of acute flaccid paralysis (AFP) in Bangladesh is 3.25 cases per 100,000 children <15 years of age (1). Guillain-Barré syndrome (GBS), an acute polyradiculoneuropathy, is the most frequent cause of AFP (2). GBS in Bangladesh is frequently preceded by an enteric infection caused by *Campylobacter jejuni* (3). Frequent exposure to enteric pathogens at an early age may increase the incidence of GBS. We hypothesized that most AFP cases in Bangladesh can be diagnosed as GBS. Our objective was to estimate the crude incidence rate of GBS among children <15 years of age in Bangladesh.

In collaboration with the World Health Organization (WHO), the Government of Bangladesh conducts active AFP surveillance. AFP is defined as acute onset of focal or general flaccid (hypotonic) weakness without other obvious cause (e.g., trauma) in children <15 years of age.

Data on the number of reported AFP cases in Bangladesh during 2006 and 2007 were obtained. On the basis of clinical and other information routinely collected through the surveillance system, we defined a GBS case as presence of an acute flaccid (hypotonic) paralysis and symmetrical weakness (4) in the absence of injury or birth trauma.

Bangladesh is divided into 6 divisions (major administrative regions) comprising 64 districts. Crude incidence data for GBS were calculated per division and per district on the basis of the population <15 years of age reported by WHO and the Government of Bangladesh.

In 2006 and 2007, a total of 1,619 and 1,844 AFP cases, respectively, were reported in children <15 years of age, of which 608 (37%) and 855 (46%) cases, respectively, fulfilled the GBS case definition. The crude incidence rate for GBS in children <15 years of age varied from 1.5 to 1.7 cases per 100,000 population in the 3 northern divisions (Dhaka, Rajshahi, and Sylhet) and from 2.1 to 2.5 per 100,000 in the 3 southern divisions (Khulna, Barisal, and Chittagong) (online Appendix Figure, [www.cdc.gov/EID/content/17/7/1317-appF.htm](http://www.cdc.gov/EID/content/17/7/1317-appF.htm)). Overall, the crude incidence rate of GBS in children <15 years of age varied from 1.5 to 2.5 cases per 100,000 population per year in the 6 divisions of Bangladesh. Incidence rates were high (>5.0/100,000) in the Meherpur and Barisal districts in southern Bangladesh. We found a seasonal fluctuation in the frequency of patients with GBS; the most cases occurred in May (n = 159) and the lowest in February (n = 84). GBS occurred predominantly among boys (59%).

Most incidence studies reported in the literature originate from Europe and North America (5). A recent review reported that the best estimate of the global incidence of GBS in children <15 years of age is