



Short communication

The Bolivian bamboo rat, *Dactylomys boliviensis* (Rodentia: Echimyidae), a new record for chromosome number in a mammal

By J. L. DUNNUM, J. SALAZAR-BRAVO, and T. L. YATES

Museum of Southwestern Biology and Department of Biology, University of New Mexico, Albuquerque, New Mexico, U.S.A.

Receipt of Ms. 20. 09. 2000

Acceptance of Ms. 30. 10. 2000

Key words: *Dactylomys boliviensis*, Rodentia, Bolivia, new record

The family Echimyidae, a highly diverse group of rodents, occurs throughout most of Central and South America. There are 16 recent genera and about 70 living species (WOODS 1993), however, new taxa continue to be described (e.g., DA SILVA 1998; PATTON et al. 2000). To date no comprehensive phylogenetic analysis is available for the group although great advances have been forthcoming (e.g., LARA et al. 1996; PATTON et al. 2000). The group is highly diversified ecologically and has had a long evolutionary history in South America (PATTERSON and PASCUAL 1972; WOODS 1982). Karyologically, less than half of the species have been analyzed but it is known that there is considerable variation in diploid (from $2n = 14$ to $2n = 96$) and fundamental numbers (from $FNa = 18$ to $FNa = 144$) (Tab. 1). One of the most specialized groups within the Echimyidae is the subfamily of bamboo rats (Dactylomyinae). WOODS (1993) placed three genera in the subfamily Dactylomyinae; *Dactylomys*, *Kannabateomys*, and *Olallamys*. The biology and evolutionary relationships of the Dactylomyinae are poorly known, likely due to their rarity to collectors and subsequent scarcity in museum collections. What

is known is well summarized in PATTON et al. (2000). Until recently (ANISKIN 1993) no information on the chromosomal complement of any member of this group was available.

As part of a long-term survey of the mammals of Bolivia, many new and important records for the country were collected (ANDERSON 1997). In July of 1992 and May of 1996, we took a total of five specimens of *Dactylomys boliviensis* (Bolivian bamboo rat) from a locality in the Yungas of La Paz (SALAZAR et al. 1994). Here we report the karyotype of this species, the highest chromosomal number known in a mammal.

The individuals were located and collected at night in a dense stand of bamboo and secondary growth within the village of La Reserva (Departamento La Paz, Nor Yungas, La Reserva, elev. 840 m, $15^{\circ} 44' S$, $67^{\circ} 31' W$) by following their distinctive calls and eye shine. The village of La Reserva lies along Rio La Reserva, a small tributary of the Caranavi River. The village is at the bottom of a valley in the subtropical montane forest that covers most of the eastern Andean slopes between 15° and $17^{\circ} S$ latitude in the Cordillera Oriental of Bolivia. The foothills at this elevation are cov-



Fig. 1. Standard karyotype of *Dactylomys boliviensis*.

ered with semi-deciduous vegetation intermingled with columnar cacti and bromeliads. The forest is drier and sparser than at higher elevations. Compared to forests at higher and lower elevations, the trees are smaller, more highly branched, and most grow in open sun. The east facing slope above the river is steep, with much vegetation, some secondary growth, and banana and tangerine cultivation. Palms and tree ferns are absent (SALAZAR et al. 1994).

Chromosomal preparations were obtained using the technique described in ANDERSON et al. (1987). Metaphase cells were photographed and scored to determine the diploid ($2n$) and fundamental numbers (FNa). One of us (JLD) scored 5 slides per animal and over 20 spreads per slide to determine chromosome numbers. The analysis of the morphology of the chromosomes was based on 10 metaphase plates from three

individuals. Nomenclature for chromosome morphology and fundamental number follows PATTON (1967).

Chromosome slides, tissue samples, and cell suspensions are deposited in the Division of Biological Materials, Museum of Southwestern Biology (MSB). Voucher specimens are deposited at MSB (MSB 68547, MSB 85627, NK 40537), the American Museum of Natural History (AMNH 264887, 264884), and the Colección Boliviana de Fauna (CBF 2608), in La Paz, Bolivia.

The standard karyotype of *Dactylomys boliviensis* is highly asymmetrical, composed of 26 pairs of metacentric or sub-metacentric autosomes and 32 pairs of acrocentric autosomes. The X chromosome is a large sub-metacentric and the Y chromosome is a medium sub-metacentric. The resulting karyotype has a diploid count of $2n = 118$ and FNa of 168 (Fig. 1). Chromo-

Table 1. Diploid (2n) and fundamental number (FN) for members of the family Echimyidae.

Taxon	2n	FN	Reference
<i>Dactylomys boliviensis</i>	118	168	this report
<i>Dactylomys dactylinus</i>	94	144	ANISKIN (1993)
<i>Echimys blainvilliei</i>	50	94	REIG (1989)
<i>Echimys dasythrix</i>	96	102	LIMA et al. (1998)
<i>Echimys semivillosus</i>	94	134	AGUILERA et al. (1998)
<i>Echimys</i> sp.	90	108	LIMA et al. (1998)
<i>Echimys</i> sp.	90	110	ANISKIN (1993)
<i>Echimys</i> sp.	90	112	REIG (1989)
<i>Isothrix bistriata</i>	60	116	PATTON et al. (2000)
<i>Isothrix bistriata</i>	60	120	LIMA et al. (1998)
<i>Isothrix pagurus</i>	22	38	PATTON and EMMONS (1985)
<i>Isothrix sinamariensis</i>	28	42	VIE et al. (1996)
<i>Makalata armata</i>	70	120	LIMA et al. (1998)
<i>Makalata didelphoides</i>	66	106	LIMA et al. (1998)
<i>Clyomys laticeps</i>	34	60	REIG (1989)
<i>Euryzgomatomys guiana</i>	46	82	ANISKIN (1993)
<i>Euryzgomatomys spinosus</i>	46	92	REIG (1989)
<i>Hoplomys gymnurus</i>	46		ANISKIN (1993)
<i>Lonchothrix emiliae</i>	60	116	ANISKIN (1993)
<i>Mesomys hispidus</i>	60	120	LIMA et al. (1998)
<i>Mesomys hispidus</i>	60	116	PATTON et al. (2000)
<i>Mesomys occultus</i>	42	54	PATTON et al. (2000)
<i>Proechimys albispinus</i>	60	116	LEAL-MESQUITA et al. (1992)
<i>Proechimys amphicoricus</i>	26	44	REIG (1989)
<i>Proechimys breviceauda</i>	28-30	48-50	GARDNER and EMMONS (1984)
<i>Proechimys canicollis</i>	24	44	GARDNER and EMMONS (1984)
<i>Proechimys cuvieri</i>	28	46	MAIA and LANGGUTH (1993)
<i>Proechimys decumanus</i>	30	54	GARDNER and EMMONS (1984)
<i>Proechimys echinothrix</i>	32	69	PATTON et al. (2000)
<i>Proechimys gardneri</i>	40	56	PATTON et al. (2000)
<i>Proechimys goeldii</i>	24	44	PATTON et al. (2000)
<i>Proechimys guarae</i>	44-50	72-76	GARDNER and EMMONS (1984)
<i>Proechimys gularis</i>	30	48	GARDNER and EMMONS (1984)
<i>Proechimys guyanensis</i>	40	54-56	GARDNER and EMMONS (1984)
<i>Proechimys iheringi</i>	62-65	117-124	REIG (1989)
<i>Proechimys kulinae</i>	34	52	PATTON et al. (2000)
<i>Proechimys mincae</i>	48	68	GARDNER and EMMONS (1984)
<i>Proechimys oconnelli</i>	32	52	GARDNER and EMMONS (1984)
<i>Proechimys oris</i>	30	52-56	GARDNER and EMMONS (1984)
<i>Proechimys pattoni</i>	40	56	PATTON et al. (2000)
<i>Proechimys poliopus</i>	42	76	GARDNER and EMMONS (1984)
<i>Proechimys quadruplicatus</i>	28	44	GARDNER and EMMONS (1984)
<i>Proechimys semispinosus</i>	30	50-54	GARDNER and EMMONS (1984)
<i>Proechimys simonsi</i>	32	58	GARDNER and EMMONS (1984)
<i>Proechimys steerei</i>	24	42	GARDNER and EMMONS (1984)
<i>Proechimys trinitatus</i>	62	80	GARDNER and EMMONS (1984)
<i>Proechimys urichi</i>	62	88	GARDNER and EMMONS (1984)
<i>Proechimys yonenagae</i>	54	104	LEAL-MESQUITA et al. (1992), ROCHA (1995)
<i>Proechimys</i> sp.	34	56	ANISKIN (1993)
<i>Proechimys</i> sp.	14-16	18	REIG (1989)
<i>Proechimys</i> sp. (Balta)	40	56	REIG (1989)
<i>Proechimys</i> sp. (Barinas)	62	74	GARDNER and EMMONS (1984)
<i>Thricomys aperoides</i>	26	48	LEAL-MESQUITA et al. (1993)
<i>Thricomys aperoides</i>	30	54	REIG (1989)
<i>Thricomys aperoides</i>	30	50	ANISKIN (1993)




Fig. 2. Karyogram of known echimoid karyotypes.

some pair nine exhibits the characteristic satellite chromosome found in other echimoid rodents.

No chromosomal information is available for *Kannabateomys*, *Olallamys*, or *D. peruanus*. ANISKIN (1993) described the karyotype of *D. dactylinus* ($2n = 94$, $Fn = 144$) from the Loreto Department in Peru. The karyotype of *D. boliviensis* differs from that of *D. dactylinus* by the presence of one additional set of meta or sub-metacentric pairs, and 10 pairs of acrocentric chromosomes although comparisons are difficult due to the fact that ANISKIN (1993) did not identify sex chromosomes. At least 14 Robertsonian rearrangements would be necessary to transform the karyotype of one species into the other.

We compiled a list of all species of echimoid rodents for which data were available

(Tab. 1) and created a karyograph (IAMI and CROZIER 1980) based on chromosomal and fundamental numbers (Fig. 2). A definite pattern of subfamily grouping is clear where two species of *Dactylomys* assume the highest positions on the plot and the echimoid rodents (*Echimyus*, *Makalata*, *Isothrix*) are positioned at an intermediate level (with the exception of *I. pagurus* and *I. sinnamariensis*). The most speciose and karyologically studied group is the Eumysopinae (represented in this sample by *Proechimys*, *Clyomys*, *Euryzygomatomys*, *Hoplomys*, *Lonchothrix*, *Mesomys*, and *Thrichomys*). For the most part these fall at the lower end of (Fig. 2). To date, no eumysopids have been found with a $2n > 65$.

LIMA et al. (1998) proposed that Robertsonian rearrangements were more important in the evolution of the karyotype of arbo-

real echimyids than other chromosomal rearrangements because karyotypes of this group appeared to show higher levels of variation in diploid numbers than in fundamental numbers. Our data do not support LIMA et al. (1998). We found statistically significant differences in the levels of variation between diploid and fundamental number for the arboreal echimyids (Kruskal-Wallis; $P < 0.004$), terrestrial echimyids (One-way ANOVA; $\alpha = 0.05$; $P < 0.004$), and for the entire echimyid radiation (Kruskal-Wallis; $P < 0.001$). However, in all cases the fundamental number varied more than the diploid number, suggesting that pericentric inversions may be more common. None the less it is quite likely that several processes may have influenced the evolution of the karyotype in this group.

Acknowledgements

This work was supported by grants from the National Science Foundation and a Research Experience for Undergraduates supplement. We would like to thank the 1992 and 1996 members of the joint CBF-MNK-AMNH-MSB field expe-

Prior to our results, the highest chromosome number reported for a mammal was $2n = 102$ in *Tympanoctomys barrerae* (CONTRERAS et al 1990). These authors also suggested that the family Octodontidae presented the greatest chromosomal diversity. While this remains true for Fundamental number, the Echimyidae now represent the family with the greatest diversity in diploid number ($2n = 14$ to $2n = 118$).

Although *Tympanoctomys* and *Dactylomys* represent terminal branches in two different families of South American hystricognath rodents with a long history on this continent, they also share another characteristic: both occupy restricted ecological niches and possess highly specialized life history traits. We concur with CONTRERAS et al. (1990) in suggesting that the high chromosomal count appears to be a derived character.

References

- AGUILERA, M.; SANGINES, N.; PEREZ-ZAPATA, A. (1998): *Echymys semivillosus*, a rodent with a very high chromosomal number. *Caryologia* **51**, 181–187.
- ANDERSON, S. (1997). Mammals of Bolivia, Taxonomy and Distribution. *Bull. Am. Mus. Nat. Hist.* **231**.
- ANDERSON, S.; YATES, T. L.; COOK, J. A. (1987): Notes on Bolivian mammals 4: The genus *Ctenomys* (Rodentia, Ctenomyidae) in the eastern lowlands. *Am. Museum Novitates* **2891**, 1–20.
- ANISKIN, V. M. (1993): Three new karyotypes of prickly chinchillas of the family Echimyidae (Rodentia). *Genetika* **29**, 1500–1507.
- CONTRERAS, L. C.; TORRES-MURA, J. C.; SPOTORNO, A. E. (1990): The largest known chromosome number for a mammal, in a South American desert rodent. *Experientia* **46**, 506–508.
- DA SILVA, M. N. F. (1998): Four new species of spiny rats of the genus *Proechimys* (Rodentia: Echimyidae) from the western Amazon of Brazil. *Proc. Biol. Soc. Washington* **111**, 436–471.
- GARDNER, A.; EMMONS, L. (1984): Species groups in *Proechimys* (Rodentia, Echimyidae) as indicated by karyology and bullar morphology. *J. Mammalogy* **65**, 10–25.
- IMAI, H. T.; CROZIER, R. H. (1980): Quantitative analysis of directionality in mammalian karyotype evolution. *Am. Nat.* **116**, 537–569.
- LARA, M. C.; PATTON, J. L.; DA SILVA, M. N. F. (1996): The simultaneous diversification of South American echimyid rodents (Hystricognathi) based on complete cytochrome b sequences. *Mol. Phylog. Evol.* **5**, 403–413.
- LIMA, J. F. DE, S.; LANGGUTH, A.; DE SOUSA, L. C. (1998): The karyotype of *Makalata didelphoides* (Rodentia, Echimyidae). *Z. Säugetierkunde* **63**, 315–318.
- LEAL-MESQUITA, E. R.; FAGUNDES, V.; YONENAGA YASSUDA, Y.; ROCHA, P. DA. (1993): Com-

- parative cytogenetic studies of 2 karyomorphs of *Trictomys aperoides* (Rodentia, Echimyidae). *Rev. Bras Genetica* 16, 639-651.
- LEAL-MESQUITA, E.R.; YONENAGA YASSUDA, Y.; CHU TIEN, H.; ROCHA, P. da. (1992): Chromosomal characterization and comparative cytogenetic analysis of two species of *Proechimys* (Echimyidae, Rodentia) from the Caatinga domain of the state of Bahia, Brazil. *Caryologia* 45, 197-212.
- MAIA, V.; LANGGUTH, A. (1993): Constitutive heterochromatin polymorphism and NORs in *Proechimys cuvieri* Petter, 1978 (Rodentia, Echimyidae). *Rev. Bras Genetica* 16, 145-154.
- PATTERSON, B.; PASCUAL, R. (1972): The fossil mammal fauna of South America. In: *Evolution, Mammals and Southern Continents* Ed. by A. KEAST, F. ERK and B. GLASS. Albany: State University of New York Press Pp. 247-309.
- PATTON, J. L. (1967): Chromosome studies of certain pocket mice, genus *Perognathus* (Rodentia: Heteromyidae). *J Mammalogy* 48, 27-37.
- PATTON, J. L.; EMMONS, L. H. (1985): A review of the genus *Isothrix* (Rodentia: Echimyidae). *Am. Museum Novitates* 2817, 1-14.
- PATTON, J. L.; DA SILVA, M. N. F.; MALCOLM, J. R. (2000): Mammals of the Rio Jurua and the evolutionary and ecological diversification of Amazonia. *Bull. Am. Mus. Nat. Hist.* 244.
- REIG, O. (1989): Karyotypic repatterning as one triggering factor in cases of explosive speciation. In: *Evolutionary Biology of Unstable Populations*. Ed. by A. FONTDEVILA. Berlin, Heidelberg, Springer-Verlag Pp. 247-289.
- ROCHA, P. L. B. da (1995): *Proechimys yonenagae*, a new species of spiny rat (Rodentia, Echimyidae) from fossil sand dunes in the Brazilian Caatinga. *Mammalia* 59, 537-549.
- SALAZAR-BRAGO, J.; CAMPBELL, M.; ANDERSON, S.; GARDNER, S.; DUNNUM, J. (1994): New records of Bolivian mammals *Mammalia* 58, 125-130.
- VIE, J.; VOLOBOUEV, V.; PATTON, J.; GRANJON, L. (1996): A new species of *Isothrix* (Rodentia, Echimyidae) from French Guiana. *Mammalia* 60, 393.
- WOODS, C. A. (1982): The history and classification of South American hystricognath rodents: Reflections on the far away and long ago. In: *Mammalian Biology in South America*. Ed. by M. MARES and H. GENOWAYS. Pittsburgh, Pennsylvania: Pymatuning Laboratory of Ecology Vol. 6, 377-392.
- WOODS, C. A. (1993): Family Echimyidae (Gray, 1825). In: *Mammal Species of the World, A Taxonomic and Geographic Reference*. Ed. by D. E. WILSON and D. M. REEDER. Washington, DC: Smithsonian Institution Press Pp. 789-799.

Authors' address:

JONATHAN L. DUNNUM, JORGE SALAZAR-BRAGO, TERRY L. YATES, Museum of Southwestern Biology, Department of Biology, University of New Mexico, Albuquerque, New Mexico 87131, U.S.A.