

(a)
Razumovskiy et al. [8]
DFT (this work)
MD, random alloy
MD, ϕ = 0.6–0.9
MD, fully decomposed (ϕ = 1)
Fe-rich α
Fe–36Cr
Cr-rich α′
f_SAW (GHz), λ = 6.4 μm
x_Cr
(b)
Fe95Cr5, 25³ cell
Fe80Cr20, 25³ cell
Fe64Cr36, 25³ cell
Fe50Cr50 (Acta, 50³ cell)
Fe50Cr50, 25³ cell (validation)
Fe5Cr95, 25³ cell
f_SAW (GHz), λ = 6.4 μm
Spinodal conversion fraction, ϕ
(c)
Fe64Cr36 decomposing (MD, 25³ cells, outer faces): Fe-rich α and Cr-rich α′ domains form as ϕ goes 0.5 → 1
Fe
Cr
ϕ = 0.5
ϕ = 0.6
ϕ = 0.7
ϕ = 0.8
ϕ = 0.9
ϕ = 1.0
(d)
DFT cells relaxed in QE: bcc unit cells and 40-atom SQS (2×2×5), compositions of the DFT tensors (VASP and QE)
Fe
Cr
↑ moment
bcc Fe
20% Cr, SQS
35% Cr, SQS
50% Cr, SQS
65% Cr, SQS
95% Cr, SQS
bcc Cr