




6-Step Guide for Poultry Farming

1



- Use a wooden cage, or small room for brooding.
- Clean the Area : Wash and disinfect the brooding area.
- Cover the floor with clean and dry sawdust or newspaper.
- Provide a source of heat (charcoal stove, kerosene lamp, or electric bulb) to keep chicks warm).

Preparing the Brooder
(Before Chicks Arrive)

2



Receiving the Chicks

- Buy healthy, active, and vaccinated chicks.
- Gently place them in the warm brooder.
- Give them clean drinking water with glucose for energy.
- Feed them chick starter feed (fine grains rich in protein).
- Always keep the place clean and dry.

3



Keeping Chicks Warm (First 2 Weeks)

- Chicks need warmth to grow well.
- If they are gathered, they are cold, increase heat.
- If they are far apart with open mouths, they are hot, reduce heat.

4



Feeding and Watering (Week 1-8)


- Feed chicks 3-4 times daily with starter feed.
- Give clean water always-never dirty water!
- Add vitamins and antibiotics (as recommended by your Vet).
- Remove wet feed to prevent disease

5



Moving to Growers' Phase
(Week 3-8)

- As chicks grow, they need more space.
- Move them to a bigger pen with good ventilation.
- Start feeding grower feed (larger grains).
- Keep feeding and watering regularly.



Vaccination and Disease
Prevention (Week 1 - 4)

- Vaccinate against common poultry diseases like Newcastle.
- Keep the pen clean always—dirty pens cause sickness.
- Remove weak or sick birds to prevent spreading disease.

WATCH OUT !!! GOOD SIGNS



Chicks should be active, alert, and eating



Birds should be gaining weight and eating well



Normal droppings
(sign of healthiness)



Always wash your hands
before and after attending
to the birds.



2 weeks

4 Weeks

6-8 weeks

Feathers should start
growing by week 2

WATCH OUT !!! BAD SIGNS



Don't allow wet floor (can cause infections)




Don't mix sick chicks with healthy ones



Don't feed spoiled or moldy feeds



Ensure no weak, sick, or isolated chicks



They should not have wet droppings (sign of illness)



Check if chicks are huddled together (too cold) or spread apart (too hot)
If cold INCREASE HEAT
If hot, REDUCE HEAT

DO'S

- 
- ➔ Provide clean drinking water with glucose upon arrival
 - ➔ Feed starter mash 3-4 times daily
 - ➔ Keep the brooder clean and dry
 - ➔ Keep track of feed consumption
 - ➔ Record vaccinations and medications given
 - ➔ Monitor growth and mortality rates
 - ➔ Wash hands before and after handling birds
 - ➔ Restrict unnecessary visitors to the poultry house
 - ➔ Clean and disinfect equipment regularly
 - ➔ Ensure no signs of illness
 - ➔ Observe normal feeding and drinking habits



DON'TS ❌

- ⌚ Don't use dirty or contaminated water
- ⌚ Don't overcrowd chicks
- ⌚ Don't place the brooder in a cold or windy location
- ⌚ Don't expose birds to extreme cold or rain
- ⌚ Don't forget to clean feeders and drinkers
- ⌚ Don't keep too many birds in a small space

MAINTENANCE

- Water regularly
- Place staking material at the four corner of the bed.
- Attach the growing vines around the stake to support the growth.

Ready for Harvest

Sign (What to look for)

- Vines growing well with large leaves.
- 4–6 weeks after planting, leaves ready for cutting.

Action

- Harvest leaves gently by hand or cut with knife.



Watch for signs of sickness:

Not eating or drinking? **Sick!**

Droopy wings, closed eyes? **Sick!**

Running nose, difficulty breathing? **Sick!**

