

# Discrete Structures

Spring 2020

**Number Theory**

**Text book:** Kenneth H. Rosen, Discrete Mathematics and Its Applications

Section: 4.1

# Divisibility and Modular Arithmetic

Section 4.1

- When one integer is divided by a second nonzero integer, the quotient may or may not be an integer. For example,  $12/3 = 4$  is an integer, whereas  $11/4 = 2.75$  is not.

# Division

**Definition:** If  $a$  and  $b$  are integers with  $a \neq 0$ , then  $a$  *divides*  $b$  if there exists an integer  $c$  such that  $b = ac$

- When  $a$  divides  $b$  we say that  $a$  is a *factor* or *divisor* of  $b$  and that  $b$  is a multiple of  $a$ .
- The notation  $a \mid b$  denotes that  $a$  divides  $b$ .
- If  $a \mid b$ , then  $\frac{b}{a}$  is an integer.
- If  $a$  does not divide  $b$ , we write  $a \nmid b$ .



# Division

**Example:** Determine whether  $3 \mid 7$  and whether  $3 \mid 12$ .

**Solution:**  $3 \nmid 7$  because  $7/3$  is not an integer.  $3 \mid 12$  because  $12/3=4$ , which is an integer.

# Properties of Divisibility

**Theorem 1:** Let  $a$ ,  $b$ , and  $c$  be integers, where  $a \neq 0$ .

- i. If  $a \mid b$  and  $a \mid c$ , then  $a \mid (b + c)$ ;
- ii. If  $a \mid b$ , then  $a \mid bc$  for all integers  $c$ ;
- iii. If  $a \mid b$  and  $b \mid c$ , then  $a \mid c$ .

**Proof:** (i) Suppose  $a \mid b$  and  $a \mid c$ , then it follows that there are integers  $s$  and  $t$  with  $b = as$  and  $c = at$ . Hence,

$$b + c = as + at = a(s + t). \quad \text{Hence, } a \mid (b + c)$$

(similarly parts (ii) and (iii) can be proved) 

**Corollary:** If  $a$ ,  $b$ , and  $c$  be integers, where  $a \neq 0$ , such that  $a \mid b$  and  $a \mid c$ , then  $a \mid mb + nc$  whenever  $m$  and  $n$  are integers.

# Division Algorithm

- When an integer is divided by a positive integer, there is a quotient and a remainder. This is traditionally called the “Division Algorithm,”.

**Division Algorithm:** If  $a$  is an integer and  $d$  a positive integer, then there are unique integers  $q$  and  $r$ , with  $0 \leq r < d$ , such that  $a = dq + r$ .

- $d$  is called the *divisor*.
- $a$  is called the *dividend*.
- $q$  is called the *quotient*.
- $r$  is called the *remainder*.

Definitions of Functions **div**  
and **mod**

$$q = a \text{ div } d$$
$$r = a \text{ mod } d$$



# Division Algorithm

## Examples:

- What are the quotient and remainder when 101 is divided by 11?

**Solution:** The quotient when 101 is divided by 11 is  $9 = 101 \text{ div } 11$ , and the remainder is  $2 = 101 \text{ mod } 11$ .

- What are the quotient and remainder when  $-11$  is divided by 3?

**Solution:** The quotient when  $-11$  is divided by 3 is  $-4 = -11 \text{ div } 3$ , and the remainder is  $1 = -11 \text{ mod } 3$ .



# Congruence Relation

**Definition:** If  $a$  and  $b$  are integers and  $m$  is a positive integer, then  $a$  is congruent to  $b$  modulo  $m$  if  $m$  divides  $a - b$ .

- The notation  $a \equiv b \pmod{m}$  says that  $a$  is congruent to  $b$  modulo  $m$ .
- We say that  $a \equiv b \pmod{m}$  is a congruence and that  $m$  is its modulus.
- Two integers are congruent mod  $m$  if and only if they have the same remainder when divided by  $m$ .
- If  $a$  is not congruent to  $b$  modulo  $m$ , we write
$$a \not\equiv b \pmod{m}$$

# Congruence Relation

**Example:** Determine whether 17 is congruent to 5 modulo 6 and whether 24 and 14 are congruent modulo 6.

**Solution:**

- $17 \equiv 5 \pmod{6}$  because 6 divides  $17 - 5 = 12$ .
- $24 \not\equiv 14 \pmod{6}$  since 6 divides  $24 - 14 = 10$  is not divisible by 6.



# More on Congruences

**Theorem 4:** Let  $m$  be a positive integer. The integers  $a$  and  $b$  are congruent modulo  $m$  if and only if there is an integer  $k$  such that  $a = b + km$ .

**Proof:**

- If  $a \equiv b \pmod{m}$ , then (by the definition of congruence)  $m \mid a - b$ . Hence, there is an integer  $k$  such that  $a - b = km$  and equivalently  $a = b + km$ .
- Conversely, if there is an integer  $k$  such that  $a = b + km$ , then  $km = a - b$ . Hence,  $m \mid a - b$  and  $a \equiv b \pmod{m}$ . ◀



# The Relationship between $(\text{mod } m)$ and $\text{mod } m$ Notations

- The use of “mod” in  $a \equiv b \pmod{m}$  and  $a \text{ mod } m = b$  are different.
  - $a \equiv b \pmod{m}$  is a relation on the set of integers.
  - In  $a \text{ mod } m = b$ , the notation **mod** denotes a function.
- The relationship between these notations is made clear in this theorem.
- **Theorem 3:** Let  $a$  and  $b$  be integers, and let  $m$  be a positive integer. Then  $a \equiv b \pmod{m}$  if and only if  $a \text{ mod } m = b \text{ mod } m$ .

# Congruences of Sums and Products

**Theorem 5:** Let  $m$  be a positive integer. If  $a \equiv b \pmod{m}$  and  $c \equiv d \pmod{m}$ , then

$$a + c \equiv b + d \pmod{m} \text{ and } ac \equiv bd \pmod{m}$$

**Proof:**

- Because  $a \equiv b \pmod{m}$  and  $c \equiv d \pmod{m}$ , by Theorem 4 there are integers  $s$  and  $t$  with  $b = a + sm$  and  $d = c + tm$ .
- Therefore,
  - $b + d = (a + sm) + (c + tm) = (a + c) + m(s + t)$  and
  - $bd = (a + sm)(c + tm) = ac + m(at + cs + stm)$ .
- Hence,  $a + c \equiv b + d \pmod{m}$  and  $ac \equiv bd \pmod{m}$ .

**Example:** Because  $7 \equiv 2 \pmod{5}$  and  $11 \equiv 1 \pmod{5}$ , it follows from Theorem 5 that

$$18 = 7 + 11 \equiv 2 + 1 = 3 \pmod{5}$$

$$77 = 7 \cdot 11 \equiv 2 * 1 = 2 \pmod{5}$$





# Algebraic Manipulation of Congruences

- Multiplying both sides of a valid congruence by an integer preserves validity.  
If  $a \equiv b \pmod{m}$  holds then  $c \cdot a \equiv c \cdot b \pmod{m}$ , where  $c$  is any integer, holds by Theorem 5 with  $d = c$ .
- Adding an integer to both sides of a valid congruence preserves validity.  
If  $a \equiv b \pmod{m}$  holds then  $c + a \equiv c + b \pmod{m}$ , where  $c$  is any integer, holds by Theorem 5 with  $d = c$ .
- Dividing a congruence by an integer does not always produce a valid congruence.

**Example:** The congruence  $14 \equiv 8 \pmod{6}$  holds. But dividing both sides by 2 does not produce a valid congruence since  $14/2 = 7$  and  $8/2 = 4$ , but  $7 \not\equiv 4 \pmod{6}$ .



# Computing the $\bmod m$ Function of Products and Sums

- We use the following corollary to Theorem 5 to compute the remainder of the product or sum of two integers when divided by  $m$  from the remainders when each is divided by  $m$ .

**Corollary:** Let  $m$  be a positive integer and let  $a$  and  $b$  be integers. Then

$$(a + b) (\bmod m) = ((a \bmod m) + (b \bmod m)) \bmod m$$

and

$$ab \bmod m = ((a \bmod m) (b \bmod m)) \bmod m.$$

# Arithmetic Modulo $m$

**Definitions:** Let  $\mathbb{Z}_m$  be the set of nonnegative integers less than  $m$ :  $\{0, 1, \dots, m-1\}$

- The operation  $+_m$  is defined as  $a +_m b = (a + b) \bmod m$ . This is *addition modulo  $m$* .
- The operation  $\cdot_m$  is defined as  $a \cdot_m b = (a \cdot b) \bmod m$ . This is *multiplication modulo  $m$* .
- Using these operations is said to be doing *arithmetic modulo  $m$* .

**Example:** Find  $7 +_{11} 9$  and  $7 \cdot_{11} 9$ .

**Solution:** Using the definitions above:

- $7 +_{11} 9 = (7 + 9) \bmod 11 = 16 \bmod 11 = 5$
- $7 \cdot_{11} 9 = (7 \cdot 9) \bmod 11 = 63 \bmod 11 = 8$



# Arithmetic Modulo $m$

- The operations  $+_m$  and  $\cdot_m$  satisfy many of the same properties of ordinary addition and multiplication of integers.
  - **Closure**: If  $a$  and  $b$  belong to  $\mathbf{Z}_m$ , then  $a +_m b$  and  $a \cdot_m b$  belong to  $\mathbf{Z}_m$ .
  - **Associativity**: If  $a$ ,  $b$ , and  $c$  belong to  $\mathbf{Z}_m$ , then  $(a +_m b) +_m c = a +_m (b +_m c)$  and  $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$ .
  - **Commutativity**: If  $a$  and  $b$  belong to  $\mathbf{Z}_m$ , then  $a +_m b = b +_m a$  and  $a \cdot_m b = b \cdot_m a$ .
  - **Identity elements**: The elements 0 and 1 are identity elements for addition and multiplication modulo  $m$ , respectively.
    - If  $a$  belongs to  $\mathbf{Z}_m$ , then  $a +_m 0 = a$  and  $a \cdot_m 1 = a$ .

*continued* →



# Arithmetic Modulo $m$

- *Additive inverses*: If  $a \neq 0$  belongs to  $\mathbf{Z}_m$ , then  $m - a$  is the additive inverse of  $a$  modulo  $m$  and  $0$  is its own additive inverse.
  - $a +_m (m - a) = 0$  and  $0 +_m 0 = 0$
- *Distributivity*: If  $a$ ,  $b$ , and  $c$  belong to  $\mathbf{Z}_m$ , then
  - $a \cdot_m (b +_m c) = (a \cdot_m b) +_m (a \cdot_m c)$  and  $(a +_m b) \cdot_m c = (a \cdot_m c) +_m (b \cdot_m c)$ .
- Multiplicative inverses have not been included since they do not always exist. For example, there is no multiplicative inverse of  $2$  modulo  $6$ .