

UV-C Industrial Wand Testing Report

(Data according Signify latest report
with Boston Univeristy)

Testing Result for XBT LED System

| | Signify and Boston Univesity's Research | 1/4'' distance | 0.5'' distance | 1'' distance |
|------------------------------|--|----------------|----------------|--------------|
| Testing Data | 5mJ (6Seconds) Equals to 30mw/Cm2 (1 second) | 45.2 mw/Cm2 | 12.84 mw/Cm2 | 5.28 mw/Cm2 |
| Time to kill the Covid-19 | | 0.66Seconds | 2.34Seconds | 5.68Seconds |

Testing Result for XST LED System

| | Signify and Boston Univesity's Research | 1/4'' distance | 0.5'' distance | 1'' distance |
|------------------------------|--|----------------|-----------------|--------------|
| Testing Data | 5mJ (6Seconds) Equals to 30mw/Cm2 (1 second) | 56.2 mw/Cm2 | 14.84 mw/Cm2 | 5.92mw/Cm2 |
| Time to kill the Covid-19 | | 0.53Seconds | 2.02Seconds | 5Seconds |

UV-C Industrial Wand Testing Report

(Data according Philips latest research
with Boston University)



Testing Result for Single 25 LED System

| | Signify and Boston University's Research 5mJ (6Seconds) Equals to 30mw/Cm2 (1 second) | Touching the surface | 0.5'' distance | 1'' distance |
|------------------------------|---|----------------------|----------------|---------------|
| Testing Data | | 20.982 mw/Cm2 | 6.478 mw/Cm2 | 2.279 mw/Cm2 |
| Time to kill the Covid-19 | | 1.43 Seconds | 4.63 Seconds | 13.16 Seconds |

Testing Result for Double 25 LED System


| | Signify and Boston University's Research 5mJ (6Seconds) Equals to 30mw/Cm2 (1 second) | Touching the surface | 0.5'' distance | 1'' distance |
|------------------------------|---|----------------------|----------------|--------------|
| Testing Data | | 29.2 mw/Cm2 | 8.757 mw/Cm2 | 4.553 mw/Cm2 |
| Time to kill the Covid-19 | | 1.03 Seconds | 3.43 Seconds | 6.59 Seconds |

Testing Result for Single 64 LED System

| | Signify and Boston University's Research 5mJ (6Seconds) Equals to 30mw/Cm2 (1 second) | Touching the surface | 0.5'' distance | 1'' distance |
|------------------------------|---|----------------------|----------------|--------------|
| Testing Data | | 32.21 mw/Cm2 | 10.2 mw/Cm2 | 5.12 mw/Cm2 |
| Time to kill the Covid-19 | | 0.93 Seconds | 2.94 Seconds | 5.86 Seconds |

Testing Result for Double 64 LED System

| | Signify and Boston University's Research 5mJ (6Seconds) Equals to 30mw/Cm2 (1 second) | Touching the surface | 0.5'' distance | 1'' distance |
|------------------------------|---|----------------------|----------------|--------------|
| Testing Data | | 33.41 mw/Cm2 | 10.89 mw/Cm2 | 5.44 mw/Cm2 |
| Time to kill the Covid-19 | | 0.90 Seconds | 2.75 Seconds | 5.51 Seconds |



Rapid inactivation of SARS-CoV-2 with deep-UV LED irradiation

Hiroko Inagaki , Akatsuki Saito , Hironobu Sugiyama , Tamaki Okabayashi & Shouichi Fujimoto

To cite this article: Hiroko Inagaki , Akatsuki Saito , Hironobu Sugiyama , Tamaki Okabayashi & Shouichi Fujimoto (2020) Rapid inactivation of SARS-CoV-2 with deep-UV LED irradiation, Emerging Microbes & Infections, 9:1, 1744-1747, DOI: [10.1080/22221751.2020.1796529](https://doi.org/10.1080/22221751.2020.1796529)

To link to this article: <https://doi.org/10.1080/22221751.2020.1796529>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group, on behalf of Shanghai Shangyixun Cultural Communication Co., Ltd



Accepted author version posted online: 16 Jul 2020.
Published online: 31 Jul 2020.



Submit your article to this journal [↗](#)



Article views: 4254



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 2 View citing articles [↗](#)



LETTER



Rapid inactivation of SARS-CoV-2 with deep-UV LED irradiation

Hiroko Inagaki^a, Akatsuki Saito^{b,c}, Hironobu Sugiyama^{a,d}, Tamaki Okabayashi^{b,c} and Shouichi Fujimoto^e

^aM&N Collaboration Research Laboratory, Department of Medical Environment Innovation, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan; ^bDepartment of Veterinary Science, Faculty of Agriculture, University of Miyazaki, Miyazaki, Japan; ^cCenter for Animal Disease Control, University of Miyazaki, Miyazaki, Japan; ^dNikkiso Co., LTD, Tokyo, Japan; ^eDepartment of Hemovascular Medicine and Artificial Organs, Faculty of Medicine, University of Miyazaki, Miyazaki, Japan

ABSTRACT

The spread of novel coronavirus disease 2019 (COVID-19) infections worldwide has raised concerns about the prevention and control of SARS-CoV-2. Devices that rapidly inactivate viruses can reduce the chance of infection through aerosols and contact transmission. This *in vitro* study demonstrated that irradiation with a deep ultraviolet light-emitting diode (DUV-LED) of 280 ± 5 nm wavelength rapidly inactivates SARS-CoV-2 obtained from a COVID-19 patient. Development of devices equipped with DUV-LED is expected to prevent virus invasion through the air and after touching contaminated objects.

ARTICLE HISTORY Received 6 June 2020; Revised 8 July 2020; Accepted 9 July 2020

KEYWORDS COVID-19; deep-UV LED; inactivation; SARS-CoV-2; fomite infection; contact transmission; antiviral efficacy

Letter

The novel coronavirus SARS-CoV-2 pandemic has spread worldwide and placed countries in emerging, rapidly transforming situations. More than 5.3 million cases of COVID-19 and 342,000 deaths had been reported to WHO by 25 May 2020 [1]. Infectious virus is detected in specimens from the respiratory tract, nasopharyngeal sites, and feces in COVID-19 patients [2]. Recently, infectious SARS-CoV-2 was isolated from the urine of a COVID-19 patient [3]. SARS-CoV-2 is detectable in aerosols for up to 3 h, up to 4 h on copper, up to 24 h on cardboard and up to 2–3 days on plastic and stainless steel [4]. In nasal mucus and sputum, this virus is undetectable after 48 h, but its RNA remained detectable for 7 days [5]. To prevent exposure to contaminated material (fomite infection) which appears to be an important transmission route after droplet infection, hand hygiene with alcohol is key and efficient, but additional measures in preventing the spread of SARS-CoV-2 infection may be needed [6,7].

A deep ultraviolet light-emitting diode (DUV-LED) instrument generating around 250–300 nm wavelength has been reported to effectively inactivate microorganisms, including bacteria, viruses and fungi [8–11], but effects on SARS-CoV-2 have not been reported. We evaluated the antiviral efficacy of irradiation by DUV-LED, generating a narrow-range wavelength (280 ± 5 nm, mean \pm SD) (Nikkiso Co., Tokyo,

Japan), which we tested against SARS-CoV-2. This wavelength was selected due to higher output (radiation) power and longer lifetime in the stage of LED development considering practicality.

A strain of SARS-CoV-2 isolated from a patient who developed COVID-19 in the cruise ship *Diamond Princess* in Japan in February 2020 [12] was obtained from the Kanagawa Prefectural Institute of Public Health (SARS-CoV-2/Hu/DP/Kng/19-027, LC528233). The virus was propagated in Vero cells cultured in minimum essential medium (MEM) containing 2% fetal bovine serum (FBS). At 48 h after infection, virus stocks were collected by centrifuging the culture supernatants of infected Vero cells at 3000 rpm for 10 min. Clarified supernatants were kept at -80°C until use. Aliquots of stock virus were diluted with phosphate-buffered saline and adjusted to 2.0×10^4 plaque-forming units (PFU)/ml. For the evaluation of DUV-LED inactivation, aliquots of virus stock (150 μl) were placed in the centre of a 60-mm Petri dish and irradiated with $3.75 \text{ mW}/\text{cm}^2$ at work distance 20 mm for a range of times ($n = 3$ each time for 1, 10, 20, 30, or 60 s, and each dose corresponding to 3.75, 37.5, 75, 112.5 or $225 \text{ mJ}/\text{cm}^2$, respectively). After irradiation with DUV-LED, approximately 120 μl of each virus stock was collected with a 200 μl tip. Virus solutions were serially diluted in 10-fold steps using serum free MEM in a 1.5 ml tube, then inoculated onto Vero monolayers in a 12-well plate. After

CONTACT Shouichi Fujimoto fujimos@med.miyazaki-u.ac.jp Department of Hemovascular Medicine and Artificial Organs, Faculty of Medicine, University of Miyazaki, 5200 Kihara, Kiyotake, Miyazaki 889-1692, Japan

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group, on behalf of Shanghai Shangyixun Cultural Communication Co., Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

adsorption of virus for 2 h, cells were overlaid with MEM containing 1% carboxymethyl cellulose and 2% FBS (final concentration). Cells were incubated for 72 h in a CO₂ incubator, then cytopathic effects were observed under a microscope. An unirradiated virus suspension was used as a negative control. To calculate PFU, cells were fixed with 10% formalin for 30 min, followed by staining with 0.1% methylene blue solution. The antiviral effects of DUV-LED irradiations were assessed using the logPFU ratio, calculated as $\log_{10} \text{PFU ratio} = \log_{10} (N_t/N_0)$, where N_t is the PFU count of the UV-irradiated sample, and N_0 is the PFU count of the sample without UV irradiation. In addition, the infectious titer reduction rate was calculated as $(1 - 1/10^{\log_{10} \text{PFU ratio}}) \times 100 (\%)$. All experiments were performed in a BSL-3 laboratory.

We observed a marked cytopathic effect in virus-infected cells without DUV-LED irradiation (Figure 1A, see “0 s”). In contrast, virus-infected cells irradiated for 60 s showed largely comparable morphology to mock cells (Figure 1A, see “60 s”). To our surprise,

the cells inoculated with virus irradiated for 1 s looked similar to mock cells with minimal cytopathic effects (Figure 1A, see “1 s”). The plaque assay (Figure 1B) revealed that short time DUV-LED irradiation rapidly inactivated SARS-CoV-2 (Figure 1C and Table 1). Of note, the infectious titer reduction rate of 87.4% was already recognized with irradiation of virus stock for 1 s, and the rate was 99.9% with irradiation for 10 s. These results suggest that DUV-LED drastically inactivated SARS-CoV-2 with irradiation for even a very short time.

UV-LEDs providing irradiation at various peak emission wavelengths, such as UV-A (320–400 nm), UV-B (280–320 nm), and UV-C (100–280 nm), have been adopted to inactivate various pathogenic species, including bacteria, viruses and fungi. Devices equipped with UV-LEDs are now beginning to be introduced into medical fields. UV-C is considered to be the most effective germicidal region of the UV spectrum, acting through the formation of photoproducts in DNA and RNA [13]. These pyrimidine dimers




Figure 1. Inhibitory effects of DUV-irradiation on SARS-CoV-2. (A) Cytopathic changes in virus-infected Vero cells without DUV-LED irradiation (0 s), or with DUV-LED irradiation for 1, 10, 20, 30 or 60 s, and each dose corresponding to 3.75, 37.5, 75, 112.5 or 225 mJ/cm², respectively. (B) Plaque formation in Vero cells. Virus solutions irradiated with DUV-LED for several durations were diluted (100-fold) and inoculated to Vero cells. A representative result is shown. (C) Time-dependent inactivation of SARS-CoV-2 by DUV-LED irradiation. The results shown are the mean and standard deviation (SD) of triplicate measurements.

Table 1. Differences in infectious titer with different DUV-LED irradiation times.

| Irradiation time | Control (no irradiation) | DUV-LED irradiation time | | | | |
|--|--------------------------|--------------------------|-------------------|-------------------|-------|-------|
| | | 1 s | 10 s | 20 s | 30 s | 60 s |
| PFU (PFU/mL) | 3.7×10^4 | 4.7×10^3 | 2.7×10^1 | 6.7×10^0 | <20 | <20 |
| Log PFU ratio ^a | – | 0.9 | 3.1 | >3.3 | >3.3 | >3.3 |
| Infection titer reduction ratio ^b (%) | – | 87.4 | 99.9 | >99.9 | >99.9 | >99.9 |

^a $\log_{10} (Nt/NO)$ where Nt is the PFU count of the UV-irradiated sample and NO is the PFU count of the sample without UV irradiation.

^b $(1 - 1/10^{\log_{10} \text{PFU ratio}}) \times 100$ (%).

interrupt transcription, translation and replication of DNA and RNA, eventually leading to inactivation of microorganisms [14]. The efficacy of this inactivation may depend not only on the wavelength, but also on factors such as the target (e.g. bacterial species), light output and environmental conditions. The DUV-LED we used has the characteristics that can irradiate a narrow-range wavelength (280 ± 5 nm) compared with ordinary lamp, and achieve both long lifetime and high output power in time. This study demonstrated for the first time the rapid inactivation of SARS-CoV-2 under DUV-LED irradiation. As shown in Figure 1A, cytopathic effects were observed in control Vero cells infected with SARS-CoV-2, but not in the cells with DUV-LED irradiation for only 10 s. Whether UV-LED can similarly inactivate SARS-CoV-2 viruses existing in bodily fluid such as saliva needs to be investigated in the future. As well as in community settings, healthcare settings are also vulnerable to the invasion and spread of SARS-CoV-2, and the stability of SARS-CoV-2 in aerosols and on surfaces [4] likely contributes to virus transmission in medical environments. Although various monoclonal antibodies and clinical trials showing some results with first drugs (remdesivir) have been published [15], no vaccines are currently available for prevention and treatment of SARS-CoV-2. By revealing that SARS-CoV-2 inactivation can be achieved with very short-term DUV-LED irradiation, this study provides useful baseline data toward securing a safer medical environment. In the future, it will be necessary to validate the relationship between the radiation distance by DUV-LED and the inactivation of target virus. Development of devices equipped with DUV-LED is expected to prevent the virus invasion through the air and after touching contaminated objects.

Acknowledgements

We wish to thank Drs. Tomohiko Takasaki and Jun-Ichi Sakuragi, from the Kanagawa Prefectural Institute of Public Health, for providing the SARS-CoV-2/Hu/DP/Kng/19-027 strain.

Contributors

H.I. and H.S. conceived the study and wrote the manuscript. A.S. and T.O. conducted the experiments

dealing with viruses. S.F. contributed to the study design, study supervision and manuscript revision.

Disclosure statement

H.S. receives part of his salary from Nikkiso Co., Ltd., Tokyo, Japan. Nikkiso supplied the deep ultraviolet light-emitting diode (DUV-LED) instrument for evaluation. Nikkiso had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The other authors declare no conflicts of interest.

ORCID

Akatsuki Saito  <http://orcid.org/0000-0002-2792-4399>

Shouichi Fujimoto  <http://orcid.org/0000-0002-0025-4030>

References

- [1] Available from: https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200525-covid-19-sitrep-126.pdf?sfvrsn=887dbd66_2
- [2] Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in different types of clinical specimens. *JAMA*. 2020;323(18):1843–1844.
- [3] Sun J, Zhu A, Li H, et al. Isolation of infectious SARS-CoV-2 from urine of a COVID-19 patient. *Emerg Microbes Infect.* 2020;9(1):991–993.
- [4] van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med*. 2020;382(16):1564–1567.
- [5] Matson MJ, Kwe Yinda C, Seifert SN, et al. Effect of environmental conditions on SARS-CoV-2 stability in human nasal mucus and sputum. *Emerg Infect Dis*. Published online 2020 June 8.
- [6] Kampf G, Todt D, Pfaender S, et al. Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *J Hosp Infect*. 2020;104(3):246–251.
- [7] Kratzel A, Todt D, V'kovski P, et al. Inactivation of severe acute respiratory syndrome Coronavirus 2 by WHO-recommended hand rub formulations and alcohols. *Emerg Infect Dis*. 2020;26(7):1592–1595.
- [8] Rattanukul S, Oguma K. Inactivation kinetics and efficiencies of UV-LEDs against *Pseudomonas aeruginosa*, *Legionella pneumophila*, and surrogate microorganisms. *Water Res*. 2018;130:31–37.
- [9] Nishisaka-Nonaka R, Mawatari K, Yamamoto T, et al. Irradiation by ultraviolet light-emitting diodes inactivates influenza A viruses by inhibiting replication and transcription of viral RNA in host cells. *J Photochem Photobiol B*. 2018;189:193–200.
- [10] Kim DK, Kang DH. UVC LED irradiation effectively inactivates aerosolized viruses, bacteria, and fungi in

- a chamber-type air disinfection system. *Appl Environ Microbiol.* [2018](#);84(17):e00944–18.
- [11] Cheng Y, Chen H, Basurto LAS, et al. Inactivation of *Listeria* and *E. coli* by Deep-UV LED: effect of substrate conditions on inactivation kinetics. *Sci Rep.* [2020](#);10:3411.
- [12] Mallapaty S. What the cruise-ship outbreaks reveal about COVID-19. *Nature.* [2020](#);580(7801):18.
- [13] Bintsis T, Litopoulou-Tzanetaki E, Robinson RK. Existing and potential applications of ultraviolet light in the food industry – a critical review. *J Sci Food Agric.* [2000](#);80:637–645.
- [14] Kim SJ, Kim DK, Kang DH. Using UVC light-emitting diodes at wavelengths of 266 to 279 nanometers to inactivate foodborne pathogens and pasteurize sliced cheese. *Appl Environ Microbiol.* [2016](#);82(1):11–17.
- [15] Tu Y-F, Chien C-S, Yarmishyn AA, et al. A review of SARS-CoV-2 and the ongoing clinical trials. *Int J Mol Sci.* [2020](#);21(7):2657.