

Discussion Feb 16th, 4pm (neuer Hörsaal)

This practical gives a practical perspective on some aspects of the quasi-geostrophic approximation. We will use the Quasi-geostrophic barotropic vorticity with and without a forcing and the two-layer model to reflect on some of the theoretical findings derived during the semester.

We devote the final exercise on Feb 16th to a discussion of your results. Please hand in your code and a small exposé consisting of some figures showing results from the model prior to this discussion. We expect a brief discussion / presentation of the exposé during this exercise.

1 Preliminaries - model infrastructure

3P

You can use the model infrastructure provided for python that is provided on the whiteboard. (Alternatively, you may develop an infrastructure with similar capabilities in a programming/scripting language of your choice.)

Make sure you understand the following aspects before you get started to implement any other aspect of this assignment:

- How is the model controlled (how does it learn about its input / parameters / configurations?) ?
- Which classes exist and what is there purpose?

1. _____ – _____

2. _____ – _____

3. _____ – _____

4. _____ – _____

- What time integration scheme is used? In wich class and routine is it implemented?

Scheme _____

Class _____

Routine _____

- What kind of output is generated to the following streams/files

stdout _____

*.log _____

*.iter _____

*.nc _____

2 Time integration

3 P

Check the time integration scheme implemented in the class `INTEGRATOR`. Setting `rhs`¹ to `exponential` (a member of the class `integrator`) evaluates to the input, i.e. we solve the equation

$$\frac{dx}{dt} = x \Rightarrow \tilde{x}(t) = \frac{x_0}{e^{t_0}} e^t.$$

- a) What is the expected result for $t_0 = 0$ and $x_0 = 1$ when integrating up to $t = 2$? Check for $\Delta t = 1$!
- b) Varying the time step τ of the integration, check the order of the time integration scheme and produce a log-log plot of the error $x_{num} - \tilde{x}$ as a function of τ .
- c) What is the order of the numerical error for the time integration scheme used?
- d) Why does the numerical error reach a limit as $\tau \rightarrow 0$?

3 Rossby waves

15 P

4 Two-layer model

20 P

¹Note that with `rhs`, we are passing a function as argument, i.e. we are using the concept of function pointers