


# Elementary Linear Algebra



## Chapter 2

Howard Anton

Copyright © 2014 by John Wiley & Sons, Inc.

All rights reserved.



# Chapter 2

## Determinants

- 2.1 Determinants by Cofactor Expansion
- 2.2 Evaluating Determinants by Row Reduction
- 2.3 Properties of Determinants; Cramer's Rule

# Section 2.1    Determinants by Cofactor Expansion

**DEFINITION 2** If  $A$  is an  $n \times n$  matrix, then the number obtained by multiplying the entries in any row or column of  $A$  by the corresponding cofactors and adding the resulting products is called the *determinant of  $A$* , and the sums themselves are called *cofactor expansions of  $A$* . That is,

$$\det(A) = a_{1j}C_{1j} + a_{2j}C_{2j} + \cdots + a_{nj}C_{nj} \quad (7)$$

[cofactor expansion along the  $j$ th column]

and

$$\det(A) = a_{i1}C_{i1} + a_{i2}C_{i2} + \cdots + a_{in}C_{in} \quad (8)$$

[cofactor expansion along the  $i$ th row]

► **EXAMPLE 3** Cofactor Expansion Along the First Row

Find the determinant of the matrix

$$A = \begin{bmatrix} 3 & 1 & 0 \\ -2 & -4 & 3 \\ 5 & 4 & -2 \end{bmatrix}$$

by cofactor expansion along the first row.

*Solution*

$$\begin{aligned} \det(A) &= \begin{vmatrix} 3 & 1 & 0 \\ -2 & -4 & 3 \\ 5 & 4 & -2 \end{vmatrix} = 3 \begin{vmatrix} -4 & 3 \\ 4 & -2 \end{vmatrix} - 1 \begin{vmatrix} -2 & 3 \\ 5 & -2 \end{vmatrix} + 0 \begin{vmatrix} -2 & -4 \\ 5 & 4 \end{vmatrix} \\ &= 3(-4) - (1)(-11) + 0 = -1 \end{aligned}$$

► **EXAMPLE 4** Cofactor Expansion Along the First Column

Let  $A$  be the matrix in Example 3, and evaluate  $\det(A)$  by cofactor expansion along the first column of  $A$ .

*Solution*

$$\begin{aligned} \det(A) &= \begin{vmatrix} 3 & 1 & 0 \\ -2 & -4 & 3 \\ 5 & 4 & -2 \end{vmatrix} = 3 \begin{vmatrix} -4 & 3 \\ 4 & -2 \end{vmatrix} - (-2) \begin{vmatrix} 1 & 0 \\ 4 & -2 \end{vmatrix} + 5 \begin{vmatrix} 1 & 0 \\ -4 & 3 \end{vmatrix} \\ &= 3(-4) - (-2)(-2) + 5(3) = -1 \end{aligned}$$

This agrees with the result obtained in Example 3.

# A technique for determinants of 2x2 and 3x3 matrices only

## ▶ EXAMPLE 7 A Technique for Evaluating 2 x 2 and 3 x 3 Determinants

$$\begin{vmatrix} 3 & 1 \\ 4 & -2 \end{vmatrix} = \begin{vmatrix} 3 & 1 \\ 4 & -2 \end{vmatrix} = (3)(-2) - (1)(4) = -10$$

$$\begin{vmatrix} 1 & 2 & 3 \\ -4 & 5 & 6 \\ 7 & -8 & 9 \end{vmatrix} = \begin{vmatrix} 1 & 2 & 3 & 1 & 2 \\ -4 & 5 & 6 & -4 & 5 \\ 7 & -8 & 9 & 7 & -8 \end{vmatrix}$$
$$= [45 + 84 + 96] - [105 - 48 - 72] = 240 \quad \blacktriangleleft$$



# Row Reduction and Determinants

**THEOREM 2.2.3** Let  $A$  be an  $n \times n$  matrix.

- (a) If  $B$  is the matrix that results when a single row or single column of  $A$  is multiplied by a scalar  $k$ , then  $\det(B) = k \det(A)$ .
- (b) If  $B$  is the matrix that results when two rows or two columns of  $A$  are interchanged, then  $\det(B) = -\det(A)$ .
- (c) If  $B$  is the matrix that results when a multiple of one row of  $A$  is added to another row or when a multiple of one column is added to another column, then  $\det(B) = \det(A)$ .

Table 1

Relationship	Operation
$\begin{vmatrix} ka_{11} & ka_{12} & ka_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = k \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$ $\det(B) = k \det(A)$	The first row of $A$ is multiplied by $k$ .
$\begin{vmatrix} a_{21} & a_{22} & a_{23} \\ a_{11} & a_{12} & a_{13} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = - \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$ $\det(B) = -\det(A)$	The first and second rows of $A$ are interchanged.
$\begin{vmatrix} a_{11} + ka_{21} & a_{12} + ka_{22} & a_{13} + ka_{23} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$ $\det(B) = \det(A)$	A multiple of the second row of $A$ is added to the first row.

# Section 2.3

## Cramer's Rule

### THEOREM 2.3.7 Cramer's Rule

*If  $A\mathbf{x} = \mathbf{b}$  is a system of  $n$  linear equations in  $n$  unknowns such that  $\det(A) \neq 0$ , then the system has a unique solution. This solution is*

$$x_1 = \frac{\det(A_1)}{\det(A)}, \quad x_2 = \frac{\det(A_2)}{\det(A)}, \dots, \quad x_n = \frac{\det(A_n)}{\det(A)}$$

*where  $A_j$  is the matrix obtained by replacing the entries in the  $j$ th column of  $A$  by the entries in the matrix*

$$\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_n \end{bmatrix}$$

# Section 2.3

## Cramer's Rule

► **EXAMPLE 8** Using Cramer's Rule to Solve a Linear System

Use Cramer's rule to solve

$$\begin{aligned}x_1 + \quad + 2x_3 &= 6 \\-3x_1 + 4x_2 + 6x_3 &= 30 \\-x_1 - 2x_2 + 3x_3 &= 8\end{aligned}$$

*Solution*

$$A = \begin{bmatrix} 1 & 0 & 2 \\ -3 & 4 & 6 \\ -1 & -2 & 3 \end{bmatrix}, \quad A_1 = \begin{bmatrix} 6 & 0 & 2 \\ 30 & 4 & 6 \\ 8 & -2 & 3 \end{bmatrix},$$
$$A_2 = \begin{bmatrix} 1 & 6 & 2 \\ -3 & 30 & 6 \\ -1 & 8 & 3 \end{bmatrix}, \quad A_3 = \begin{bmatrix} 1 & 0 & 6 \\ -3 & 4 & 30 \\ -1 & -2 & 8 \end{bmatrix}$$

Therefore,

$$x_1 = \frac{\det(A_1)}{\det(A)} = \frac{-40}{44} = \frac{-10}{11}, \quad x_2 = \frac{\det(A_2)}{\det(A)} = \frac{72}{44} = \frac{18}{11},$$
$$x_3 = \frac{\det(A_3)}{\det(A)} = \frac{152}{44} = \frac{38}{11} \quad \blacktriangleleft$$