

Based on National Curriculum 2022-23

Textbook of

# Mathematics

GRADE

10



National Book Foundation

Ministry of Federal Education and Professional Training

Government of Pakistan

#### **Government Approval**

Approved by the National Curriculum Council (NCC), Ministry of Federal Education and Professional Training, Islamabad  
vide letter No. F.1(1-10)/2024/NCP-NOC/NBF/Maths, dated 19th September, 2024

#### **© 2025 National Book Foundation (NBF) as Federal Textbook Board**

All rights to this publication are strictly reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means—electronic, mechanical, photocopying, recording, or otherwise—without the prior express written consent of the publisher. Unauthorized use, duplication, or distribution is strictly prohibited and may result in legal action.

#### **A Textbook of Mathematics for Grade 10 Based on National Curriculum of Pakistan (NCP) 2022-23**

##### **Authors**

Dr. Saleem Ullah (Managing Author)  
Dr. Khalid Mahmood, Ms. Nabeela Azad, Ms. Hafsa Jabeen (Co-Authors)

##### **Contributors/Collaborators**

Muhammad Ahmad Nawaz (Designer)

---

##### **Supervision of Curriculum and Review Process**

Dr. Shafquat Ali Janjua  
Joint Educational Advisor, National Curriculum Council (NCC)  
Ministry of Federal Education and Professional Training, Government of Pakistan, Islamabad

##### **NCC Review Committee Members**

Mr. Abdul Rauf, EST, FG Public School, Mehfooz Road, Rawalpindi, FGEI's  
Major General Retired Malik Muhammad Abbas, Ms. Uzma Malik, Psychologist

##### **Desk Officer**

Zehra Khushal, (Assistant Educational Advisor), National Curriculum Council (NCC)

---

##### **NBF Textbooks Development Supervision**

Murad Ali Mohmand  
Managing Director, National Book Foundation (NBF)

##### **In-Charge, NBF Textbooks Development**

Mansoor Ahmad, Assistant Director

##### **Printed in Pakistan**

First Edition: First Impression: February, 2025 | Pages: 300 | Quantity: 100000

Price: 400/-, Code: STE-724, ISBN: 978-969-37-1726-6

Printer: Al-Qazi Printers, Peshawar



For details on additional publications from the National Book Foundation, please visit our website at [www.nbf.org.pk](http://www.nbf.org.pk)

You can also reach us by phone at 051 9261125 or via email at [books@nbf.org.pk](mailto:books@nbf.org.pk)

For feedback or corrections, kindly send your comments to '[nbftextbooks@gmail.com](mailto:nbftextbooks@gmail.com)' and '[textbooks@snc.gov.pk](mailto:textbooks@snc.gov.pk)'

##### **Note**

All illustrations, artwork, and images in this book are intended solely for educational and promotional purposes, benefitting the public interest.

# PREFACE

Textbook of Mathematics for Grade 10 has been developed by NBF according to the National Curriculum of Pakistan 2022- 2023. The aim of this experimentation skills is to enhance learning abilities through inculcation of logical thinking in learners, and to develop higher order thinking processes by systematically building upon the foundation of learning from the previous grades. A key emphasis of the present experimentation skills is on creating real life linkages of the concepts and methods introduced. This approach was devised with the intent of enabling students to solve daily life problems as they go up the learning curve and for them to fully grasp the conceptual basis that will be built upon in subsequent grades.

After amalgamation of the efforts of experts and experienced author, this book was reviewed and finalized after extensive reviews by professional educationists. Efforts were made to make the contents student friendly and to develop the concepts in interesting ways.

The National Book Foundation is always striving for improvement in the quality of its books. The present book features an improved design, better illustration and interesting activities relating to real life to make it attractive for young learners. However, there is always room for improvement and the suggestions and feedback of students, teachers and the community are most welcome for further enriching the subsequent editions of this book.

May Allah guide and help us (Ameen).

Murad Ali Mohmand

Managing Director

# Applications of Mathematics

**Complex Numbers:** Complex numbers play a critical role in fields such as engineering, electronics, and quantum physics. They are especially valuable for modeling oscillations in alternating current (AC) circuits, which are foundational to modern electrical systems, and for analyzing waveforms and resonances that occur naturally in many types of physical systems.

**Quadratic Equations:** Quadratic equations model a range of phenomena. In physics, they describe projectile motion; in finance, they are used for profit and cost optimization; and in architecture, they inform structural designs by calculating load distributions and parabolic shapes for stability.

**Matrices and Determinants:** Matrices are essential in solving complex systems of equations and executing transformations in areas like computer graphics and machine learning. Determinants, closely related to matrix invertibility, provide insights into properties of shapes and can be used to calculate area, volume, and to determine if linear systems have unique solutions.

**Linear and Quadratic Inequalities:** Linear and quadratic inequalities express ranges of values that satisfy given conditions, such as acceptable income levels, speed limits, or temperature thresholds. In real life, they are essential in budgeting, optimizing resources in industries, setting safety regulations, and environmental monitoring, where certain parameters must stay within safe or efficient boundaries.

**Algebraic Fractions:** Algebraic fractions simplify relationships found in finance (like calculating compound interest), physics (such as wave motion and light refraction), and data science (for probability ratios and odds in statistical models). They allow complex relationships to be expressed in simpler, comparable terms.

**Functions and Graphs:** Functions are mathematical tools that map inputs to outputs, visualized through graphs to represent relationships between variables, such as speed over time, cost versus quantity, or population growth over years. These graphical representations are fundamental in economics, biology, and physics, where understanding variable relationships is key to predicting and modeling real-world changes.

**Vectors:** Vectors combine magnitude and direction, making them indispensable in applications like navigation, physics (force and velocity calculations), and engineering (stress analysis in materials). In computer graphics, they help in rendering movements and 3D transformations, contributing to realistic visualizations.

**Trigonometry:** Trigonometry is crucial for solving right and non-right triangles. Its applications span navigation, construction, and astronomy, where precise distance and angle measurements are essential, from mapping locations to constructing stable structures.

**Chords and Arcs of a Circle:** Understanding chords and arcs forms the basis of circular geometry, used extensively in design, architecture, and mechanical engineering, where curves and circular components (like gears or wheels) must be accurately measured and constructed.

**Tangents and Its Construction:** A tangent touches a circle at one point, forming a right angle with the radius at that point. This principle has practical applications in road and race track design gear construction, and analyzing rotational motion, where precision at points of contact is crucial. Constructing tangents, arcs, and circular sections is fundamental in structural design, architecture, bridge engineering, and urban planning.

**Statistics:** Statistics, through measures like mean, median, mode, range, and standard deviation, help summarize and interpret data trends. Statistics play an essential role in business forecasting, healthcare analytics, social sciences, and environmental studies, where decisions must be made based on reliable data insights.

## Table of Contents

Sr. No.	Description	P. No.
1	Complex Numbers	6
2	Quadratic Equations	27
3	Matrices And Determinants	55
4	Linear and Quadratic Inequalities	78
5	Algebraic Fractions	97
6	Functions And Graphs	115
7	Vectors in Plane	140
8	Application of Trigonometry	160
9	Chord and Arcs of a Circle	196
10	Tangents and Angles of a Circle	215
11	Practical Geometry of Circles	232
12	Basic Statistics	245
13	Answers	279

## 01

# Complex Numbers

**After studying this unit students will be able to:**

- Recognize complex numbers, conjugate and modulus of a complex numbers.
- Apply basic operations on complex numbers.
- Define commutative laws, associative laws and distributive laws for complex numbers.
- Recognize additive and multiplicative identity and find additive and multiplicative inverse of complex numbers.
- Find real and imaginary parts of  $\left(\frac{x_1 + iy}{x_2 + iy}\right)^n$  for  $n = \pm 1, \pm 2, \dots$
- Find solution of equations having complex values.
- Apply the geometric interpretation of a complex numbers.
- Solve daily life problems involving complex numbers.

Until the sixteenth century, mathematicians were puzzled by square roots of a negative numbers.

For example, the solutions of  $x^2 - 2 = 0$  are  $\sqrt{2}$ , and  $-\sqrt{2}$ . But what are the solution of  $x^2 + 2 = 0$ ? To find the solution of this equation, imaginary numbers were defined.

A French mathematician René Descartes was the first to emphasize the imaginary nature of numbers, positing that “one can imagine as many (numbers) as already mentioned in each equation, but sometimes, there is no quantity that matches what we imagine.”

However, in 16th century an Italian mathematician Gerolamo Cardano, proved that having a negative term inside a square root can lead to the solution of an equation.


Later, in the 18th century, mathematician Carl Friedrich Gauss consolidated Cardano's premises, in addition to developing a treatise on complex numbers in a plane and thereby established the modern bases of the term.



René Descartes



Gerolamo Cardano



Carl Friedrich Gauss

# Imaginary and Complex Numbers

Negative numbers do not have square roots in the real number system. However, a larger number system that contains the real number system is designed so that negative numbers do have square roots. This system is called the complex number system and it makes use of a number that is a square root of  $-1$ . We call this new number  $i$ .

Let us find the solution of:

$$x^2 + 1 = 0 \quad (\text{i})$$

From (i),  $x^2 = -1$  which implies  $x = \pm\sqrt{-1} = \pm i$

The number  $i$  is the solution of  $x^2 + 1 = 0$ , and is defined as:

$i = \sqrt{-1}$  where  $i^2 = -1$

Since the square of a real number is not negative, therefore  $i$  is not a real number and is called the imaginary unit.

Using  $i$  we can define square root of any negative number.

## Interesting Information

Girolamo Cardano began his career as a doctor and studied, taught and wrote mathematics as a side line. He held important positions at The University of Pavia and Bologna in Italy and wrote many works on arithmetic, astronomy, physics and medicine.

To express the roots of negative numbers in terms of  $i$ , we can use the fact that in the complex numbers,  $\sqrt{-x} = \sqrt{-1}\sqrt{x} = i\sqrt{x}$ . When  $x$  is a positive real number and  $i$  is the imaginary number, the number  $i\sqrt{x}$  is called a pure imaginary number.

**Example:** Express each number in terms of  $i$ .

a.  $\sqrt{-5}$

b.  $\sqrt{-9}$

c.  $\sqrt{-17}$

d.  $-\sqrt{-64}$

e.  $3i \times 5i$

f.  $\sqrt{-5} \times \sqrt{-20}$

**Solution:**

a.  $\sqrt{-5} = \sqrt{-1 \times 5} = \sqrt{-1} \times \sqrt{5} = i\sqrt{5} = \sqrt{5}i$

b.  $\sqrt{-9} = \sqrt{-1 \times 9} = \sqrt{-1} \times \sqrt{9} = i \times 3 = 3i$

We prefer to write  $ia$  as  $ai$ , where  $a$  is rational number.

c.  $-\sqrt{-17} = -\sqrt{-1 \times 17} = -\sqrt{-1} \times \sqrt{17} = -i\sqrt{17} = -\sqrt{17}i$

d.  $-\sqrt{-64} = -\sqrt{-1 \times 64} = -\sqrt{-1} \times \sqrt{64} = -i\sqrt{64} = -i8 = -8i$

e.  $3i \times 5i = 15i^2 = 15 \times (-1) = -15 \quad \text{where } (i^2 = -1)$

f.  $\sqrt{-5} \times \sqrt{-20} = \sqrt{-1 \times 5} \times \sqrt{-1 \times 20} = (\sqrt{-1} \times \sqrt{5}) \times (\sqrt{-1} \times \sqrt{20})$   
 $= i\sqrt{5} \times i\sqrt{20} = i^2\sqrt{5 \times 20} = -1\sqrt{100} = -10$

## Imaginary Number

An imaginary number is a number that can be written  $a+bi$ , where  $a$  and  $b$  are real numbers,  $b \neq 0$ .


## Complex Numbers

When  $a$  and  $b$  are real numbers and  $b$  is allowed to be 0, the number  $a+bi$  is said to be complex.

A complex number is any number that can be written in the form of  $a+bi$ , where  $a$  and  $b$  are any real numbers and both  $a$  and  $b$  can be zero.

### Key Fact:

- The numbers  $2+5i$ ,  $7-11i$ ,  $18i$  are imaginary as  $b \neq 0$  in these numbers. But they are also complex numbers.
- Complex number like  $5i$  in which  $a=0$  and  $b \neq 0$  are imaginary number with no real part. Such numbers are called pure imaginary numbers.
- As 6 can be written as:  $6+0i$ . Therefore, every real number is a complex number with 0 as imaginary part.
- The set of complex numbers is denoted by  $C$  and  $R \subseteq C$ .



## Addition and Subtraction of Complex Numbers

The complex numbers obey the commutative, associative and distributive law. Thus, we can add and subtract them as we do binomials. Moreover, the sum and difference of two complex numbers is also a complex number. If  $z_1 = a + bi$  and  $z_2 = c + di$  are two complex numbers then their sum and difference is defined as:

**Sum:** Add real part in real part and imaginary part in imaginary part.

$$z_1 + z_2 = (a + bi) + (c + di) = (a + c) + (b + d)i$$

**Difference:** Subtract real part in real part and imaginary part in imaginary part.

$$z_1 - z_2 = (a + bi) - (c + di) = (a - c) + (b - d)i$$

**Example:** Simplify:

a.  $(7 + 5i) + (4 + 3i)$

b.  $(5 + 6i) - (7 - 4i)$

**Solution:**

a.  $(7 + 5i) + (4 + 3i)$

$$= 7 + 5i + 4 + 3i$$

= 7 + 4 + 5i + 3i (combine real and

= 11 + 8i      iminiary parts)

b.  $(5 + 6i) - (7 - 4i)$

$$= 5 + 6i - 7 + 4i$$

= 5 - 7 + 6i + 4i (combining real and

= -2 + 10i      iminiary parts)

## Multiplication of Complex Numbers

If  $a + bi$  and  $c + di$  are two complex numbers then their product can be found by using distributive law.

$$(a + bi)(c + di) = a(c + di) + bi(c + di) = ac + adi + bci + bdi^2 \\ = ac + adi + bci - bd = (ac - bd) + (ad + bc)i \quad \dots \dots \text{ (as } i^2 = -1\text{)}$$

Therefore, the product of two complex numbers is also a complex number.

For complex numbers, the property  $\sqrt{a}\sqrt{b} = \sqrt{ab}$  does not hold in general, but it does when  $a$  is negative and  $b$  is a non-negative number.

To express square root of negative numbers, we first express them in terms of  $i$ . For example:

$$\sqrt{-3} \times \sqrt{-7} = \sqrt{-1}\sqrt{3} \times \sqrt{-1}\sqrt{7} = i\sqrt{3} \times i\sqrt{7} = i^2\sqrt{21} = -\sqrt{21} : \text{correct.}$$

But,  $\sqrt{-2} \times \sqrt{-5} = \sqrt{-(2)(-5)} = \sqrt{10}$  is wrong.

**Example:** Multiply and simplify:

a.  $\sqrt{-2}\sqrt{-11}$

b.  $-2i \cdot 5i$

c.  $-3i(2 - 8i)$

d.  $\sqrt{-9}\sqrt{-36}$

e.  $(3 - 2i)(1 + 5i)$

f.  $(5 - i)(9 - 3i)$

**Solution:**

a.  $\sqrt{-2}\sqrt{-11} = \sqrt{-1}\sqrt{2} \times \sqrt{-1}\sqrt{11} = i\sqrt{2} \times i\sqrt{11} = i^2\sqrt{2 \times 11} = -\sqrt{22}$

b.  $-2i \cdot 5i = -10(i^2) = -10(-1) = 10 \quad (\text{as } i^2 = -1)$

- c.  $-3i(2 - 8i) = -6i + 24(i^2) = -6i + 24(-1) = -6i - 24 = -24 - 6i$   
d.  $\sqrt{-9}\sqrt{-36} = \sqrt{-1 \times 9}\sqrt{-1 \times 36} = \sqrt{-1}\sqrt{9} \times \sqrt{-1}\sqrt{36} = i3 \times i6 = 18i^2 = -18$   
e.  $(3 - 2i)(1 + 5i) = 3 + 15i - 2i - 10i^2 = 3 + 13i - 10(-1) = 13 + 13i$   
f.  $(5 - i)(9 - 3i) = 45 - 15i - 9i + 3i^2 = 45 - 24i + 3(-1) = 42 - 24i$

### Powers of $i$

We now want to simplify certain expressions involving higher powers of  $i$ . Simplifying powers of  $i$  can then be done by using the fact that  $i^2 = -1$  and expressing the given power of  $i$  in terms of  $i^2$ . Consider:

$$\begin{aligned}i^3 &= i^2 \cdot i = (-1) \cdot i = -i \\i^5 &= i^4 \cdot i = (i^2)^2 \cdot i = (-1)^2 \cdot i = i \\i^7 &= i^6 \cdot i = (i^2)^3 \cdot i = (-1)^3 \cdot i = -i\end{aligned}$$

$$\begin{aligned}i^4 &= (i^2)^2 = (-1)^2 = 1 \\i^6 &= (i^2)^3 = (-1)^3 = -1 \\i^8 &= (i^2)^4 = (-1)^4 = 1\end{aligned}$$

The powers of  $i$  cycle themselves through the values of  $i, -1, -i$  and  $1$ .

**Example:** Simplify:

a.  $i^{11}$

b.  $i^{22}$

c.  $i^{61}$

d.  $i^{100}$

**Solution:**

$$\begin{aligned}a. \quad i^{11} &= (i^2)^5 \cdot i = (-1)^5 \cdot i = (-1) \cdot i = -i \\b. \quad i^{22} &= (i^2)^{11} = (-1)^{11} = -1 \\c. \quad i^{61} &= (i^2)^{30} \cdot i = (-1)^{30} \cdot i = (1) \cdot i = i \\d. \quad i^{100} &= (i^2)^{50} = (-1)^{50} = 1\end{aligned}$$

### Key Fact:

Imaginary numbers appear in the field of engineering and physical sciences.

**Example:** Write the expressions in the form of  $a + ib$ .

a.  $2 + i^3$

b.  $19 + 7i^2$

c.  $i^{12} + i^{15}$

d.  $4i^3 + 8i^6$

**Solution:**

a.  $2 + i^3 = 2 + i^2 \cdot i = 2 + (-1) \cdot i = 2 - i$

b.  $19 + 7i^2 = 19 + 7(-1) = 19 - 7 = 12 + 0i$

c.  $i^{12} + i^{15} = (i^2)^6 + (i^2)^7 \cdot i = (-1)^6 + (-1)^7 \cdot i = 1 + (-1) \cdot i = 1 - i$

d.  $4i^3 + 8i^6 = 4i^2 \cdot i + 8(i^2)^3 = 4(-1) \cdot i + 8(-1)^3 = -4i + 8(-1) = -4i - 8 = -8 - 4i$

### Conjugates of Complex Numbers

Conjugate of complex number is a complex number obtained by changing the sign of the imaginary part. In simple words, conjugate of a complex number is a number that has the same real part as the original complex number, and the imaginary part has the same magnitude but opposite sign.

The conjugate of a complex number  $a + ib$  is  $a - ib$ .

For example, the conjugate of  $5 + 6i$  is  $5 - 6i$  and that of  $3 - 11i$  is  $3 + 11i$ .

Commonly, the complex number is denoted by  $z$  and its conjugate by  $\bar{z}$ .

For example, if  $z = 11 + 7i$  then  $\bar{z} = \overline{11 + 7i} = 11 - 7i$

**Example:** Find the conjugate:

a.  $5+9i$

b.  $13-5i$

c.  $-5-7i$

d.  $11i$

**Solution:**

a.  $5+9i$  the conjugate is  $5-9i$

b.  $13-5i$  the conjugate is  $13+5i$

c.  $-5-7i$  the conjugate is  $-5+7i$

d.  $11i$  the conjugate is  $-11i$

## Division of Complex Numbers

Dividing complex numbers is a little more complicated than addition, subtraction, and multiplication of complex numbers because it is difficult to divide a number by an imaginary number.

For dividing complex numbers, we need to find a term by which we can multiply the numerator and the denominator that will eliminate the imaginary part of the denominator so that we end up with a real number in the denominator.

If  $z_1 = x_1 + iy_1$  and  $z_2 = x_2 + iy_2$  are the two complex numbers, then division of  $z_1$  by  $z_2$  is written as:

$$\frac{z_1}{z_2} = \frac{x_1+iy_1}{x_2+iy_2} \quad \dots \dots \text{(i)}$$

To eliminate the imaginary part in the denominator, we multiply the numerator and denominator of (i) by the conjugate of  $x_2 + iy_2$  i.e. by  $x_2 - iy_2$

$$\frac{z_1}{z_2} = \frac{x_1+iy_1}{x_2+iy_2} \times \frac{x_2-iy_2}{x_2-iy_2} \quad \dots \dots \text{(ii)}$$

After that, we simplify the expression (ii).

## Steps for Dividing Complex Numbers

To divide the two complex numbers, follow these steps:

- First, calculate the conjugate of the complex number that is at the denominator of the fraction.
- Multiply the conjugate with the numerator and the denominator of the complex fraction.
- Apply the algebraic identity  $(a+b)(a-b) = a^2 - b^2$  in the denominator and substitute  $i^2 = -1$ .
- Apply the distributive property in the numerator and simplify.
- Separate the real part and the imaginary part of the resultant complex number.

**Example:** Divide and simplify to the form  $a+ib$ .

a.  $\frac{-7+3i}{2-5i}$

b.  $\frac{11+5i}{10i}$

c.  $\frac{1-7i}{3+4i}$

**Solution:**

a.  $\frac{-7+3i}{2-5i}$

$$= \frac{-7+3i}{2-5i} \times \frac{2+5i}{2+5i} \quad \text{multiplying and dividing by conjugate of denominator}$$

$$\begin{aligned}
 &= \frac{(-7+3i)(2+5i)}{(2-5i)(2+5i)} = \frac{-14-35i+6i+15i^2}{(2)^2-(5i)^2} \\
 &= \frac{-14-35i+6i+15(-1)}{4-25i^2} = \frac{-14-29i-15}{4-25(-1)} \\
 &= \frac{-29-29i}{4+25} = \frac{-29-29i}{29} = \frac{29(-1-i)}{29}
 \end{aligned}$$

$$=-1-i$$

b.  $\frac{11+5i}{10i}$

$$\frac{11+5i}{10i} \times \frac{-10i}{-10i} \text{ multiplying and dividing by conjugate of denominator } (10i = 0+10i)$$

$$= \frac{(11+5i)(-10i)}{(10i)(-10i)} = \frac{-110i-50i^2}{-100i^2} = \frac{-110i-50(-1)}{-100(-1)}$$

$$= \frac{-110i+50}{100} = \frac{50-110i}{100} = \frac{50}{100} - \frac{110}{100}i$$

$$= \frac{1}{2} - \frac{11}{10}i$$

c.  $\frac{1-7i}{3+4i}$

$$= \frac{1-7i}{3+4i} \times \frac{3-4i}{3-4i} \text{ multiplying and dividing by conjugate of denominator}$$

$$= \frac{(1-7i)(3-4i)}{(3+4i)(3-4i)} = \frac{3-4i-21i+28i^2}{(3)^2-(4i)^2} = \frac{3-25i+28(-1)}{9-16i^2}$$

$$= \frac{3-25i-28}{9-16(-1)} = \frac{-25-25i}{9+16} = \frac{-25-25i}{25} = \frac{25(-1-i)}{25}$$

$$=-1-i.$$

### Exercise 1.1

1. Simplify and express in terms of  $iy$ .

i.  $\sqrt{-3}$

ii.  $6\sqrt{-4}$

iii.  $\sqrt{-\frac{4}{9}}$

iv.  $-\sqrt{-20}$

v.  $4-\sqrt{-60}$

vi.  $\sqrt{-8}\sqrt{-2}$

2. Simplify:

i.  $(4-i)+(5+5i)$

ii.  $(7-6i)-(5-6i)$

iii.  $(-2+8i)-(7+3i)$

iv.  $(4-2i)-(5-2i)$

v.  $(2+4i)(1+2i)$

vi.  $(1-4i)(2-3i)$

vii.  $-8i(2-2i)$

viii.  $(3+2i)^2$

ix.  $(3-6i)(3+6i)$

x.  $(-5-3i)^2$

xi.  $(1+\sqrt{2}i)(1-\sqrt{3}i)$

xii.  $(\sqrt{2}+i)(\sqrt{2}-i)$

3. Simplify:

i.  $i^9$

ii.  $i^{13}$

iii.  $i^{28}$

iv.  $(-i)^{21}$

v.  $(3i)^3$

vi.  $(-2i)^4$

4. Simplify in the form of  $a+bi$ .

i.  $9+i^6$

ii.  $-17+i^5$

iii.  $i^4 - 13i$

iv.  $i^5 + 21i$

v.  $i^5 + i^7$

vi.  $i^{74} - i^{100}$

5. Divide and simplify in the form of  $a+bi$ .

i.  $\frac{3}{4-i}$

ii.  $\frac{3i}{6+5i}$

iii.  $\frac{3-i\sqrt{5}}{3+i\sqrt{5}}$

iv.  $\frac{2+7i}{5i}$

v.  $\frac{4+5i}{4-5i}$

vi.  $\frac{3+2i}{2+i}$

vii.  $\frac{5+i}{1+2i}$

viii.  $\frac{a+ib}{a-ib}$

ix.  $\frac{1+i}{(1-i)^2}$

x.  $\frac{(2+2i)^2}{(1+i)^2}$

### Developing skilled knowledge

- ✓ What is an imaginary number?
- ✓ Is an imaginary number real number?
- ✓ Provide examples of numbers that are pure imaginary numbers.
- ✓ What is the pattern formed by the powers of  $i$ ?
- ✓ Can we say product of  $(2-3i)(2-3i)$  is imaginary?
- ✓ If conjugate of  $z$  is  $\bar{z}$ , then is  $z\bar{z}$  a real number?

### Properties of Complex Numbers

Let  $z = a + bi$ ,  $w = c + di$  and  $v = e + fi$  be any three complex numbers, then the following properties hold.

- $z + w = w + z$  (commutative law of addition)
- $z \times w = w \times z$  (commutative law of multiplication)
- $z + 0 = 0 + z = z$  (existence of additive identity where  $0 = 0 + 0i$ )
- $z \times 1 = 1 \times z = z$  (existence of multiplicative identity where  $1 = 1 + 0i$ )
- $(z + w) + v = z + (w + v)$  (associative law of addition)
- $(z \times w) \times v = z \times (w \times v)$  (associative law of multiplication)
- $z(w + v) = zw + zv$  (distributive law of multiplication over addition)
- $z(w - v) = zw - zv$

#### Challenge

Prove the above properties by taking,  $z = 1 + 2i$ ,  $w = 3 - i$  and  $v = 2 + 2i$

## Additive Inverse of Complex Numbers

If  $z = a + bi$  is a complex number then its additive inverse is  $-z = -a - bi$ .

In the same way, additive inverse of  $-z = -a - bi$  is  $z = a + bi$ .

Sum of a complex number and its additive inverse is zero.

$$z + (-z) = (a + bi) + (-a - bi) = a + bi - a - bi = 0 = 0 + 0i$$

### Example:

Find the additive inverse of  $3 - 2i$ .

### Solution:

Let  $a + bi$  be the additive inverse of  $3 - 2i$ .

Then by additive inverse rule:

$$(3 - 2i) + (a + bi) = 0 + 0i$$

$$(3 + a) + (-2i + bi) = 0 + 0i$$

$$(3 + a) + (-2 + b)i = 0 + 0i$$

Comparing real and imaginary parts of both sides, we get:

$$3 + a = 0; -2 + b = 0 \Rightarrow a = -3; b = 2$$

Thus, additive inverse of  $3 - 2i$  is  $-3 + 2i$ .

## Equality of Complex Numbers

If two complex numbers  $a + bi$  and  $c + di$  are equal then:

$$a = c \text{ and } b = d$$

### Direct Method:

We can apply the rule that additive inverse is the opposite sign of  $3 - 2i$  that is  $-3 + 2i$ .

Similarly the additive inverse of  $-7 + 11i$  is  $7 - 11i$ .

## Multiplicative Inverse of Complex Numbers

The multiplicative inverse of  $a + bi$  is  $\frac{1}{a+bi}$ .

### Example: Find the multiplicative inverse of $7 - 5i$ .

**Solution:** The multiplicative inverse of  $7 - 5i$  is  $\frac{1}{7-5i}$ .

Now, simplifying the expression  $\frac{1}{7-5i} = \frac{1}{7-5i} \times \frac{7+5i}{7+5i}$  multiplying by conjugate

$$= \frac{7+5i}{(7)^2 - (5i)^2} \Rightarrow \frac{7+5i}{49 - 25i^2} \Rightarrow \frac{7+5i}{49 - 25(-1)}$$

$$= \frac{7+5i}{49+25} = \frac{7+5i}{84} = \frac{7}{84} + \frac{5}{84}i$$

$$= \frac{1}{12} + \frac{5}{84}i$$

Verification:  $(7 - 5i) \times \left(\frac{1}{12} + \frac{5}{84}i\right) = (7 - 5i) \left(\frac{7+5i}{84}\right)$

$$= \frac{(7)^2 - (5i)^2}{84} = \frac{49 - 25i^2}{84} = \frac{49 - 25(-1)}{84}$$

$$= \frac{49+25}{84} = \frac{84}{84} = 1$$

### Key Fact:

The product of a complex number and its multiplicative inverse is 1.

## Properties of Conjugate of Complex Number

### Key Fact:

a.  $z\bar{z}$  is a real number.

$$\begin{aligned} \text{Consider } z = 7 - 6i \text{ then } \bar{z} &= \overline{7 - 6i} = 7 + 6i \\ z\bar{z} &= (7 - 6i)(7 + 6i) = (7)^2 - (6i)^2 = 49 - 36i^2 \\ &= 49 - 36(-1) = 49 + 36 = 85 \end{aligned}$$

The complex number  $a + ib$  can be written as  $(a, b)$ ,  $a$  is a real part and  $b$  is imaginary part.

b.  $\overline{z_1 z_2} = \bar{z}_1 \cdot \bar{z}_2$

Consider  $z_1 = 1 - 2i$  and  $z_2 = 4 + 3i$ , then:

$$\bar{z}_1 = 1 + 2i \text{ and } \bar{z}_2 = 4 - 3i$$

$$\begin{aligned} \text{Now, LHS} &= z_1 \cdot z_2 = (1 - 2i)(4 + 3i) = 4 + 3i - 8i - 6i^2 \\ &= 4 - 5i - 6(-1) = 4 - 5i + 6 = 10 - 5i \end{aligned}$$

$$\overline{z_1 \cdot z_2} = \overline{10 - 5i} = 10 + 5i$$

$$\begin{aligned} \text{RHS} &= \bar{z}_1 \cdot \bar{z}_2 = (1 + 2i)(4 - 3i) = 4 - 3i + 8i - 6i^2 \\ &= 4 + 5i - 6(-1) = 4 + 5i + 6 = 10 + 5i \end{aligned}$$

Hence,  $\overline{z_1 \cdot z_2} = \bar{z}_1 \cdot \bar{z}_2$

c.  $\left(\frac{z_1}{z_2}\right) = \frac{\bar{z}_1}{\bar{z}_2}$

Consider  $z_1 = -1 + 5i$  and  $z_2 = 2 + i$

Then,  $\bar{z}_1 = -1 - 5i$ ,  $\bar{z}_2 = 2 - i$

$$\begin{aligned} \text{Taking } \frac{z_1}{z_2} &= \frac{-1+5i}{2+i} = \frac{-1+5i}{2+i} \times \frac{2-i}{2-i} = \frac{(-1+5i)(2-i)}{(2+i)(2-i)} \\ &= \frac{-2+i+10i-5i^2}{(2)^2-(i)^2} = \frac{-2+11i-5(-1)}{4-(-1)} = \frac{-2+11i+5}{4+1} = \frac{3+11i}{5} = \frac{3}{5} + \frac{11}{5}i \end{aligned}$$


$$\text{LHS} = \left(\frac{z_1}{z_2}\right) = \frac{3}{5} - \frac{11}{5}i$$

$$\begin{aligned} \text{RHS} &= \frac{\bar{z}_1}{\bar{z}_2} = \frac{-1-5i}{2-i} = \frac{-1-5i}{2-i} \times \frac{2+i}{2+i} = \frac{(-1-5i)(2+i)}{(2-i)(2+i)} \\ &= \frac{-2-i-10i-5i^2}{(2)^2-(i)^2} = \frac{-2-11i-5(-1)}{4-(-1)} = \frac{-2-11i+5}{4+1} = \frac{3-11i}{5} = \frac{3}{5} - \frac{11}{5}i \end{aligned}$$

Hence,  $\left(\frac{z_1}{z_2}\right) = \frac{\bar{z}_1}{\bar{z}_2}$

## Graphical Representation of Complex Number


In complex number  $z = a + ib$ , there are two parts of  $z$ ,  $a$  is a real part and  $b$  is an imaginary part associated with  $i$ . On the graphical representation of complex number  $z$ , the real part of  $z$  is taken along  $x$ -axis and imaginary part is taken along  $y$ -axis and considered as to real axis and imaginary axis respectively. The coordinate plane itself is called the complex plane or  $z$ -plane.



**Example:** Represent the numbers in the complex plane.

$$2 + 3i, 3 - 2i, -3 + i, -2 - 3i$$

**Solution:**



We observe that the complex numbers appear in all the four quadrants due to the negative and positive signs of their real and imaginary parts.

**Key Fact:**

The complex plane is also called "The Argand Diagram" after the French-Swiss mathematician Jean Robert Argand (1768-1822) who was the first to represent a complex number geometrically as point in the plane.

**Key Fact:**

If  $x = 0$ , then  $x + yi = yi$  is an imaginary number located on vertical axis.

If  $y = 0$ , then  $x + yi = x$  is a real number located on horizontal axis.

**Absolute Value or Modulus of a Complex Number**


Let  $z = (a, b) = a + ib$  be a complex number.

The absolute value or modulus of  $z$ , denoted by  $|z|$  is defined as:

$$|z| = \sqrt{a^2 + b^2}$$

which is a real number. From the diagram, it is clear that the modulus of a complex number is the distance OP of

complex number from the origin.



**Key Fact:**

If,  $z = a + ib$ , then  $|z| = \sqrt{(\text{real part})^2 + (\text{img part})^2} = \text{Real Number}$

**Example:** Find the absolute value of the complex numbers:

a.  $5i$

b.  $2$

c.  $4 - 7i$

d.  $-3 + 2i$

**Solution:**

a.  $z = 5i = 0 + 5i = |z| = \sqrt{0^2 + 5^2} = \sqrt{25} = 5$

b.  $z = 2 = 2 + 0i = |z| = \sqrt{2^2 + 0^2} = \sqrt{4} = 2$

c.  $z = 4 - 7i = |z| = \sqrt{(4)^2 + (-7)^2} = \sqrt{16 + 49} = \sqrt{65}$

d.  $z = -3 + 2i = |z| = \sqrt{(-3)^2 + (2)^2} = \sqrt{9 + 4} = \sqrt{13}$

### Properties of Modulus

If  $z$  is a complex number, then following properties holds:

a.  $|z| = |-z| = |\bar{z}| = |-\bar{z}|$       b.  $\bar{z} \cdot z = |z|^2$

a. Consider a complex number  $z = 2 - 5i$

$$|z| = \sqrt{(2)^2 + (-5)^2} = \sqrt{4 + 25} = \sqrt{29} \dots \dots \text{(i)}$$

$$-z = -(2 - 5i) = -2 + 5i$$

$$|-z| = \sqrt{(-2)^2 + (5)^2} = \sqrt{4 + 25} = \sqrt{29} \dots \dots \text{(ii)}$$

$$-\bar{z} = -2 - 5i$$

$$|\bar{z}| = \sqrt{2^2 + 5^2} = \sqrt{4 + 25} = \sqrt{29} \dots \dots \dots \text{(iii)}$$

$$\bar{z} = \overline{2 - 5i} = 2 + 5i$$

$$|-\bar{z}| = \sqrt{(-2)^2 + (-5)^2} = \sqrt{4 + 25} = \sqrt{29} \dots \text{(iv)}$$

From (i), (ii), (iii), and (iv):  $|z| = |-z| = |\bar{z}| = |-\bar{z}|$

b. Consider  $z = 2 - 5i$

$$\bar{z} = 2 + 5i \text{ then,}$$

$$\begin{aligned}\bar{z} \cdot z &= (2 + 5i)(2 - 5i) \\ &= (2)^2 - (5i)^2 = 4 - 25i^2 \\ &= 4 - 25(-1) = 4 + 25 \\ &= 29 \dots \dots \text{(i)}\end{aligned}$$

$$\begin{aligned}|z| &= \sqrt{(2)^2 + (-5)^2} \\ &= \sqrt{4 + 25} = \sqrt{29}\end{aligned}$$

$$|z|^2 = (\sqrt{29})^2 = 29 \dots \dots \text{(ii)}$$

From (i) and (ii),

$$\bar{z} \cdot z = |z|^2$$

**Example:** Find the real and imaginary parts of the complex numbers:

a.  $(2-3i)^2$

b.  $(1+2i)^{-2}$

c.  $\frac{(\sqrt{2}+i\sqrt{3})^2}{1+i}$

d.  $\frac{(1-i)^2}{(1+i)^2}$

**Solution:**

$$\begin{aligned}\text{a. } (2-3i)^2 &= (2)^2 + (3i)^2 - 12i = 4 + 9i^2 - 12i = 4 + 9(-1) - 12i \\ &= 4 - 9 - 12i = -5 - 12i. \text{ Real part} = -5, \text{ Imaginary part} = -12\end{aligned}$$

$$\begin{aligned}\text{b. } (1+2i)^{-2} &= \frac{1}{(1+2i)^2} = \frac{1}{(1)^2 + (2i)^2 + 4i} = \frac{1}{1 + 4i^2 + 4i} = \frac{1}{1 + 4(-1) + 4i} \\ &= \frac{1}{1-4+4i} = \frac{1}{-3+4i} = \frac{1}{-3+4i} \times \frac{-3-4i}{-3-4i} \\ &= \frac{-3-4i}{(-3)^2 - (4i)^2} = \frac{-3-4i}{9-16i^2} = \frac{-3-4i}{9-16(-1)} \\ &= \frac{-3-4i}{9+16} = \frac{-3-4i}{25} = \frac{-3}{25} - \frac{4}{25}i. \quad \text{Real part} = \frac{-3}{25}, \quad \text{Imaginary part} = \frac{-4}{25}\end{aligned}$$

c. 
$$\begin{aligned} \frac{(\sqrt{2}+i\sqrt{3})^2}{1+i} &= \frac{(\sqrt{2})^2 + (i\sqrt{3})^2 + 2i\sqrt{2}\sqrt{3}}{1+i} = \frac{2+3i^2+2i\sqrt{3}\times 2}{1+i} \\ &= \frac{2+3(-1)+2i\sqrt{6}}{1+i} = \frac{2-3+2i\sqrt{6}}{1+i} = \frac{-1+2i\sqrt{6}}{1+i} \\ &= \frac{-1+2i\sqrt{6}}{1+i} \times \frac{1-i}{1-i} = \frac{(-1+2i\sqrt{6})(1-i)}{(1)^2-(i)^2} = \frac{-1+2i\sqrt{6}+i-2i^2\sqrt{6}}{1-i^2} \\ &= \frac{-1+(2\sqrt{6}+1)i+2\sqrt{6}}{1-(-1)} = \frac{-1+2\sqrt{6}+(2\sqrt{6}+1)i}{2} = \frac{-1+2\sqrt{6}}{2} + \frac{(2\sqrt{6}+1)}{2}i \end{aligned}$$

Real part =  $\frac{-1+2\sqrt{6}}{2}$ , Imaginary part =  $\frac{(2\sqrt{6}+1)}{2}$

d. 
$$\frac{(1-i)^2}{(1+i)^2} = \frac{(1)^2 + (i)^2 - 2i}{(1)^2 + (i)^2 + 2i} = \frac{1-1-2i}{1-1+2i} = \frac{-2i}{2i} = -1$$

Real part = -1, Imaginary part = 0


## Geometrical Representation of Algebraic Operations

The graphical representation of algebraic operations in the complex number system makes it easier to understand the concepts.

### (i) Geometrical Representation of Addition

Geometrically, addition of two complex numbers  $z_1$  and  $z_2$  can be visualized as addition of the vectors by using the *parallelogram law*. The vector sum  $z_1 + z_2$  is represented by the diagonal of the parallelogram formed by the two original vectors.

If two points P and Q represent complex numbers  $z_1$  and  $z_2$  respectively in the Argand Plane, then the sum  $z_1 + z_2$  is represented by the point R of the diagonal OR of parallelogram OPRQ having OP and OQ as two adjacent sides.




### (ii) Geometrical Representation of Subtraction

The representation of the difference of two complex numbers, is slightly complicated than the addition of the complex numbers. The easiest way of presenting the subtraction is to think of addition of a negative vector.

If we want to represent  $z_1 - z_2$ , then the easiest way of representing it would be to think of adding a negative vector  $z_1 + (-z_2)$ .

The negative vector is the same as the positive one, the only difference being that the negative vector points in the opposite direction.

The difference vector  $z_1 - z_2$  is represented in the adjoining figure.



#### Key Facts:

It is important to note here that the vector representing the difference of the vectors  $z_1 - z_2$  may also be drawn joining the end point of  $z_2$  to the tip of  $z_1$  instead of the origin. This kind of representation does not alter the meaning or interpretation of the difference operator.

#### Check Point:

Can you think about a geometric interpretation of the addition of three or more complex numbers?

#### Example:



Represent sum and difference of  $z_1 = 4 + 2i$  and  $z_2 = 2 + 3i$  graphically.

#### Solution:

Given,  $z_1 = 4 + 2i = (4, 2)$  and  $z_2 = 2 + 3i = (2, 3)$

$$\text{Sum} = z_1 + z_2 = (4 + 2i) + (2 + 3i) = 6 + 5i = (6, 5)$$

$$\text{Difference} = z_1 - z_2 = (4 + 2i) - (2 + 3i) = 2 - i = (2, -1)$$




### (iii) Geometrical Representation of Multiplication and Division

Consider,  $z_1 = 4 + 3i$  and  $z_2 = 2 - 2i$ .

$$\begin{aligned} \text{Now, } z_1 z_2 &= (4 + 3i)(2 - 2i) \\ &= 8 - 8i + 6i - 6i^2 \\ &= 8 - 2i + 6 = 14 - 2i \\ &= (14, -2) \end{aligned}$$


Product  $z_1 z_2$  is shown in the adjoining diagram.



Again, consider  $z_1 = 4 + 6i$   
and  $z_2 = -2i$  then:

$$\begin{aligned} z_1 \div z_2 &= (4 + 6i) \div (-2i) \\ &= -3 + 2i \end{aligned}$$

The result is shown in the diagram.



#### Exercise 1.2

- Find the additive inverse of each complex number.
  - $-4+5i$
  - $-3-3i$
  - $5-5i$
  - $4i$
- Show that each pair of complex numbers are multiplicative inverse of each other.
  - $2+3i, \frac{2-3i}{13}$
  - $5-4i, \frac{5+4i}{41}$
  - $6+8i, \frac{3-4i}{50}$
- Find the multiplicative inverse of each complex number.
  - $1+i$
  - $7-3i$
  - $10-12i$
  - $\frac{2}{5-i}$
  - $\frac{-i}{2-3i}$
  - $a-bi$
- Find the product of each complex number and its conjugate.
  - $4$
  - $1-i$
  - $7i$
  - $6-2i$
  - $10+9i$
  - $-4-11i$
- If  $z_1 = 1 - 2i$  and  $z_2 = 2 + i$ :
  - Show that:
    - $\overline{z_1 \cdot z_2} = \overline{z_1} \cdot \overline{z_2}$
    - $\overline{\left(\frac{z_1}{z_2}\right)} = \frac{\overline{z_1}}{\overline{z_2}}$

$$\text{iii. } |z_1| = |-z_1| = |\bar{z}_1| = |-\bar{z}_1| \quad \text{iv. } z_2 \cdot \bar{z}_2 = |z_2|^2$$

b. Find:

$$\text{i. } |z_1 + z_2|$$

$$\text{ii. } |z_1 z_2|$$

$$\text{iii. } \left| \frac{z_1}{z_2} \right|$$

6. Represent the numbers in the complex plane.

$$\text{a. } -1-3i$$

$$\text{b. } 2+4i$$

$$\text{c. } -3+2i$$

$$\text{d. } 2-3i$$

$$\text{e. } 2i$$

$$\text{f. } -3i$$

$$\text{g. } 2$$

7. Separate into real and imaginary parts of each complex number.

$$\text{a. } (\sqrt{2}-\sqrt{3}i)^2$$

$$\text{b. } (\sqrt{2}+i)^2$$

$$\text{c. } \frac{(2+3i)^2}{1-3i}$$

$$\text{d. } \left[ \frac{1}{2} + \frac{\sqrt{3}}{2}i \right]^2$$

$$\text{e. } \frac{1-i}{(i)^2}$$

$$\text{f. } \frac{1}{i(1-i)^2}$$

$$\text{g. } \frac{(1+i)^2}{(1-2i)^2}$$

8. Taking any complex number and show that:

$$\text{a. } z \cdot \bar{z} \text{ is a real number.}$$

$$\text{b. } z^2 + (\bar{z})^2 \text{ is a real number.}$$

$$\text{c. } (z - \bar{z})^2 \text{ is a real number.}$$

$$\text{d. } |z| \text{ and } |\bar{z}| \text{ are real numbers.}$$

$$\text{e. } z^2 - (\bar{z})^2 \text{ is an imaginary number.}$$

9. Represent sum and difference of complex numbers graphically.

$$\text{(i) } z_1 = 5 + 3i \text{ and } z_2 = 2 - 3i \quad \text{(ii) } z_1 = -3 + 2i \text{ and } z_2 = 4 + 3i$$

10. Represent product of complex numbers graphically.

$$\text{(i) } z_1 = 4 + 2i \text{ and } z_2 = -2 + 3i \quad \text{(ii) } z_1 = -2 + 4i \text{ and } z_2 = 3 - i$$

11. Represent  $z_1 \div z_2$  graphically when:

$$\text{(i) } z_1 = 6 - 4i \text{ and } z_2 = 3 \quad \text{(ii) } z_1 = -4 - 6i \text{ and } z_2 = 1 + i$$

## Applications of Complex Numbers

Complex numbers are used in many fields in real life. Major Fields where complex numbers are used are:

Electronics

Electromagnetism

Computer science engineering

Civil engineering

Mechanical engineering

Control systems .

In modelling and problem solving, we find many equations whose roots are complex numbers. Dealing such kind of problems, we must have conceptual understanding of complex numbers. Practically, negative numbers do not have square roots in the number system. However, a large number system that contains the real number system is designed so that negative numbers do have square roots.

**Example:** Solve  $x^2 + 5 = 0$

**Solution:**  $x^2 + 5 = 0$

$$x^2 = -5$$

$$x = \pm\sqrt{-5} = \pm\sqrt{-1 \times 5} = \pm\sqrt{-1} \times \sqrt{5}$$
$$= \pm\sqrt{5}i$$

The solution  $+\sqrt{5}i$  and  $-\sqrt{5}i$  are both pure imaginary numbers.

Sometimes we need factorization of the equation, but in real number system in some expressions making factors are not possible. By using the complex numbers properties, we can make factorization possible.

**Example:** Factorize:

a.  $z^2 + 9$

b.  $3z^2 + 5$

**Solution:**

a.  $z^2 + 9$

b.  $3z^2 + 5$

$$= (z)^2 - 9i^2$$

$$= (\sqrt{3})^2 z^2 - i^2 (\sqrt{5})^2$$

$$= (z)^2 - (3i)^2$$

$$= (\sqrt{3}z)^2 - (i\sqrt{5})^2$$

$$= (z - 3i)(z + 3i)$$

$$= (\sqrt{3}z - i\sqrt{5})(\sqrt{3}z + i\sqrt{5})$$

The factors of  $z^2 + 9$  are  $(z - 3i)(z + 3i)$

The factors are  $(\sqrt{3}z - i\sqrt{5})(\sqrt{3}z + i\sqrt{5})$

**Example:** Mr. Waqas is an electrical engineer designing the electrical circuits for a new office building. There are three basic things to be considered in an electrical circuit: The flow of the electric current I, the resistance to that flow Z, called impedance, and electromotive force E, called voltage. Their quantities are related in the formula  $E = IZ$ . The current of the circuit Waqas is designing is to be  $(35 - 40J)$  amp. Electrical engineers use the letter J to represent the imaginary unit. Find the impedance of the circuit if the voltage is to be  $(430 - 330J)$  volts.

**Solution:** We know that  $E = IZ$  .....(i)

We know it is given that  $I = 35 - 40J$ ,  $E = 430 - 330J$

Substituting the values in (i)  $E = IZ$

and  $Z = \frac{E}{I}$

$$Z = \frac{430 - 330J}{35 - 40J}$$

$$= \frac{430 - 330J}{35 - 40J} \times \frac{35 + 40J}{35 + 40J}$$

$$= \frac{15050 + 12200J - 11550J - 13200J^2}{1225 - 1600J^2}, \quad J^2 = -1$$

**Check Point:**

$$x^2 + 5 = 0$$

Solution is  $\pm\sqrt{5}i$ .

Verify the solution.

$$= \frac{2825 + 5650J}{2825} = 1 + 2J$$

The impedance will be  $1 + 2J$  ohms.

**Example:** If  $z_1 = 5.5 + 5i$ ,  $z_2 = 7.7 + 7i$ , then calculate:

- a.  $z_1 + z_2$       b.  $z_1 z_2$       c.  $\frac{z_1}{z_2}$

**Solution:**

$$a. \quad z_1 + z_2 = 5.5 + 5i + 7.7 + 7i = 13.2 + 12i$$

$$b. \quad z_1 z_2 = (5.5 + 5i)(7.7 + 7i) = 42.35 + 38.5i + 38.5i + 35i^2 \\ \equiv 7.35 + 77i$$

$$c. \frac{z_1}{z_2} = \frac{5.5+5i}{7.7+7i} = \frac{5.5+5i}{7.7+7i} \times \frac{7.7-7i}{7.7-7i} = \frac{(5.5+5i)(7.7-7i)}{(7.7)^2 - (7i)^2}$$

$$= \frac{42.35 - 38.5i + 38.5i - 35i^2}{53.9 - 49i^2} = \frac{77.35}{102.9}$$

## Solution of System of Equations with Complex Coefficients

In this section, we shall find solution of different equations in complex variables either with real or complex coefficients. Consider the following equation:

Where  $p$ ,  $q$  and  $r$  are complex numbers (every real number is a complex number). The equation (i) is called a linear equation in two complex variables  $z$  and  $w$ .

$$p_1z + q_1w = r_1$$

$$p_2 z + q_2 w = r_2$$

These two equations together form a system of linear equations in two variables  $z$  and  $w$ . The linear equations in two variables are also called simultaneous linear equations.

**Example:** Solve:

$$z + w = 3i$$

$$2z + 3w = 2$$

**Solution:** Given equations are:

Multiply equation (i) by 2 to equate the coefficient of  $z$  with equation (ii) and subtracting

$$\begin{array}{r} 2z + 2w = 6i \\ \underline{+2z + 3w = \pm 2} \\ -w = 6i - 2 \\ w = 2 - 6i \end{array}$$

(Substitute in (i) to find  $z$ )

$$\begin{aligned} z + 2 - 6i &= 3i \\ z &= 9i - 2 \end{aligned}$$

Solution is:  $z = 9i - 2$ ,  $w = 2 - 6i$

### Exercise 1.3

Solve:

1.  $x^2 + 7 = 0$       2.  $x^2 + 9 = 0$       3.  $x^2 + 100 = 0$

Determine whether the given complex number is a solution of the equation.

4.  $1+2i$ ,  $x^2 - 2x + 5 = 0$       5.  $1-2i$ ,  $x^2 - 2x + 5 = 0$

6.  $1-i$ ,  $x^2 + 2x + 2 = 0$       7.  $i$ ,  $x^2 + 1 = 0$

Factorize the expressions:

8.  $x^2 + 16$       9.  $a^2 + b^2$       10.  $x^2 + 25y^2$

Solve the following system of linear equations.

11.  $z - 4w = 3i$       12.  $3z + (2+i)w = 11 - i$   
 $2z + 3w = 11 - 5i$        $(2-i)z - w = -1 + i$

13. In an electrical circuit, the flow of the electric current  $I$ , the impedance  $Z$  and the voltage  $E$ , are related by the formula  $E = IZ$ .

a. Find  $I$ , given the values:

- $E = (70 + 220J)$  volts,  $Z = (16 + 8J)$  ohms
- $E = (85 + 110J)$  volts,  $Z = (3 - 4J)$  ohms

b. Find  $Z$  given the value:

- $E = (-50 + 100J)$  volts,  $I = (-6 - 2J)$  amp
- $E = (100 + 10J)$  volts,  $I = (-8 + 3J)$  amp

c. Evaluate  $\frac{1}{z - z^2}$  when  $z = \frac{1-i}{10}$

- Recognizing complex numbers, and conjugate and modulus of a complex numbers.
- Applying basic operations on complex numbers.
- Defining commutative laws, associative laws and distributive laws for complex numbers.
- Recognizing additive and multiplicative identity and finding additive and multiplicative inverse of complex numbers.
- Finding real and imaginary parts of  $\left(\frac{x_1+iy_1}{x_2+iy_2}\right)^n$  for  $n = \pm 1, \pm 2, \dots$
- Finding solution of equations having complex values.
- Applying the geometric interpretation of a complex numbers.
- Solving daily life problems involving complex numbers.

## MISCELLANEOUS EXERCISE-1

1. Tick the correct option.

- $\sqrt{-1}$  is equal to:
  - 1
  - 1
  - $i$
  - $-i$
- If  $x < 0$ , then  $\sqrt{x}$  is:
  - Real
  - Complex
  - Irrational
  - Rational
- Conjugate of  $\sqrt{x} - i\sqrt{y}$  is:
  - $\sqrt{x} + i\sqrt{y}$
  - $x - iy$
  - $x - y$
  - $x + iy$
- If  $z = x + iy$ , then  $\bar{z}z$  is:
  - Imaginary
  - Complex
  - Non-negative number
  - Negative number
- $\sqrt{-25} + \sqrt[3]{8}$  is equal to:
  - $-5 + \sqrt{8}$
  - $2 + 5i$
  - $-5 + 2i$
  - $2\sqrt{2} + 5i$
- $1 + (-i)^9 = ?$ 
  - $1+i$
  - $1+\sqrt{-1}$
  - $1-i$
  - $-i$
- $\frac{2}{1-i} = ?$ 
  - $\frac{1+i}{2}$
  - $\frac{(1+i)^2}{2}$
  - $1-i$
  - $1+i$
- $(-xi)^{19} = ?$ 
  - $-x^{19}i$
  - $x^{19}i$
  - $-i^{19}$
  - $-x^{19}$

- ix. If  $z = 3 + 4i$ , then  $|z|^2$  is:
- 5
  - $\sqrt{5}$
  - 25
  - 16
- x. The solution of  $x^2 + 4 = 0$  is:
- $2i$
  - $-2i$
  - $\pm 2$
  - $\pm 2i$
2. Simplify:
- $(-2 + 4i) - (8 - 5i)$
  - $(-3 + 4i) + (-7i + 4)$
3. Find the product:
- $(x + iy)(2 + 3i)$
  - $(-3 + 6i)(-6 + 3i)$
4. Write in the form of  $a + bi$ , then find its conjugate:
- $3\sqrt{2} - \sqrt{-7}$
  - $\sqrt{-2}$
5. Find  $z \cdot \bar{z}$ :
- $z = \frac{-1}{2} + i$
  - $z = 14 - 7i$
6. Simplify:
- $\frac{-3-i}{-3+i}$
  - $\frac{1+3i}{i\sqrt{5}}$
7. Factorize:
- $2x^2 + 18$
  - $-x^2 - 25y^4$
8. Solve:
- $3x^2 + 15 = 0$
  - $6y^2 + 36 = 0$

## 02

# Quadratic Equations

**After studying this unit students will be able to:**

- Define quadratic equations.
- Solve quadratic equations by graphical method, factorization and completing squares.
- Derive quadratic formula and use it to solve quadratic equations.
- Solve different types of equations which are reducible to quadratic form.
- Solve simultaneous equations. When one is homogenous and one is linear.
- Know nature of roots of a quadratic equation.
- Find relation between roots and coefficients of a quadratic equation.
- Define and evaluate symmetric functions of roots of a quadratic equation.
- Form a quadratic equation when its roots are given.

# Algebra

standard exponential expression  
inequality formula operations slope variable  
trinomial equation linear  
rational y-intercept quadratic binomial

Algebra is your trusted tool that helps you carry out of various activities of daily importance. There is hardly any line of work that does not employ the concept of algebra. We will look at the variables and equation, you surely will not be wondered about why you need to leave there in a nutshell. Algebra and equations prepare you for hardly all aspects of life and stays with you right from your infancy to your adulthood.

# Quadratic Equations

Saleem wants to build a swimming pool surrounded by a sidewalk of uniform width. He wants the dimensions of the pool and sidewalk to be 16 meters by 20 meters. The pool has an area of 192 square meters. How wide should the sidewalk be?

Let  $x$  meters be the width of the sidewalk.

The length of a pool is  $20 - 2x$  meters.

The width of the pool is  $16 - 2x$  meters.

The area of the pool can be expressed as the product of the length and width.

Area = Length  $\times$  Width

$$= (20 - 2x)(16 - 2x)$$

$$= 4x^2 - 72x + 320$$

The area of the pool can be expressed as  $4x^2 - 72x + 320$  square meters. Since the area of the pool is 192 square meters, replacing area with 192 results in the equation  $192 = 4x^2 - 72x + 320$ . Saleem would probably never use a quadratic equation like  $192 = 4x^2 - 72x + 320$  to express area of a pool. However, in this unit we will study many other formulas in science and business that involve this type of equations.

## Quadratic Equation

It is an equation that can be written in the form of  $ax^2 + bx + c = 0$ , where  $a \neq 0$ . We say that equations like this have a degree of 2. Since the greatest exponent of the variable is 2. We observe that this quadratic equation only has one variable and all the exponents are positive.

## Standard Quadratic Equation

$ax^2 + bx + c = 0$  is a standard form of the quadratic equation in one variable  $x$ . The equation  $x^2 + 6x + 5 = 0$  is in standard form.

$(x+1)^2 = 4$  and  $x^2 + 3x = 18$  are examples of quadratic equation in one variable  $x$  but they are not in standard form.

**Example:** Write the following quadratic equations in standard form.

(i).  $ax^2 + bx = c$     (ii).  $(3x - 1)^2 = 4$     (iii).  $(x + 1)^2 - (2x - 1)^2 = 2$

### Solution:

i.  $ax^2 + bx = c$

$$ax^2 + bx - c = 0$$

ii.  $(3x - 1)^2 = 4$

$$9x^2 - 6x + 1 = 4$$

$$9x^2 - 6x + 1 - 4 = 0$$

$$9x^2 - 6x - 3 = 0$$

$$3(3x^2 - 2x - 1) = 0, \text{ as } 3 \neq 0$$

$$3x^2 - 2x - 1 = 0$$

iii.  $(x + 1)^2 - (2x - 1)^2 = 2$


$$(x^2 + 2x + 1) - (4x^2 - 4x + 1) = 2$$

$$x^2 + 2x + 1 - 4x^2 + 4x - 1 - 2 = 0$$

$$-3x^2 + 6x - 2 = 0$$

$$-1(3x^2 - 6x + 2) = 0$$

$$3x^2 - 6x + 2 = 0 \quad \text{as } -1 \neq 0$$



## Solution of Quadratic Equations

The values of variables for which an equation becomes a true sentence are called solutions or roots of the equation. i.e. if  $P(a) = c$ , then  $x = a$  is root of the equation  $P(x) = c$ . The set of roots of an equation is called the solution set. A quadratic equation in one variable has two roots. These roots of the quadratic equation can be found in many ways. In this section we will study four methods.

- i. Graphing method
- ii. Factorization Method
- iii. Completing Square Method
- iv. Using Quadratic Formula

### i. Graphical Method

One way to determine the roots of a quadratic equation is to graph the related quadratic function. Write the equation in general form like:

$$192 = 4x^2 - 72x + 320$$


$0 = 4x^2 - 72x + 128$  subtract 192 from each side.

$0 = x^2 - 18x + 32$  dividing each side by 4.

The related function for this equation is

$$f(x) = y = x^2 - 18x + 32.$$

$x$	2	4	16	20
$y$	32	-24	0	72



Notice that the function crosses the x-axis twice and the function has a degree of 2. These intercept points are called the zero of a function. The x values, 2 and 16 are the roots of the equation  $x^2 - 18x + 32 = 0$


**Example:** Graph the related quadratic function

$y = x^2 - 2x - 15$  by finding and graphing the ordered pairs that satisfy the function.

**Solution:**

$x$	-3	-1	1	5
$y$	0	-12	-16	0

The graphs tells that the solutions are



-3 and 5, where the curve intersects the x-axis.

## ii. Factoring Method

Another way of solving a quadratic equation is by factoring. The factoring method depends on the zero-product property.

### Zero Product Property:

For any real number a and b, if  $ab = 0$ , then either  $a = 0$  or  $b = 0$ .

- i. To solve the quadratic equation by factorization first write it in standard form, then factorize the polynomial on the left-hand side of the equation.
- ii. Use zero product property.
- iii. Now solve linear equations and write solution set.

For example, to solve  $x^2 = 4$  by factorization we do the following steps:

$$\begin{aligned}x^2 &= 4 \\x^2 - 4 &= 0 && \text{(standard form)} \\(x)^2 - (2)^2 &= 0 \\(x - 2)(x + 2) &= 0 && \text{(factorization)} \\x - 2 = 0 \text{ or } x + 2 &= 0 && \text{(linear equations)} \\x = 2 \text{ or } x = -2 & && \text{(roots)} \\S.S. &= \{-2, 2\} && \text{(solutionset)}\end{aligned}$$

**Example:** Factorize to solve the equations.

i.  $x^2 - 18x + 32 = 0$

ii.  $x^2 - 2x - 15 = 0$

**Solution:**

i.  $x^2 - 18x + 32 = 0$   
 $x^2 - 16x - 2x + 32 = 0$   
 $x(x - 16) - 2(x - 16) = 0$   
 $(x - 2)(x - 16) = 0$   
 $x - 2 = 0 \text{ or } x - 16 = 0$   
 $x = 2 \text{ or } x = 16$   
 $S.S = \{2, 16\}$

ii.  $x^2 - 2x - 15 = 0$   
 $x^2 - 5x + 3x - 15 = 0$   
 $x(x - 5) + 3(x - 5) = 0$   
 $(x - 5)(x + 3) = 0$   
 $x - 5 = 0 \text{ or } x + 3 = 0$   
 $x = 5 \text{ or } x = -3$   
 $S.S = \{-3, 5\}$

## iii. Completing Square Method

In a city, a square field has been maintained for playing. Each side of the field is 110 m long. The management has decided to equally expand two of the sides of the field for parking spaces so that the entire region has an area of 14000  $m^2$ . What is the width of the parking strip being added?

Let  $x$  represents the width of each parking strip.  
After expanding each side of the field would be  
 $110+x$  meters long.

The equation expresses the area of the new field,  
which is still square.


$$(x + 110)^2 = 14000$$

$$\sqrt{(x + 110)^2} = \sqrt{14000}$$

$$x + 110 = \pm\sqrt{14000}$$

$$x = -110 \pm \sqrt{14000}$$

$$x = 8.32m$$



Where the negative value is neglected.

Quadratic equations can be solved in the same way as long as one side of the equation contains a perfect square. When the equation does not contain a perfect square, we can use process called completing the square to create a perfect square.

For an expression of the form  $x^2 + bx$ , we can add a constant  $c$  in the expression so that the expression  $x^2 + bx + c$  is a perfect square trinomial. This process is called completing square.

### Procedure

- Step I:** Write the equation in standard form.
- Step II:** Shift constant term to the right hand side of the equation.
- Step III:** To make the left hand side a complete square, add a suitable term to both sides of the equation. If the coefficient of  $x^2$  is 1, then it is easy to complete the square. Just add the square of half the coefficient of  $x$  and get a complete square.
- Step IV:** Take the square root on both sides and form two linear equations.
- Step V:** Solve the linear equations and find roots.

### Example:

Solve  $3x^2 - 11x - 4 = 0$  by completing square.

### Solution:

$$3x^2 - 11x - 4 = 0$$

$$3x^2 - 11x = 4 \quad \text{Shift 4 right side}$$

$$x^2 - \frac{11}{3}x = \frac{4}{3} \quad \text{dividing each side by 3.}$$

### Check Point:

Describe the error.

$$x^2 - 14x = 11$$

$$x^2 - 14x + 49 = 11$$

$$(x - 7)^2 = 11$$

$$x - 7 = \pm\sqrt{11}$$

$$x = 7 \pm \sqrt{11}$$

$$x^2 - \frac{11}{3}x + \frac{121}{36} = \frac{4}{3} + \frac{121}{36} \text{ adding } \frac{121}{36} \text{ on both sides.}$$

$$\left(x - \frac{11}{6}\right)^2 = \frac{169}{36} \text{ factor}$$

$$\sqrt{\left(x - \frac{11}{6}\right)^2} = \sqrt{\frac{169}{36}} \text{ taking square root}$$

$$x - \frac{11}{6} = \pm \frac{13}{6}$$

$$x - \frac{11}{6} = \frac{13}{6} \text{ or } x - \frac{11}{6} = -\frac{13}{6}$$

$$x = 4 \text{ or } x = -\frac{1}{3} \quad S. S. = \{-\frac{1}{3}, 4\}$$

### Example:

Solve  $3x^2 + 6x - 9 = 0$  by the completing square method.

### Solution:

$$3x^2 + 6x - 9 = 0$$

$$3x^2 + 6x = 9$$

$$x^2 + 2x = 3 \quad (\text{dividing by 3})$$

$$x^2 + 2x + 1 = 3 + 1 \quad (\text{adding 1 to both sides})$$

$$(x + 1)^2 = 4$$

$$x + 1 = \pm \sqrt{4} \quad (\text{taking square root})$$

$$x + 1 = \pm 2$$

$$x = \pm 2 - 1$$

$$S. S. = \{1, -3\}$$

### iv. Quadratic Formula

We have learnt several ways to solve quadratic equations. Each has its limitations. The question we think is that is there any formula that will work for any quadratic equation? The answer is 'yes' and the formula is called the quadratic formula. The formula is derived from solving the general form of a quadratic equation for  $x$ .

#### Key Fact:

### Derivation of the Quadratic Formula

Standard form of quadratic equation is:  $ax^2 + bx + c = 0$

$$ax^2 + bx = -c$$

$$x^2 + \frac{b}{a}x = -\frac{c}{a} \quad (\text{Divide by } a \text{ both sides})$$

$$x^2 + \frac{b}{a}x + \left(\frac{b}{2a}\right)^2 = \left(\frac{b}{2a}\right)^2 - \frac{c}{a} \quad (\text{Add } \left(\frac{b}{2a}\right)^2 \text{ both sides})$$

The solution of a quadratic equation of the form

$$ax^2 + bx + c = 0 \text{ with } a \neq 0 \text{ is}$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x^2 + 2x \frac{b}{2a} + \left(\frac{b}{2a}\right)^2 = \left(\frac{b}{2a}\right)^2 - \frac{c}{a}$$

$$\left(x + \frac{b}{2a}\right)^2 = \frac{b^2 - 4ac}{4a^2} \quad (\text{Simplify the left side})$$

$$x + \frac{b}{2a} = \pm \sqrt{\frac{b^2 - 4ac}{4a^2}} \quad (\text{Taking square root both sides})$$

$$x = -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{\sqrt{4a^2}} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

This equation is known as quadratic formula.

**Example:** Solve  $2x^2 + 7x = 4$  using quadratic formula.

**Solution:**  $2x^2 + 7x = 4$

$$2x^2 + 7x - 4 = 0 \quad \text{General form}$$

$$a = 2, b = 7, c = -4$$

$$\text{We know that: } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-7 \pm \sqrt{(7)^2 - 4(2)(-4)}}{2(2)}$$

$$x = \frac{-7 \pm \sqrt{49 + 32}}{4}$$

$$x = \frac{-7 \pm \sqrt{81}}{4} \Rightarrow \frac{-7 \pm 9}{4}$$

$$x = \frac{-7 + 9}{4}, x = \frac{-7 - 9}{4}$$

$$x = \frac{2}{4}, x = \frac{-16}{4}$$

$$x = \frac{1}{2}, x = -4$$

$$\text{S. S.} = \left\{ -4, \frac{1}{2} \right\}$$

### Procedure

To solve a quadratic equation by using the quadratic formula, we proceed as follows:

**Step I:** Write the equation in standard form.

**Step II:** Compare it with  $ax^2 + bx + c = 0$ , to get the values of  $a, b$  and  $c$ .

**Step III:** Write the quadratic formula for  $x$ .

**Step IV:** Put values of  $a, b$  and  $c$ , then simplify.

**Step V:** Write solution set.

### Exercise 2.1

1. Write the following quadratic equation in standard form:

$$\text{i. } (x+2)(x-3) = 5 \quad \text{ii. } (x-5)^2 - (2x+4)^2 = 7 \quad \text{iii. } x = x(x-1)$$

2. Solve the following equations by factoring method.

$$\text{i. } (x-1)(x-4) = 0 \quad \text{ii. } x^2 - 2x + 1 = 0$$

iii.  $x^2 - 7x - 8 = 0$

iv.  $x^2 - 4x + 4 = (2x - 7)^2$

v.  $\left(2x + \frac{7}{4}\right)^2 = \frac{48x^2 + 529}{16}$

3. Solve the following equations by completing the square method:

i.  $x^2 + 4x - 32 = 0$

ii.  $x^2 + 8x = 0$

iii.  $x^2 + 6x - 9 = 0$

iv.  $3x^2 + 12x + 8 = 0$

v.  $x^2 + x + 1 = 0$

vi.  $4x^2 - 8x - 5 = 0$

4. Solve the following equations by quadratic formula:

i.  $x^2 - 9 = 0$

ii.  $2x^2 + 5x + 1 = 0$

iii.  $x^2 - 23x - 24 = 0$

iv.  $(x+1)^2 = (2x-1)^2$

v.  $\frac{x+1}{2} - \frac{x(x+2)}{3} = 0$

vi.  $(x-2)(x-6) = (2x+1)(x+1)$

5. Solve  $x^2 + 6x = -9$  by graphing and factoring method.

6. Graph the function  $y = x^2 + 2x + 4$  and verifying solution by completing square method.

7. Explain each term:

i. Solution

ii. root

iii. Zero of a function

iv. x-intercept

8. Can a quadratic equation have more than two solutions? Why or why not?

9. Plans for rectangular ice-skating rink that is 30m by 60m have to be revised to double the area of the playground by adding strips of the same width to a side and an end of the area to form a rectangle. Find the width of the strips. What are the dimensions of the playground now?

10. If a car has an initial speed of 20m/s and a constant acceleration of 2m/s<sup>2</sup>, determine the amount of time it takes to travel 145m. (Use:  $S = v_i t + \frac{1}{2} a t^2$ )

## Equation Reducible to Quadratic Form

Some equations are not quadratic, but to solve them we can reduce them to quadratic form.

The example below illustrates solving an equation that is not a quadratic equation itself, but contains a factor that is a quadratic. The method we use to solve the quadratic part of that equation.

**Example:** A cuboid box has a base that is  $x$  cm long. The width of the base is 2 cm less than its length. The height is 3 cm greater than the length of the base. The measure of the volume is 6 times the measure of the length of the base. Find the dimensions of the box.

### Solution:

The volume of the rectangular box is the product of the length, width and height of box.

Here, Length =  $x$ , Width =  $x - 2$ , Height =  $x + 3$

∴ The volume is:

$$V = \text{Length} \times \text{Width} \times \text{Height} = lwh$$

$$V = 6 \times \text{the measure of the length of the base} \dots \text{(i)}$$

$$\text{As, } V = x(x - 2)(x + 3) \dots \text{(ii)}$$

$$\therefore 6x = x(x - 2)(x + 3) \quad (\text{from (i) and (ii)})$$

$$6x = x^3 + x^2 - 6x \Rightarrow x^3 + x^2 - 12x = 0$$

$$x(x^2 + x - 12) = 0 \Rightarrow x(x - 3)(x + 4) = 0$$

We use the zero-product property

$$x = 0, x + 4 = 0, x - 3 = 0$$

$$x = 0, x = -4, x = 3$$

Since we are dealing with the dimensions of a box,  
we ignore 0 and  $-4$ , the solution is:

$$\text{length } x = 3$$

$$\text{width } = x - 2 = 3 - 2 = 1$$

$$\text{height } = x + 3 = 3 + 3 = 6$$

Hence, the dimensions of the box are  $3\text{cm} \times 1\text{cm} \times 6\text{cm}$ .

We have many types of equations that are not quadratic, but we can write in a form that resembles a quadratic equation. For example, the equation  $9x^4 - 8x^2 - 1 = 0$  can be written as:  $9(x^2)^2 - 8x^2 - 1 = 0$ . Equations that can be written in this way are said to be equations reducible to quadratic form.

Some of equations that can be reduced to quadratic equations, are given below.

### Equations of the Form $ax^4 + bx^2 + c = 0$

It is an equation of degree 4. It can be reduced to quadratic form by using the substitution  $x^2 = y$ . In this way the above equation becomes  $ay^2 + by + c = 0$ . Solve it by any of the three methods discussed earlier. Replace  $y$  with its substitute  $x^2$ . Find values of  $x$  by taking the square root and write the solution set.

**Example:** Solve  $9x^4 - 8x^2 - 1 = 0$ .

**Solution:**

$$9x^4 - 8x^2 - 1 = 0$$

Put  $x^2 = y$  which gives  $x^4 = y^2$

$$\therefore 9y^2 - 8y - 1 = 0$$

$$9y^2 - 9y + y - 1 = 0$$


$$9y(y - 1) + 1(y - 1) = 0$$

$$(y - 1)(9y + 1) = 0$$

$$y - 1 = 0 \text{ or } 9y + 1 = 0$$

Back substitution gives:

$$x + 3$$



$$x + 3$$

$$x$$

$$x - 2$$

$$x + 2$$

$$x + 3$$

### Challenge

The rectangle has an area of 42 square centimeters. What is the value of length of sides?

$$x^2 - 1 = 0 \quad \text{or} \quad 9x^2 + 1 = 0$$

$$x = \pm\sqrt{1} \quad \text{or} \quad x = \pm\sqrt{\frac{-1}{9}}$$

$$x = \pm 1 \quad \text{or} \quad x = \pm \frac{i}{3}$$

### Equations of the Form $aP(x) + \frac{b}{P(x)} = c$

To solve this type of equation, use a substitution  $P(x) = y$ . Multiply the equation by  $y$  and form a quadratic equation involving the variable  $y$ . Solve the new equation for  $y$  and replace  $y$  by  $P(x)$ . In this way, you will get two simple equations involving variable  $x$ . Solve these equations and write the solution set.

**Example:** Solve:  $3x + \frac{4}{x} = 7$

$$\text{Solution: } 3x + \frac{4}{x} = 7$$

$$3x^2 + 4 = 7x$$

$$3x^2 - 7x + 4 = 0$$

$$3x^2 - 3x - 4x + 4 = 0$$

$$3x(x - 1) - 4(x - 1) = 0 \Rightarrow x - 1 = 0 \quad \text{or} \quad 3x - 4 = 0$$

$$x = 1 \quad \text{or} \quad x = \frac{4}{3}$$

$$\therefore S.S. = \left\{ 1, \frac{4}{3} \right\}$$

**Example:** Solve  $2(x^2 + 1) - \frac{2}{x^2 + 1} = 3$

$$\text{Solution: } 2(x^2 + 1) - \frac{2}{x^2 + 1} = 3$$

$$\text{Put } x^2 + 1 = y$$

$$\text{So, } 2y - \frac{2}{y} = 3$$

$$2y^2 - 3y - 2 = 0$$

$$2y(y - 2) + 1(y - 2) = 0$$

$$y - 2 = 0 \quad \text{or} \quad 2y + 1 = 0$$

### Challenge

(i) Solve for real roots,  $y^3 - 27 = 0$

(ii) Solve:

$$x^2 + 3x + 7 = \frac{6}{x^2 + 3x + 2}$$

Back substitution: When  $y = 2$

$$x^2 + 1 - 2 = 0$$

$$x^2 - 1 = 0$$

$$x = \pm 1$$

$$\text{Solution} = \pm 1, \pm \sqrt{\frac{3}{2}}i$$

When  $y = \frac{-1}{2}$

$$x^2 + 1 = -\frac{1}{2}$$

$$2x^2 = -3 \Rightarrow x^2 = \frac{-3}{2}$$

$$x = \pm \sqrt{\frac{3}{2}}i$$

## Reciprocal Equations

If an equation is not affected by the replacement of a variable by its reciprocal, then it is called a reciprocal equation. The reciprocal of a root of such an equation is also its root.

Consider the equation  $ax^4 + bx^3 + cx^2 + bx + a = 0$ . → (i)

$$a\left(\frac{1}{x}\right)^4 + b\left(\frac{1}{x}\right)^3 + c\left(\frac{1}{x}\right)^2 + b\left(\frac{1}{x}\right) + a = 0 \quad (x \text{ is replaced by } \frac{1}{x})$$

$$\frac{a}{x^4} + \frac{b}{x^3} + \frac{c}{x^2} + \frac{b}{x} + a = 0 \quad \therefore$$

$$a + bx + cx^2 + bx^3 + ax^4 = 0 \quad (\text{equation is multiplied by } x^4)$$

$$ax^4 + bx^3 + cx^2 + bx + a = 0. \quad \rightarrow \text{(ii)}$$

Equations (i) and (ii) are same.

To solve the reciprocal equation (i), arrange it in the form

$$a\left(x^2 + \frac{1}{x^2}\right) + b\left(x + \frac{1}{x}\right) + c = 0$$

Put  $y = x + \frac{1}{x}$  and find the value of  $x^2 + \frac{1}{x^2} = y^2 - 2$ .

In this way equation (i) is reduced to quadratic form. Solve this equation for  $y$  and then replace  $y$  with its substitute  $x + \frac{1}{x}$ . Finally find values of  $x$  and write solution set.

**Example:** Solve the equation:

$$6x^4 - 35x^3 + 62x^2 - 35x + 6 = 0.$$

**Solution:**

$$6x^4 - 35x^3 + 62x^2 - 35x + 6 = 0.$$

$$6x^2 - 35x + 62 - \frac{35}{x} + \frac{6}{x^2} = 0 \quad (\text{dividing by } x^2)$$

$$6(x^2 + \frac{1}{x^2}) - 35(x + \frac{1}{x}) + 62 = 0 \quad \rightarrow \text{(i)}$$

$$\text{Let } x + \frac{1}{x} = y$$

$$\left(x + \frac{1}{x}\right) = y^2$$

$$x^2 + \frac{1}{x^2} + 2 = y^2$$

$$x^2 + \frac{1}{x^2} = y^2 - 2$$

Put in equation (i)

$$6(y^2 - 2) - 35y + 62 = 0$$

$$6y^2 - 12 - 35y + 62 = 0$$

### Critical Thinking

**Example:** Solve  $y - 7\sqrt{y} - 8 = 0$ ,  $y \in R$

**Solution:**  $y - 7\sqrt{y} - 8 = 0$

$$\text{Let } \sqrt{y} = z$$

$$\text{Then } y = z^2$$

Equation will become:

$$z^2 - 7z - 8 = 0$$

$$z = \frac{-(-7) \pm \sqrt{(-7)^2 - 4(1)(-8)}}{2(1)}$$

$$z = \frac{+7 \pm \sqrt{49 + 32}}{2} = \frac{+7 \pm \sqrt{81}}{2}$$

$$z = \frac{+7 \pm 9}{2}$$

$$z = \frac{+7+9}{2}, z = \frac{+7-9}{2}$$

$$z = 8, z = -1$$

Back substitution gives:

$$\sqrt{y} = 8, \sqrt{y} = -1$$

$y = 64$ , (No real roots for  $\sqrt{y} = -1$ )

So, the solution is = 64

$$6y^2 - 35y + 50 = 0$$

$$6y^2 - 15y - 20y + 50 = 0$$

$$3y(2y - 5) - 10(2y - 5) = 0$$

$$(2y - 5)(3y - 10) = 0 \Rightarrow 2y - 5 = 0 \text{ or } 3y - 10 = 0$$

$$2\left(x + \frac{1}{x}\right) - 5 = 0 \quad \text{or} \quad 3\left(x + \frac{1}{x}\right) - 10 = 0 \quad (\text{Substituting the value of } y)$$

$$2x + \frac{2}{x} - 5 = 0 \quad \text{or} \quad 3x + \frac{3}{x} - 10 = 0$$

$$2x^2 + 2 - 5x = 0 \quad \text{or} \quad 3x^2 + 3 - 10x = 0$$

$$2x^2 - 5x + 2 = 0 \quad \text{or} \quad 3x^2 - 10x + 3 = 0$$

$$2x^2 - 4x - x + 2 = 0 \quad \text{or} \quad 3x^2 - 9x - x + 3 = 0$$

$$2x(x - 2) - 1(x - 2) = 0 \quad \text{or} \quad 3x(x - 3) - 1(x - 3) = 0$$

$$(x - 2)(2x - 1) = 0 \quad \text{or} \quad (x - 3)(3x - 1) = 0$$

$$x - 2 = 0 \quad \text{or} \quad 2x - 1 = 0 \quad \text{or} \quad x - 3 = 0 \quad \text{or} \quad 3x - 1 = 0$$

$$x = 2 \quad \text{or} \quad x = \frac{1}{2} \quad \text{or} \quad x = 3 \quad \text{or} \quad x = \frac{1}{3} \Rightarrow S.S. = \{2, \frac{1}{2}, 3, \frac{1}{3}\}$$

### Exponential Equations

An equation involving expressions of the form  $k^x$  is called an exponential equation where k is a constant. In exponential equations the exponents are variables. We will discuss only those exponential equations which can be reduced to quadratic form.

Exponential equations reducible to quadratic form are of the type:

$$ak^{2x} + bk^x + c = 0, a \neq 0.$$

We use the substitution  $k^x = y$  to reduce the above equation to quadratic form. Then we find y and replace it with  $k^x$ . In this way, roots of the exponential equations are found.

**Example:** Solve the following exponential equations

i.  $2^{2x} - 2 \times 2^x + 1 = 0$

ii.  $9^x - 3^{x+1} - 4 = 0$

**Solution:**

i.  $2^{2x} - 2 \times 2^x + 1 = 0 \rightarrow \text{(i)}$

ii.  $9^x - 3^{x+1} - 4 = 0 \rightarrow \text{(i)}$

Put  $2^x = y$ , so  $2^{2x} = y^2$

$3^{2x} - 3^x \times 3 - 4 = 0$

∴ Equation (i) becomes:

Put  $3^x = y$ , so  $3^{2x} = y^2$

$y^2 - 2y + 1 = 0$

∴ Equation (i) becomes:

$(y - 1)^2 = 0$

$y^2 - 3y - 4 = 0$

$y - 1 = 0$

$y^2 - 4y + y - 4 = 0$

$y = 1$

$y(y - 4) + 1(y - 4) = 0$

Replace y by its substitute  $2^x$

$(y - 4)(y + 1) = 0$

$y = 4 \text{ or } y = -1$

$3^x = 4 \text{ or } 3^x = -1$

$$2^x = 1 = 2^0$$

$$x = 0$$

$$S.S. = \{0\}$$

Converting  $3^x = 4$  into logarithmic form, we get

$$x = \log_3 4$$

Solution of  $3^x = -1$  is not possible.

$$\therefore S.S. = \{\log_3 4\}$$

### Equations of the Form $(x+a)(x+b)(x+c)(x+d) = k; a+b=c+d$

In this type, we multiply the pairs of factors after rearranging them in such a way that the variable terms of both products are the same or one of them is multiple of the other. Choose a substitute for the common terms and form a quadratic equation. Solve the new equation and then replace the substitute. Find the values of original variable involved in the given equation and write the solution set.

#### Example:

Solve the following equations by using suitable substitution.

$$(x+1)(x+2)(x+3)(x+4) = 24$$

#### Solution:

$$(x+1)(x+2)(x+3)(x+4) = 24$$

$$(x+1)(x+4)(x+2)(x+3) = 24$$

$$(x^2 + 5x + 4)(x^2 + 5x + 6) = 24$$

$$\text{Put } x^2 + 5x = y$$

$$(y+4)(y+6) = 24$$

$$y^2 + 10y + 24 = 24$$

$$y^2 + 10y = 0$$

$$y(y+10) = 0$$

$$y = 0 \text{ or } y + 10 = 0$$

$$\text{For } y = 0$$

$$x^2 + 5x = 0$$

$$x(x+5) = 0$$

$$x = 0 \text{ or } x+5 = 0$$

$$x = 0 \text{ or } x = -5$$

$$S.S. = \{0, -5, \frac{-5 \pm \sqrt{15}i}{2}\}$$

$$\text{For } y + 10 = 0$$

$$x^2 + 5x + 10 = 0$$

Here,  $a = 1, b = 5$  and  $c = 10$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \frac{-5 \pm \sqrt{5^2 - 4(1)(10)}}{2(1)}$$

$$x = \frac{-5 \pm \sqrt{25 - 40}}{2}$$

$$x = \frac{-5 \pm \sqrt{-15}}{2}$$

$$x = \frac{-5 \pm \sqrt{15}i}{2}$$

## Exercise 2.2

**1.** Reduce the following equations to quadratic form using suitable substitution.

- i.  $ax^4 + bx^2 + c = 0$
- ii.  $9x^6 - 3x^3 + 7 = 0$
- iii.  $3x + \frac{4}{6x-2} = -1$
- iv.  $(x+1)^2 + \frac{3}{(x+1)^2} = 4$ .
- v.  $3\left(x^2 + \frac{1}{x^2}\right) + 8\left(x + \frac{1}{x}\right) + 11 = 0$
- vi.  $3x^4 + 7x^3 + 5x^2 - 7x + 3 = 0$
- vii.  $ak^{2x} + bk^x + c = 0$
- viii.  $8 \times 4^x - 7 \times 2^x - 1 = 0$
- ix.  $(x-1)(x-2)(x+3)(x-6) = 6$
- x.  $(2x-1)(2x-7)(x-3)(x-1) = 8$

**2.** Solve the following equations by reducing them to quadratic form.

- i.  $x^4 - 20x^2 + 64 = 0$
- ii.  $x^4 + 16x^2 - 225 = 0$
- iii.  $x^{\frac{2}{5}} + 5x^{\frac{1}{5}} + 6 = 0$
- iv.  $3x^2 + \frac{4}{x^2} = 7$
- v.  $5(x+1) + \frac{3}{x+1} = 8$
- vi.  $5x^2 + \frac{36}{5x^2 + 4} = 16$
- vii.  $2x^4 - x^3 - 6x^2 - x + 2 = 0$
- viii.  $12x^4 + 11x^3 - 146x^2 + 11x + 12 = 0$
- ix.  $4\left(x^2 + \frac{1}{x^2}\right) - \left(x - \frac{1}{x}\right) - 11 = 0$
- x.  $2^{2x} - 34 \times 2^x + 64 = 0$
- xi.  $3^{2x} - 12 \times 3^x + 27 = 0$
- xii.  $5^{2x} - 150 \times 5^x + 3125 = 0$
- xiii.  $(x+2)(x-3)(x+10)(x+5) = -396$
- xiv.  $x(x-1)(x+2)(x+3) = 40$
- xv.  $(x-2)(x-6)(x+4)(x+8) + 256 = 0$

**3.** Solve the following equations by factoring without using substitution.

- i.  $2x^4 - 3x^2 + 1 = 0$
- ii.  $8x^6 - 7x^3 - 1 = 0$  (Find only real roots)
- iii.  $x^2 + \frac{1}{x^2} = 2$
- iv.  $4 \times 2^{2x} - 4 \times 2^x + 1 = 0$

- Write two examples of equations that are not quadratic, but can be written in quadratic form.
- How would you solve the equation  $(y-4)^2 - 7(y-4) = -6$ ? Write explanation and then solve the equation.
- Solve  $y^3 = 125$ .

## Nature of Roots of a Quadratic Equation

We have already derived the quadratic formula by solving the general form of quadratic equation for  $x$ . We know that the solutions of a quadratic equation of the form  $ax^2 + bx + c = 0$  with  $a \neq 0$ , are given by this formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$


To define discriminant, first we solve the following examples by using the quadratic formula:

**Example A:** Solve.  $x^2 - 3x - 28 = 0$

**Solution:** We have  $a = 1$ ,  $b = -3$  and  $c = -28$

$$\begin{aligned} x &= \frac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(-28)}}{2(1)} \\ &= \frac{3 \pm \sqrt{121}}{2} \\ x &= \frac{3 \pm 11}{2} = 7, -4 \end{aligned}$$

The roots of equation are rational and unequal.




**Example B:** Solve  $x^2 - 8x + 16 = 0$

**Solution:** We have  $a = 1$ ,  $b = -8$  and  $c = 16$

$$\begin{aligned} x &= \frac{-(-8) \pm \sqrt{(-8)^2 - 4(1)(16)}}{2(1)} \\ &= \frac{8 \pm 0}{2} \\ x &= \frac{8+0}{2}, x = \frac{8-0}{2} \end{aligned}$$

$$x = 4, x = 4$$

The roots of equation are rational and equal.




**Example C:** Solve  $3y^2 - 5y + 9 = 0$

**Solution:** We have  $a = 3$ ,  $b = -5$  and  $c = 9$

$$y = \frac{-(-5) \pm \sqrt{(-5)^2 - 4(3)(9)}}{2(3)}$$

$$= \frac{5 \pm \sqrt{-83}}{6} = \frac{5 \pm \sqrt{83}i}{6}$$

Since radical contains a negative value,  
therefore, the roots of equation are imaginary.




**Example D:** Solve  $3t^2 - 6t + 2 = 0$

**Solution:** We have  $a = 3$ ,  $b = -6$  and  $c = 2$

$$t = \frac{-(-6) \pm \sqrt{(-6)^2 - 4(3)(2)}}{2(3)}$$

$$= \frac{6 \pm \sqrt{12}}{6}$$

The roots are irrational and unequal.



These four examples demonstrate a pattern that is useful in determining the nature of the roots of a quadratic equation. In the quadratic formula, the expression under the sign ' $b^2 - 4ac$ ', is called the discriminant. The discriminant tells the nature of roots of a quadratic equation.

Equation	Value of the discriminant	Roots	Nature of Roots
$x^2 - 3x - 28 = 0$ (Exp. A)	$b^2 - 4ac = 121$	$7, -4$	Roots are rational and unequal.
$x^2 - 8x + 16 = 0$ (Exp. B)	$b^2 - 4ac = 0$	$4, 4$	Roots are rational and equal.
$3y^2 - 5y + 9 = 0$ (Exp. C)	$b^2 - 4ac = -83$	$\frac{5 \pm \sqrt{83}i}{6}$	Roots are imaginary/complex.
$3t^2 - 6t + 2 = 0$ (Exp. D)	$b^2 - 4ac = 24$	$\frac{3 \pm \sqrt{3}}{3}$	Roots are irrational and unequal.

The above chart shows that if the value of the discriminant is a perfect square or 0, the roots are real and rational. Other positive discriminant will yield irrational roots. A negative discriminant means roots will be imaginary/complex.

**Example:** Find the value of discriminant and describe the nature of roots.

a.  $2x^2 + x - 3 = 0$       b.  $x^2 + 8 = 0$

**Solution:**

<p>a. <math>2x^2 + x - 3 = 0</math>  <math>a = 2, b = 1, c = -3</math>  <math>\text{Disc} = b^2 - 4ac</math>  <math>= (1)^2 - 4(2)(-3) = 25</math></p>	<p>b. <math>x^2 + 8 = 0</math>  <math>a = 1, b = 0, c = 8</math>  <math>\text{Disc} = b^2 - 4ac</math>  <math>= (0)^2 - 4(1)(8) = -32</math></p>
--	--

The value of discriminant is positive and a perfect square. So, the given equation has two real roots and they are rational and unequal.

The value of discriminant is negative.  
So, the given equation has two imaginary roots.

#### Check Point:

Verify the nature of roots of  $9x^2 - 12x + 4 = 0$ , by solving it.

**Example:** Find the value of m, when  $x^2 - 3x + m = 0$  has equal roots.

**Solution:**  $x^2 - 3x + m = 0$

Here  $a = 1, b = 3, c = m$

It is given that equation has equal roots (same root) so the discriminant is zero.

i.e.  $\text{Disc} = b^2 - 4ac = 0$

$$(3)^2 - 4(1)(m) = 0$$

$$9 - 4m = 0 \quad \text{or} \quad 4m = 9 \quad \text{or} \quad m = \frac{9}{4}$$

### Exercise 2.3

- Find the discriminant of the following quadratic equations.
  - $x^2 + 6x - 27 = 0$
  - $x^2 - x - 12 = 0$
  - $8x^2 + 2x + 1 = 0$
  - $12x^2 - 11x - 15 = 0$
- Discuss the nature of roots of the following quadratic equations.
  - $x^2 - 2x - 15 = 0$
  - $x^2 + 3x - 4 = 0$
  - $12x^2 + x - 20 = 0$
  - $x^2 + 2x + 8 = 0$
  - $x^2 + 3x - 9 = 0$
- For what value of k,  $9x^2 - kx + 16 = 0$  is a perfect square?
- If roots of  $x^2 + kx + 9 = 0$  are equal, find k?
- Show that the roots of  $2x^2 + (mx - 1)^2 = 3$ , are equal if  $3m^2 + 4 = 0$ .

6. Find the value of "m" when roots of the following quadratic equations are equal.
- $x^2 - 6x + m = 0$
  - $m^2 x^2 + (2m + 1)x + 1 = 0$
  - $(m + 3)x^2 + (m + 1)x + m + 1 = 0$
7. Show that the roots of the equation  $(a^2 + b^2)x^2 + 2(ac + bd)x + c^2 + d^2 = 0$  are imaginary. Moreover, it shows repeated roots if  $ad = bc$ .
8. Show that the roots of the equation  $(ax + c)^2 = 4bx$  will be equal, if  $b = ac$ .
9. Show that the roots of the following equations are real.
- $mx^2 - 2mx + m - 1 = 0$
  - $bx^2 + ax + a - b = 0$
10. Show that the roots of the following equation are real.  
 $(a + b)x^2 - ax - b = 0$

## Roots and Coefficients of the Quadratic Equation

Let  $S_1, S_2$  be the roots of  $ax^2 + bx + c = 0$ , then using the quadratic formula:

$$S_1 = \frac{-b + \sqrt{b^2 - 4ac}}{2a}, S_2 = \frac{-b - \sqrt{b^2 - 4ac}}{2a}$$

$$\begin{aligned}\text{Sum of roots} &= S_1 + S_2 = \frac{-b + \sqrt{b^2 - 4ac}}{2a} + \frac{-b - \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-b + \sqrt{b^2 - 4ac} - b - \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-2b}{2a} = \frac{-b}{a}\end{aligned}$$

$$\begin{aligned}\text{and product of roots} &= S_1 S_2 = \left( \frac{-b + \sqrt{b^2 - 4ac}}{2a} \right) \left( \frac{-b - \sqrt{b^2 - 4ac}}{2a} \right) \\ &= \frac{(-b)^2 - (\sqrt{b^2 - 4ac})^2}{4a^2} \\ &= \frac{4ac}{4a^2} = \frac{c}{a}\end{aligned}$$

$$\therefore \text{Sum of the roots } S_1 + S_2 = -\frac{b}{a} = -\frac{\text{coefficient of } x}{\text{coefficient of } x^2}$$

$$\text{and product of roots } S_1 S_2 = \frac{c}{a} = \frac{\text{constant term}}{\text{coefficient of } x^2}$$

**Example:** Write down the sum and product of the roots of quadratic equation:

- $x^2 - x + 2 = 0$
- $x^2 - 7x + 5 = 0$
- $3x^2 - 11x - 4 = 0$
- $6s^2 + 2s + 3 = 0$

**Solution:**

(i) $x^2 - x + 2 = 0$ Here $a = 1, b = -1, c = 2$ Sum of the roots = $S_1 + S_2$ $= \frac{-b}{a} = \frac{-(\text{-}1)}{1} = 1$ Product of roots = $S_1 S_2$ $= \frac{c}{a} = \frac{2}{1} = 2$	(ii) $3x^2 - 11x - 4 = 0$ $a = 3, b = -11, c = -4$ Sum of roots = $S_1 + S_2$ $= \frac{-b}{a} = \frac{-(\text{-}11)}{3} = \frac{11}{3}$ Product of roots = $S_1 S_2$ $= \frac{c}{a} = \frac{-4}{3}$
(iii) $x^2 - 7x + 5 = 0$ $a = 1, b = -7, c = 5$ $S_1 + S_2 = \frac{-b}{a} = \frac{-(\text{-}7)}{1} = 7$ $S_1 S_2 = \frac{c}{a} = \frac{5}{1} = 5$	(iv) $6s^2 + 2s + 3 = 0$ $a = 6, b = 2, c = 3$ $S_1 + S_2 = \frac{-b}{a} = \frac{-(2)}{6} = \frac{-1}{3}$ $S_1 S_2 = \frac{c}{a} = \frac{3}{6} = \frac{1}{2}$

**Example:** If  $S_1, S_2$  are the roots of  $x^2 - 8x + 20 = 0$ . Find the values of

$$(i) S_1^2 + S_2^2 \quad (ii) (S_1 - S_2)^2 \quad (iii) \frac{1}{S_1+1} + \frac{1}{S_2+1}$$

**Solution:** Here  $a = 1, b = -8, c = 20$

Since  $S_1, S_2$  are the roots of  $x^2 - 8x + 20 = 0$ ,

$$\therefore S_1 + S_2 = \frac{-b}{a} = \frac{-(\text{-}8)}{1} = 8 \quad \text{and} \quad S_1 S_2 = \frac{c}{a} = \frac{20}{1} = 20$$

$$(i) \text{ Now } S_1^2 + S_2^2 = S_1^2 + S_2^2 + 2S_1 S_2 - 2S_1 S_2 \\ = (S_1 + S_2)^2 - 2S_1 S_2 \\ = (8)^2 - 2(20) \\ = 64 - 40 = 24$$

$$(ii) (S_1 - S_2)^2 = S_1^2 + S_2^2 - 2S_1 S_2 \\ = S_1^2 + S_2^2 + 2S_1 S_2 - 4S_1 S_2 \\ = (S_1 + S_2)^2 - 4S_1 S_2 \\ = (8)^2 - 4(20) \\ = 64 - 80 = -16$$

$$(iii) \frac{1}{S_1+1} + \frac{1}{S_2+1} = \frac{S_2+1+S_1+1}{(S_2+1)(S_1+1)} \\ = \frac{S_1+S_2+2}{S_1 S_2 + (S_1 + S_2) + 1} \\ = \frac{8+2}{20+8+1} = \frac{10}{29}$$

**Note:**

Instead of  $S_1, S_2$ , we can use  $\alpha$  and  $\beta$ .

## Formation of an Equation whose Roots are Given

We have solved several quadratic equations and found the roots. Here, we may know the roots of the quadratic equation but we do not know the equation itself. Consider the roots of a quadratic quadratic equation are 4 and  $-\frac{1}{3}$ .

let  $x = 4$  and  $x = -\frac{1}{3}$  are roots and  $x - 4 = 0$  and  $x + \frac{1}{3} = 0$  are factors of required equation.

$$(x - 4)(3x + 1) = 0 \quad \text{or} \quad 3x^2 + x - 12x - 4 = 0$$
$$3x^2 - 11x - 4 = 0$$

Which is the required equation.

If  $S_1, S_2$  are the roots of required quadratic equation, then

Let  $x = S_1$  and  $x = S_2$   
 $\Rightarrow x - S_1 = 0, x - S_2 = 0$   
 $(x - S_1)(x - S_2) = 0$   
 $x^2 - (S_1 + S_2)x + S_1 S_2 = 0,$

$$S_1 + S_2 = \text{Sum of roots} = S \quad \text{and} \quad S_1 S_2 = \text{Product of roots} = P$$

Then,  $x^2 - Sx + P = 0 \dots \dots \dots \text{(i)}$

Above formula (i) is used to form a quadratic equation when its roots are given.

Let us solve some examples to understand the concept.

**Example:** Form a quadratic equation whose roots are  $\frac{1}{4}$  and  $-1$ .

**Solution:** The roots are  $\frac{1}{4}$  and  $-1$ .

Let  $x = \frac{1}{4}$  and  $x = -1$

$$x - \frac{1}{4} = 0 \quad \text{and} \quad x + 1 = 0$$

$$4x - 1 = 0, \quad x + 1 = 0$$

Factors of quadratic equation are  $(4x - 1)$  and  $(x + 1)$ , we have

$$(4x - 1)(x + 1) = 0$$

$$4x^2 + 4x - x - 1 = 0$$

$$4x^2 + 3x - 1 = 0, \text{ required quadratic equation.}$$

**Example:** Form a quadratic equation with roots:

- (i) 5 and 6      (ii)  $-\frac{5}{4}$  and  $\frac{16}{5}$       (iii)  $3+2i$  and  $3-2i$

**Solution:**

- (i) Since 5 and 6 are roots then

- (iii)  $3+2i$  and  $3-2i$

$$S = \text{sum of roots} = 5 + 6 = 11$$

$$P = \text{product of the roots} = 5(6) = 30$$

The required equation is

$$S = \text{Sum} = 3 + 2i + 3 - 2i = 6$$

$$P = \text{Product} = (3 + 2i)(3 - 2i)$$

$$= 9 - 4i^2 = 13$$

The equation is:

$$x^2 - 6x + 13 = 0$$

Substituting these values of S and P in (i), we have

$$x^2 - 11x + 30 = 0$$

$$(ii) \quad -\frac{5}{4} \text{ and } \frac{16}{5}$$

$$\text{Sum } = S = -\frac{5}{4} + \frac{16}{5} = \frac{13}{3}$$

$$\text{Product} = P = -\frac{5}{4} \times \frac{16}{5} = -4$$

The equation is:  $x^2 - \frac{13}{3}x + 4 = 0$

$$\text{or } 3x^2 - 13x + 12 = 0$$

**Example:** If  $S_1, S_2$  are the roots of  $ax^2+bx+c=0$ . Form the equation whose roots are twice to the roots of the given equation.

**Solution:** If  $S_1, S_2$  are the roots of  $ax^2 + bx + c = 0$  then

$$S_1 + S_2 = \frac{-b}{a} , \quad S_1 S_2 = \frac{c}{a}$$

The new roots are  $2S_1$  and  $2S_2$ .

$$\text{Sum of new roots} = S = 2S_1 + 2S_2$$

$$= 2(S_1 + S_2) = 2\left(\frac{-b}{a}\right) = \frac{-2b}{a}$$

$$\text{Product of new roots} = P = (2S_1)(2S_2) = 4S_1S_2$$

$$= 4 \left( \frac{c}{a} \right) = \frac{4c}{a}$$

Now substituting the values of  $S$  and  $P$  in  $x^2 - Sx + P \equiv 0$ , we get

$$x^2 - \left(\frac{-2b}{a}\right)x + \frac{4c}{a} = 0$$

$$\text{or } ax^2 + 2bx + 4c = 0$$

## Exercise 2.4

1. Find the sum and product of the roots of the quadratic equation.

  - $x^2 - 5x + 2 = 0$
  - $-4x^2 - 6x - 2 = 0$
  - $5x^2 - 2x + 2 = 0$
  - $-4x^2 - 8x - 9 = 0$
  - $16y^2 - 17y - 12 = 0$
  - $0.3x^2 - 7.7x + 1.8 = 0$

2. Form a quadratic equation with roots:

  - $1, -\frac{8}{3}$
  - $\sqrt{3}, 2\sqrt{3}$
  - $2+\sqrt{3}, 2-\sqrt{3}$
  - $5i, -5i$
  - $7+2i, 7-2i$

3. If  $S_1, S_2$  are the roots of  $3x^2 - 2x + 4 = 0$ ; find the value of:

  - $\frac{1}{S_1^2} + \frac{1}{S_2^2}$
  - $S_1^2 + S_2^2$
  - $2S_1 + 2S_2 + 4$
  - $\frac{1}{S_1} + \frac{1}{S_2}$
  - $\frac{S_1}{S_2} + \frac{S_2}{S_1}$
  - $S_1S_2^2 + S_1^2S_2$
  - $S_1^3S_2 + S_1S_2^3$
  - $(S_1 - 3)(S_2 - 3)$

4. if  $S_1, S_2$  are the roots of  $7x^2 + 10x + 7 = 0$ , form the equations whose roots are:

  - $S_1^2, S_2^2$
  - $\frac{1}{S_1}, \frac{1}{S_2}$
  - $S_1^3S_2, S_1S_2^3$
  - $S_1 - \frac{1}{S_1}, S_2 - \frac{1}{S_2}$
  - $2S_1 + 1, 2S_2 + 1$
  - $\frac{S_1}{S_2}, \frac{S_2}{S_1}$
  - $S_1 + S_2, \frac{1}{S_1} + \frac{1}{S_2}$
  - $S_1^2 + 1, S_2^2 + 1$
  - $S_1^2 + S_2, S_1 + S_2^2$

5. If  $S_1, S_2$  are the roots of  $x^2 + 6x + 3 = 0$ , form the equation whose roots are  $(S_1 + S_2)^2, (S_1 - S_2)^2$

6. If  $S_1, S_2$  are the roots of  $2x^2 + 6x - 3 = 0$ , form the equation whose roots are  $S_1 - \frac{3}{S_2^2}, S_2 - \frac{3}{S_1^2}$

7. Find  $k$  if  $S_1$  and  $S_1 - 5$  are the roots of  $x^2 - 3kx + 5 = 0$ .

8. Find  $k$  such that 3 is a root of  $x^2 + kx - 21 = 0$ .

## **Simultaneous Equations**

A system of equations having a common solution is called a system of simultaneous equations. The set of all the ordered pairs  $(x, y)$  which satisfies the system of equations is called the solution of the system. System of two equation involving two variables.

## **System of two Equations Involving two Variables**

### **Case I: When one Equation is Linear and One Quadratic**

If one of the equations is linear, we can find the value of one variable in terms of the other variable from linear equation. Substituting this value of one variable in the quadratic equation, we can solve it. The procedure is illustrated through the following example.

**Example:** Solve the system of equations  $x + y = 7$  and  $x^2 - xy + y^2 = 13$

**Solution:** Given Equations are

$$x + y = 7 \dots\dots\dots(i) \quad x^2 - xy + y^2 = 13 \dots\dots\dots(ii)$$

From equation (i)

$$x = 7 - y$$

Substituting in equation (ii)

$$(7 - y)^2 - (7 - y)y + y^2 = 13$$

$$49 - 14y + y^2 - 7y + y^2 + y^2 = 13$$

$$3y^2 - 21y + 36 = 0$$

$$y^2 - 7y + 12 = 0$$

$$y^2 - 4y - 3y + 12 = 0$$

$$y(y - 4) - 3(y - 4) = 0$$

$$(y - 4)(y - 3) = 0$$

$$y - 4 = 0 \text{ or } y - 3 = 0$$

$$y = 4, y = 3$$

Putting  $y = 3$ , in (i), we get  $x = 7 - 3 = 4$

Putting  $y = 4$ , in (i), we get  $x = 7 - 4 = 3$

Hence solution set is  $\{(4, 3), (3, 4)\}$

### Case II: When both of the Equations are Quadratic

The equations in this case are classified as:

- Both the equations contain only  $x^2$  and  $y^2$  terms.
- One of the equations, is homogeneous in  $x$  and  $y$ .
- Both the equations are non-homogeneous.

The methods of solving these types of equations are explained through the following examples.

**Example:** Solve the system of equations  $x^2 + y^2 = 25$  and  $2x^2 + 3y^2 = 66$

**Solution:** Given equations are

$$x^2 + y^2 = 25 \quad (i) \quad 2x^2 + 3y^2 = 66 \quad (ii)$$

Multiplying equation (i) by 2 and subtracting from equation (ii).

$$\begin{array}{r} 2x^2 + 3y^2 = 66 \\ \pm 2x^2 \pm 2y^2 = 50 \\ \hline y^2 = 16 \end{array}$$

Putting  $y^2 = 16$  in equation (i)

$$x^2 + 16 = 25$$

$$x^2 = 9$$

Now from  $x^2 = 9$ , we have  $x = \pm 3$

and from  $y^2 = 16$ , we have  $y = \pm 4$

Hence, solution set is  $\{(\pm 3, \pm 4)\}$

### Example:

Solve the system of equations  $x^2 + y^2 = 20$  and  $6x^2 + xy - y^2 = 0$

### Solution:

Given equations are

$$x^2 + y^2 = 20 \quad (\text{i})$$

$$6x^2 + xy - y^2 = 0 \quad (\text{ii})$$

From equation (ii)

$$-y^2 + xy + 6x^2 = 0$$

$$y^2 - xy - 6x^2 = 0$$

$$y^2 - 3xy + 2xy - 6x^2 = 0$$

$$y(y - 3x) + 2x(y - 3x) = 0$$

$$(y - 3x)(y + 2x) = 0$$

$$y - 3x = 0 \quad \text{or} \quad y + 2x = 0$$

$$y = 3x \dots\dots \text{(iii)} \quad y = -2x \dots\dots \text{(iv)}$$

Putting  $y = 3x$  in (i)

$$x^2 + (3x)^2 = 20$$

$$x^2 + 9x^2 = 20$$

$$10x^2 = 20$$

$$x^2 = 2$$

$$x = \pm\sqrt{2}$$

When

$$x = \sqrt{2}$$

From (iii)

$$y = 3\sqrt{2}$$

When

$$x = -\sqrt{2},$$

From (iii)

$$y = -3\sqrt{2}$$

Putting  $y = -2x$  in (i)

$$x^2 + (-2x)^2 = 20$$

$$x^2 + 4x^2 = 20$$

$$5x^2 = 20$$

$$x^2 = 4$$

$$x = \pm 2$$

$$x = 2$$

$$y = -4$$

$$x = -2$$

$$y = 4$$

Hence, solution set is  $\{(\sqrt{2}, 3\sqrt{2}), (-\sqrt{2}, -3\sqrt{2}), (2, -4), (-2, 4)\}$

Solve the following simultaneous equations:

1.  $2x + y = 1; \quad x^2 + y^2 = 10$
2.  $3x - 2y = 1; \quad x^2 + xy - y^2 = 1$
3.  $3x + y + 3 = 0; \quad (x+1)^2 - 4(x+1) - 6 = y$
4.  $(x+3)^2 - (y-2)^2 = 10; \quad x + y = 4$
5.  $2x^2 - 8xy + 6y^2 = 0; \quad x^2 + y^2 = 45$
6.  $6x^2 - 5xy - y^2 = 0; \quad y^2 + 4xy = 30$
7.  $4x^2 + 4y^2 = 65; \quad 6x - 2y = 5$
8.  $x^2 + y = 0; \quad x + y = -2$
9.  $x^2 + 3y^2 = 14; \quad 3x^2 + y^2 = 6$
10.  $2x^2 - 5y^2 = 8; \quad x^2 + 2y^2 = 13$

### Test Taking Tip:

Most standardized tests have a time limit, so you must budget your time carefully. Some questions will be much easier than others. If you cannot answer a question within a few minutes go on to the next one. If there is still time left when you get to the end of the test, go back to the ones that you skipped.

## Real Problems Involving Quadratic Equations

There are many problems which lead to quadratic equations. To form an equation, we consider symbols for unknown quantities in the problems.

In order to solve problems, we must have

- (i) Suppose the unknown quantities to be  $x$  or  $y$ .
- (ii) Translate the problem into symbols and form the equation satisfying the conditions.

The method of solving the problem will be illustrated through the following examples.

**Example:** The length of a room is 3 meters greater than its breadth. If the area of the room is 180 square meters, find length and breadth of the room.

**Solution:** Let the breadth of room =  $x$  metres

and the length of room =  $x + 3$  metres

Area of the room =  $x(x + 3)$  square metres

By the conditions of the equation

$$x(x + 3) = 180 \quad \text{or} \quad x^2 + 3x - 180 = 0$$

$$(x + 15)(x - 12) = 0 \quad \text{or} \quad x = -15, x = 12$$

As breadth cannot be negative, therefore, we take  $x = 12$ .

$$\text{Length} = x + 3 = 12 + 3 = 15$$

$\therefore$  Breadth of the room = 12m and length of room = 15m

**Example:**

The sum of the Cartesian coordinates of a point is 6 and the sum of their squares is 20. Find the coordinate of the point.

**Solution:**

Let  $(x, y)$  be the coordinates of required point, then by the given condition we get

$$x + y = 6 \quad (i)$$

$$x^2 + y^2 = 20 \quad (ii)$$

From(i)  $x + y = 6 \Rightarrow y = 6 - x$  putting in (ii)

$$x^2 + y^2 = 20 \Rightarrow x^2 + (6-x)^2 = 20$$

$$x^2 + 36 - 12x + x^2 = 20$$

$$x^2 - 6x + 8 = 0$$

$$x^2 - 4x - 2x + 8 = 0$$

$$x(x-4) - 2(x-4) = 0$$

$$(x-4)(x-2) = 0$$

$$x-4=0, \quad x-2=0$$

$$x=4, \quad x=2$$

When  $x=4, y=6-x=6-4=2$

When  $x=2, y=6-x=6-2=4$

$\therefore$  The coordinates of the point are 4 and 2, the points may be (4, 2) and (2, 4).

**Problem Solving Plan:**

- |                        |                         |
|------------------------|-------------------------|
| • Explore the problem. | • Plan the solution.    |
| • Solve the problem.   | • Examine the solution. |

**Exercise 2.6**

- Product of two consecutive positive even numbers is 120. Find the numbers.
- The difference of a positive number and its square is 380. Find the number.
- The difference of cubes of two consecutive positive numbers is 91. Find them.
- The sum of the Cartesian coordinates of a point is 9 and the sum of their squares is 45. Find the coordinates of the point.
- The sum of two numbers is 11 and the product is 30. Find the numbers.
- The sum of the squares of two consecutive positive odd integers is 34. Find the integers.
- The sum of ages of a father and his son is 50 years. Ten years ago, the father was 9 times as old as his son. Find the present age of father and son.
- A two-digit number is decreased by 45 when the digits are reversed. If the sum of the digits is 11, find the number.
- Sum of two numbers is 20. Find the numbers if the sum of first number and square of other is 40.
- The reciprocal of the sum of reciprocals of two numbers is  $\frac{12}{5}$ . Find the numbers if their sum is 10.

11. A group of 1025 students form two square patterns during morning assembly. One square pattern contains 5 more students than the other. Find the number of students in each pattern.
12. Sum of squares of two positive consecutive numbers is 145. The difference of their squares is 17. Find the numbers.
13. A ball is thrown upwards its height  $h(t)$  (in meters) after  $t$  seconds is modeled by  $d(t) = -5t^2 + 20t + 2$ . Find the time when the ball hit the ground.
14. The stopping distance  $d(x)$  (in meters) of a car traveling at  $x$  km/h is modeled by  $d(x) = 0.05x^2 + 0.4x$ . If the stopping distance is 30 meters, find the speed of the car.
15. A valuable stamp 4cm wide and 5cm long. The stamp is to be mounted on a sheet of paper that is  $5\frac{1}{2}$  times the area of the stamp. Determine the dimensions of the paper that will ensure a uniform border around the stamp.

### I have Learnt

- $ax^2 + bx + c = 0, a \neq 0$  is standard form of quadratic equation in one variable  $x$ .
- There are three methods for solving a quadratic equation.
  - i. Factorization Method    ii. Completing Square Method    iii. The Quadratic Formula
- Some equations are not quadratic but they can be reduced in quadratic form.
- In the quadratic formula, the expression under the radical sign ' $b^2 - 4ac$ ', is called the discriminant. The discriminant tells the nature of roots of quadratic equation.
- Sum of the roots of a quadratic equation =  $S = -\frac{b}{a} = -\frac{\text{coefficient of } x}{\text{coefficient of } x^2}$
- Product of roots of a quadratic equation =  $P = \frac{c}{a} = \frac{\text{constant term}}{\text{coefficient of } x^2}$
- Formula used to form a quadratic equation when its roots are given is  $x^2 - Sx + P = 0$
- A system of equations having a common solution is called a system of simultaneous equations.
- The set of all the ordered pairs  $(x, y)$  which satisfies the system of equations is called the solution of the system.

### MISCELLANEOUS EXERCISE-2

1. Encircle the correct option in the following.
- Which of the following is a quadratic equation?
    - $ax + b = c$
    - $ax^2 + bx + c$
    - $ax^2 + bx + c = 0, a \neq 0$
    - $ax^2 + bx + c = 0, a = 0$




## 03

## Matrices And Determinants

**After studying this unit students will be able to:**

- Display information in the form of matrices of order 2.
- Calculate product of scalar quantity and a matrix.
- Solve situations involving sum, difference and product of two matrices.
- Evaluate the determinant and inverse of a matrix of order 2 by 2.
- Solve the simultaneous linear equations in two variables using matrix inversion method and Cramer's rule.
- Explain with examples, how mathematics plays a key role in the development of new scientific theories and technologies.
- Apply concepts of matrices to real world problems.

Matrices are used for solving system of linear equations by several methods. These methods are used in computer programs, in traffic flow, schedule air line flights, engineering, accounting, economics etc



# Introduction of Matrices

In a mathematical quiz between girls and boys, Class 8 (girls) scored 95 points, class 8 (boys) scored 97 points, Class 9 (girls) scored 93 points, class 9 (boys) scored 90 points, class 10 (girls) scored 96 points and class 10 (boys) scored 94 points. This information can be displayed as under:

$$Q = \begin{bmatrix} 95 & 93 & 96 \\ 97 & 90 & 94 \end{bmatrix} \begin{matrix} \text{girls} \\ \text{boys} \end{matrix}$$

**8<sup>th</sup>    9<sup>th</sup>    10<sup>th</sup>**

This method is no doubt easy to write and manipulate. In 1850, Sylvester introduced such rectangular arrangement and he named it 'Matrix'. Later Hamilton and Cayley made further significant contributions to the *Matrix Algebra*. Today matrices are being used in almost every academic discipline in building models, organizing data and solving real world problems.

## Matrix (plural Matrices)


A Matrix is a rectangular arrangement of numbers.

e.g.  $\begin{bmatrix} 1 \end{bmatrix}$ ,  $\begin{bmatrix} 3 & 4 \\ 5 & 6 \end{bmatrix}$ ,  $\begin{bmatrix} 5 & 6 \end{bmatrix}$ ,  $\begin{bmatrix} 0 \end{bmatrix}$

### Check Point:

If  $A = [a]$  is a matrix, with only one row and only one column. Can it be called a row matrix or a column matrix?

## A Model Matrix



### Key Fact:

- The numbers used in a matrix are called 'elements' or 'entries' of the matrix.
- Elements of a matrix are written within the square brackets in a definite order, in rows and columns.
- Capital alphabets are used to name a matrix while elements of a matrix are usually represented by small alphabets and numbers.

Row: Horizontal arrangement of elements in a matrix is called a row.

$$\begin{bmatrix} 5 & 2 & 6 \\ 3 & 1 & 7 \end{bmatrix}$$

First row and second row of a matrix are represented by R<sub>1</sub> and R<sub>2</sub> respectively.

Column: Vertical arrangement of elements is in a matrix called a column.

$$\begin{bmatrix} 5 & 2 & 6 \\ 3 & 1 & 7 \end{bmatrix}$$

First column, second column and third column of a matrix are represented by C<sub>1</sub>, C<sub>2</sub>, C<sub>3</sub> respectively.

The number of rows and columns may be equal or unequal in any matrix, however the number of elements in different rows and in different columns remains the same. In  $\begin{bmatrix} 5 & 2 & 6 \\ 3 & 1 & 7 \end{bmatrix}$ , there are 2 rows and 3 columns. There are 3 elements in each row and 2 elements in each column.

### Order of Matrix

If 'A' is a matrix with 'm' number of rows and 'n' number of columns, then order of the matrix is m-by-n.

Order of  $\begin{bmatrix} 5 \\ 3 \end{bmatrix}$  is 2-by-1 i.e. 2 rows and 1 column.

### Equal Matrices

Two matrices are said to be equal if and only if their:

- (i) order is same
- (ii) corresponding elements are same.

### Example:

Check whether the following pairs of matrices are equal or not. Mention the reason in each case.

$$(i) \begin{bmatrix} \sqrt{4} \\ (-5)^2 \end{bmatrix} = \begin{bmatrix} 2 \\ 25 \end{bmatrix}$$

- Matrices on both sides of equality are of order 2-by-1
- Corresponding elements of the matrices on both sides of equality are also equal.  
Hence both matrices are equal.

$$(ii) A = \begin{bmatrix} 13 \\ 17 \end{bmatrix}, B = [13 \quad 17]$$

Order of matrix A is 2-by-1. Order of matrix B is 1-by-2. Order is different, we can say  $A \neq B$ .

$$(iii) E = \begin{bmatrix} 5 & 2 \\ 3 & 6 \end{bmatrix}, F = \begin{bmatrix} 5 & 2 \\ 3 & 3 \end{bmatrix}$$

Since the corresponding elements are not equal.  $\therefore E \neq F$ .

### Example:

Find unknowns from the following if possible.

$$\begin{bmatrix} x+2 & 5 \\ 3 & y-4 \end{bmatrix} = \begin{bmatrix} 9 & 5 \\ 3 & 7 \end{bmatrix}$$

Comparing the corresponding elements of two equal matrices, we get

$$x+2=9, 5=5, 3=3, y-4=7.$$

Now from  $x+2=9$ , we have  $x=7$  and from  $y-4=7$ , we have  $y=11$ .

### Key Fact:

Two matrices are equivalent if and only if they have same rank.

### Check Point:

Can you find the value of x from  $\begin{bmatrix} x & 0 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 3 \\ 3 & 0 \end{bmatrix}$ .

Mention the reason, if your answer is no.

## Types of Matrices

### Row and Column Matrices

A matrix having only one row is called a row matrix while a matrix having only one column is called a column matrix.

Row matrices:  $[5 \quad 3]$ ,  $[a+s+d]$ ,  $\begin{bmatrix} 5 & 6 \\ 3 & 7 \end{bmatrix}$

Column matrices:  $\begin{bmatrix} 5 \\ 6 \end{bmatrix}$ ,  $\begin{bmatrix} a+s \\ s+d \end{bmatrix}$ ,  $\begin{bmatrix} 5 \\ 6 \end{bmatrix}$ .

## Rectangular Matrix

A matrix with unequal number of rows and columns is called a rectangular matrix.

$$\begin{bmatrix} a+b & c \\ d+e & f \\ g+h & i \end{bmatrix}, \begin{bmatrix} \frac{1}{3} & \frac{4}{3} & \frac{7}{3} \\ \frac{2}{3} & \frac{5}{3} & \frac{8}{3} \end{bmatrix} \text{ and } \begin{bmatrix} 5 \\ 6 \end{bmatrix} \text{ are all rectangular matrices.}$$

## Square Matrix

A matrix with equal number of rows and columns is called a square matrix.

$$\begin{bmatrix} 5 & 3 \\ 1 & 2 \end{bmatrix}, \begin{bmatrix} 1 & 9 & 5 \\ 0 & 8 & 2 \\ 2 & 5 & -1 \end{bmatrix} \text{ and } [5] \text{ are all square matrices.}$$

## Null Matrix (Zero Matrix, Additive Identity Matrix)

A matrix of any order, having all the elements equal to zero, is called a null matrix.

A null matrix is represented by capital English alphabet 'O', while its elements are all zeros.

Null matrices may have any order.

$$O_2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, O_1 = [0], O_{12} = [0 \ 0] \text{ are all null matrices.}$$

## Diagonal Matrix

A square matrix in which every element except the primary (principal) diagonal elements is zero is called a diagonal matrix.

$$A = \begin{bmatrix} \sqrt{3} & 0 \\ 0 & 15 \end{bmatrix}, B = \begin{bmatrix} 0 & 0 \\ 0 & 5 \end{bmatrix}, C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 5 \end{bmatrix} \text{ are diagonal matrices but } D = \begin{bmatrix} 0 & 5 \\ 3 & 0 \end{bmatrix} \text{ is not a diagonal matrix.}$$

## Scalar Matrix

A diagonal matrix in which all the elements of primary diagonal are equal and non-zero is called a scalar matrix.

$P = \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$ , is a scalar matrix of order 2, but  $Q = \begin{bmatrix} 0 & 5 \\ 5 & 0 \end{bmatrix}$  is not a scalar matrix, since its primary diagonal elements are zeros and secondary diagonal elements are nonzero.

Some more examples of scalar matrices are:

$$\begin{bmatrix} 3 & 0 \\ 0 & \sqrt{9} \end{bmatrix}, \begin{bmatrix} 2 & 0 \\ 3-3 & 4 \div 2 \end{bmatrix}, \begin{bmatrix} 3+7 & 0 & 0 \\ 0 & \frac{20}{2} & 0 \\ 0 & 0 & 2 \times 5 \end{bmatrix}$$

## Unit Matrix (Multiplicative Identity Matrix)

A scalar matrix in which all the primary diagonal elements are equal to 1, is called a unit matrix. Unit matrix is represented by I.

$$I_1 = [1], I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

**Check Point:**

Write the most appropriate type of following matrices.

$$A = \begin{bmatrix} 15 & 0 \\ 0 & \frac{45}{3} \end{bmatrix}, \quad B = \begin{bmatrix} 0 & 3 \\ 3 & 0 \end{bmatrix}, \quad C = \begin{bmatrix} \frac{\sqrt{9}}{3} & 5 - 5 \\ \frac{5}{5} - 1 & \frac{15}{5 \times 3} \end{bmatrix}$$

**Transpose of a Matrix**

If rows (columns) of a matrix P are changed into columns (rows) then the resulting matrix is called transpose of the matrix P. It is represented by  $P^t$ .

**Example:**

If a matrix G represents number of gold medals and silver medals won by three friends, as under:

Gold medals	Silver medals	
$\begin{bmatrix} 5 & 3 \\ 2 & 6 \\ 1 & 7 \end{bmatrix}$	$\begin{array}{l} Azka \\ Aemen \\ Khansa \end{array}$	

then find  $G^t$ .

**Solution:**

Azka	Aemen	Khansa
$\begin{bmatrix} 5 & 2 & 1 \\ 3 & 6 & 7 \end{bmatrix}$	$\begin{array}{l} Gold \text{ medals} \\ Silver \text{ medals} \end{array}$	

**Key Fact:**

If order of a matrix P is m-by-n, then order of  $P^t$  is n-by-m.

**Symmetric Matrices and Skew Symmetric Matrices**

For a square matrix A

A is Symmetric if,  $A = A^t$ .A is Skew Symmetric if,  $A = -A^t$ .**Example:**

From the following, check for symmetric and skew symmetric matrices.

If  $S = \begin{bmatrix} 5 & 3 \\ 3 & 5 \end{bmatrix}$  then,  $S^t = \begin{bmatrix} 5 & 3 \\ 3 & 5 \end{bmatrix}$ .

As  $S = S^t$ , therefore S is a symmetric matrix.

If  $T = \begin{bmatrix} 5 & 2 \\ 6 & 11 \end{bmatrix}$  then,  $T^t = \begin{bmatrix} 5 & 6 \\ 2 & 11 \end{bmatrix}$

As  $T \neq T^t$ , therefore T is not a symmetric matrix.

If  $V = \begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}$

then,  $V^t = \begin{bmatrix} 0 & -2 \\ 2 & 0 \end{bmatrix} = -\begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix} = -V^t$

As  $V = -V^t$ , therefore V is a skew symmetric matrix.

**Check Point:**

If  $\begin{bmatrix} -3 & x \\ 4 & 2 \end{bmatrix}^t = \begin{bmatrix} -3 & 4 \\ 2 & 2 \end{bmatrix}$ ,

then find  $x = ?$

**Key Fact:**

- All the square matrices of order 1 are symmetric e.g. [9].
- $(A^t)^t = A$ .
- All the symmetric matrices are square but every square matrix is not necessarily symmetric.

**Exercise 3.1**

1. Write the number of rows, columns and order of the given matrices.

(a)  $[1 + 2 \quad 4 + 5]$  (b)  $\begin{bmatrix} a+s & 0 \\ 0 & d+s \end{bmatrix}$  (c)  $\begin{bmatrix} 5 \\ 7 \end{bmatrix}$  (d)  $[4 + 2 \times 1 + 5]$

2. Check whether the following pairs of matrices are equal or not.

(a)  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} \sqrt{25} - \sqrt{25} & \frac{\sqrt{25}}{5} \\ \frac{\sqrt{16}}{4} & \sqrt{16 - 16} \end{bmatrix}$  (b)  $[1 \ 2 \ 3], [1+2+3]$

(c)  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}, [1]$  (d)  $\begin{bmatrix} 1+0 & 0+0 \\ 0 \times 0 & 1-0 \end{bmatrix}, \begin{bmatrix} \frac{5 \times 5}{25} & \frac{3 \times 0}{7} \\ 2 - 17 + 15 & 17 \times \frac{1}{17} \end{bmatrix}$

3. Hyder, Hassan and Ahmed scored 7 points, 11 points and 10 points respectively in a mathematical quiz. Display the data in a row matrix (R) and a column matrix (C).

4. Write the most appropriate type of the following matrices.

(a)  $\begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix}$  (b)  $\begin{bmatrix} 0 & 0 \\ 0 & 3+0 \end{bmatrix}$  (c)  $\begin{bmatrix} 15 \times 3 \\ 9 \times 5 \end{bmatrix}$

(d)  $\begin{bmatrix} \frac{\sqrt{625}}{25} & 17 + 15 - 32 \\ 18 - 19 + 1 & \frac{3}{\sqrt{9}} \end{bmatrix}$  (e)  $\begin{bmatrix} 6+2 & 0 \\ -3+3 & 8 \end{bmatrix}$

5. Check for symmetric and skew symmetric matrices from the following.

(a)  $\begin{bmatrix} 5 & 2 & 3 \\ 2 & 9 & 6 \\ 3 & 6 & 3 \end{bmatrix}$  (b)  $\begin{bmatrix} 0 & 9 \\ -9 & 0 \end{bmatrix}$  (c)  $\begin{bmatrix} 0 & -2 & -4 \\ 2 & 0 & -2 \\ 4 & 2 & 0 \end{bmatrix}$

(d)  $\begin{bmatrix} 0 & -7 \\ 7 & 0 \end{bmatrix}$  (e)  $\begin{bmatrix} 0 & -5 \\ -5 & 0 \end{bmatrix}$

6. Write a symmetric and a skew symmetric matrix of order 3.

**Addition, Subtraction and Scalar Multiplication of Matrices****Addition and Subtraction of Matrices**

Two matrices can be added or subtracted only if their order is same.

Consider  $A = \begin{bmatrix} 5 \\ 3 \end{bmatrix}$ ,  $B = \begin{bmatrix} 5 & 3 \end{bmatrix}$ ,  $C = \begin{bmatrix} 6 \\ 3 \end{bmatrix}$ . Here addition and subtraction of A and B is impossible

because their order is not same.

However, addition and subtraction of A and C is possible as their order is same.

**Key Fact:**

- In the process of addition or subtraction of matrices, only corresponding elements are added or subtracted.
- Matrices of same order are conformable for addition and subtraction.

**Example :** If  $A = \begin{bmatrix} 2 & -3 \\ 6 & 5 \end{bmatrix}$  and  $B = \begin{bmatrix} 7 & 8 \\ 9 & -2 \end{bmatrix}$  then find  $A + B$ ,  $A - B$ ,  $B + A$ ,  $B - A$ .

$$\text{Solution: } A + B = \begin{bmatrix} 2 & -3 \\ 6 & 5 \end{bmatrix} + \begin{bmatrix} 7 & 8 \\ 9 & -2 \end{bmatrix} = \begin{bmatrix} 2 - 3 + 7 & 8 \\ 6 + 9 - 2 & 5 \end{bmatrix} = \begin{bmatrix} 7 & 8 \\ 13 & 3 \end{bmatrix}$$

$$A - B = \begin{bmatrix} 2 & -3 \\ 6 & 5 \end{bmatrix} - \begin{bmatrix} 7 & 8 \\ 9 & -2 \end{bmatrix} = \begin{bmatrix} 2 - 7 & -3 - 8 \\ 6 - 9 & 5 - (-2) \end{bmatrix} = \begin{bmatrix} -5 & -11 \\ -3 & 7 \end{bmatrix}$$

$$\text{Similarly, } B + A = \begin{bmatrix} 7 & 8 \\ 9 & -2 \end{bmatrix} + \begin{bmatrix} 2 & -3 \\ 6 & 5 \end{bmatrix} = \begin{bmatrix} 7 + 2 & 8 - 3 \\ 9 - 2 + 6 & -2 + 5 \end{bmatrix} = \begin{bmatrix} 9 & 5 \\ 13 & 3 \end{bmatrix}$$

$$B - A = \begin{bmatrix} 7 & 8 \\ 9 & -2 \end{bmatrix} - \begin{bmatrix} 2 & -3 \\ 6 & 5 \end{bmatrix} = \begin{bmatrix} 7 - 2 & 8 + 3 \\ 9 - 6 & -2 - 5 \end{bmatrix} = \begin{bmatrix} 5 & 11 \\ 3 & -7 \end{bmatrix}$$

**Check Point:**

Can you add  
 $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$  and  $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ ?

**Scalar Multiplication of a Matrix**

Scalar multiplication of a matrix means multiplication of a matrix with a constant.

If a matrix 'C' is multiplied by 2, then each of its element is doubled.

$$\text{If } C = \begin{bmatrix} 2 & 0 \\ 6 & 0 \end{bmatrix} \text{ then, } 2C = 2 \times \begin{bmatrix} 2 & 0 \\ 6 & 0 \end{bmatrix} = \begin{bmatrix} 2 \times 2 & 2 \times 0 \\ 2 \times 6 & 2 \times 0 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ 12 & 0 \end{bmatrix}$$

**Commutative Property of Matrix Addition**

If A and B are two matrices of same order, then  $A + B = B + A$ , is called commutative property of matrix addition.

**Example:**

If  $P = \begin{bmatrix} 5 & 3 \\ 2 & 9 \end{bmatrix}$  and  $Q = \begin{bmatrix} 7 & 2 \\ 3 & 9 \end{bmatrix}$ , then verify the commutative property of matrix addition.

**Solution:**

Commutative property of addition for above matrices is  $P + Q = Q + P$ .

$$\text{L.H.S : } P + Q = \begin{bmatrix} 5 & 3 \\ 2 & 9 \end{bmatrix} + \begin{bmatrix} 7 & 2 \\ 3 & 9 \end{bmatrix} = \begin{bmatrix} 5 + 7 & 3 + 2 \\ 2 + 3 & 9 + 9 \end{bmatrix} = \begin{bmatrix} 12 & 5 \\ 5 & 18 \end{bmatrix}$$

$$\text{R.H.S: } Q + P = \begin{bmatrix} 7 & 2 \\ 3 & 9 \end{bmatrix} + \begin{bmatrix} 5 & 3 \\ 2 & 9 \end{bmatrix} = \begin{bmatrix} 7 + 5 & 2 + 3 \\ 3 + 2 & 9 + 9 \end{bmatrix} = \begin{bmatrix} 12 & 5 \\ 5 & 18 \end{bmatrix}$$

As,  $P + Q = Q + P$ , therefore commutative property of matrix addition is verified.

**Check Point:**

Can you verify the commutative property of matrix addition for:

$$A = [x] \text{ and } B = [x \ y]$$

**Associative Property of Matrix Addition**

If A, B and C are three matrices of same order, then  $(A + B) + C = A + (B + C)$ , is called associative property of matrix addition.

**Example:** If  $R = \begin{bmatrix} 9 & 7 \\ 5 & 3 \\ 8 & 6 \end{bmatrix}$ ,  $S = \begin{bmatrix} 10 & 12 \\ 13 & 7 \\ 5 & 2 \end{bmatrix}$  and  $T = \begin{bmatrix} 1 & 5 \\ 6 & 7 \\ 5 & -4 \end{bmatrix}$ , then verify associative property of matrix addition.

**Solution:** Associative property of addition for matrices is  $(R+S)+T = R+(S+T)$ .

$$\text{L.H.S: } (R+S)+T = \left( \begin{bmatrix} 9 & 7 \\ 5 & 3 \\ 8 & 6 \end{bmatrix} + \begin{bmatrix} 10 & 12 \\ 13 & 7 \\ 5 & 2 \end{bmatrix} \right) + \begin{bmatrix} 1 & 5 \\ 6 & 7 \\ 5 & -4 \end{bmatrix} = \begin{bmatrix} 9+10 & 7+12 \\ 5+13 & 3+7 \\ 8+5 & 6+2 \end{bmatrix} + \begin{bmatrix} 1 & 5 \\ 6 & 7 \\ 5 & -4 \end{bmatrix}$$

$$= \begin{bmatrix} 19 & 19 \\ 18 & 10 \\ 13 & 8 \end{bmatrix} + \begin{bmatrix} 1 & 5 \\ 6 & 7 \\ 5 & -4 \end{bmatrix} = \begin{bmatrix} 19+1 & 19+5 \\ 18+6 & 10+7 \\ 13+5 & 8+(-4) \end{bmatrix} = \begin{bmatrix} 20 & 24 \\ 24 & 17 \\ 18 & 4 \end{bmatrix}$$

$$\text{R.H.S: } R+(S+T) = \begin{bmatrix} 9 & 7 \\ 5 & 3 \\ 8 & 6 \end{bmatrix} + \left( \begin{bmatrix} 10 & 12 \\ 13 & 7 \\ 5 & 2 \end{bmatrix} + \begin{bmatrix} 1 & 5 \\ 6 & 7 \\ 5 & -4 \end{bmatrix} \right)$$

$$= \begin{bmatrix} 9 & 7 \\ 5 & 3 \\ 8 & 6 \end{bmatrix} + \left( \begin{bmatrix} 10+1 & 12+5 \\ 13+6 & 7+7 \\ 5+5 & 2+(-4) \end{bmatrix} \right)$$

$$= \begin{bmatrix} 9 & 7 \\ 5 & 3 \\ 8 & 6 \end{bmatrix} + \begin{bmatrix} 11 & 17 \\ 19 & 14 \\ 10 & -2 \end{bmatrix} = \begin{bmatrix} 9+11 & 7+17 \\ 5+19 & 3+14 \\ 8+10 & 6-2 \end{bmatrix} = \begin{bmatrix} 20 & 24 \\ 24 & 17 \\ 18 & 4 \end{bmatrix}$$

As, L.H.S = R.H.S, therefore associative property of matrix addition is verified.

### Additive Identity in Matrices

Additive identity is such a matrix which causes no change in any matrix A while 'adding to' or 'subtracting from' it. Any matrix and its additive identity matrix have the same order.

If  $A = \begin{bmatrix} 3 \\ 7 \\ 2 \end{bmatrix}$ , then  $O = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$  is the additive identity of A.

It is observed that:  $A + O = O + A = A$

If  $B = [\sqrt{7}]$ , then the associated additive identity matrix will be  $O_1 = [0]$ .

#### Check Point:

Taking matrices A and B of same order, verify that:

- (i)  $k(A+B) = kA + kB$
- (ii)  $(h+k)A = hA + kA$
- (iii)  $(hk)B = h(kB)$

where h and k are constants.

### Additive Inverse of a Matrix

Additive inverse of a matrix 'A' is such a matrix which when added to A, gives additive identity matrix of the same order.

**Example (a):** If  $X = \begin{bmatrix} 5 & 0 \\ -9 & 4 \end{bmatrix}$  then find ' $-X$ '.

**(b):** Find T from  $5 \times \begin{bmatrix} 3 & 2 \\ 1 & 0 \end{bmatrix} - 2T = \begin{bmatrix} 5 & 4 \\ 3 & 2 \end{bmatrix}$

**Solution (a):** If  $X = \begin{bmatrix} 5 & 0 \\ -9 & 4 \end{bmatrix}$  then  $-X = \begin{bmatrix} -5 & 0 \\ 9 & -4 \end{bmatrix}$

$$\text{(b)} \quad 5 \times \begin{bmatrix} 3 & 2 \\ 1 & 0 \end{bmatrix} - 2T = \begin{bmatrix} 5 & 4 \\ 3 & 2 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 5 \times 3 & 5 \times 2 \\ 5 \times 1 & 5 \times 0 \end{bmatrix} - 2T = \begin{bmatrix} 5 & 4 \\ 3 & 2 \end{bmatrix}$$

#### Key Fact:

- Matrices A and B are called additive inverses (negative) of each other if  $A+B=O$ .
- Additive inverse of a matrix has the same order as that of the matrix.

$$\Rightarrow \begin{bmatrix} 15 & 10 \\ 5 & 0 \end{bmatrix} - 2T = \begin{bmatrix} 5 & 4 \\ 3 & 2 \end{bmatrix} \Rightarrow -2T = \begin{bmatrix} 5 & 4 \\ 3 & 2 \end{bmatrix} - \begin{bmatrix} 15 & 10 \\ 5 & 0 \end{bmatrix}$$

$$\Rightarrow -2T = \begin{bmatrix} 5-15 & 4-10 \\ 3-5 & 2-0 \end{bmatrix} = \begin{bmatrix} -10 & -6 \\ -2 & 2 \end{bmatrix}$$

$$\Rightarrow T = \frac{-1}{2} \begin{bmatrix} -10 & -6 \\ -2 & 2 \end{bmatrix} = \begin{bmatrix} \frac{-10}{2} & \frac{-6}{2} \\ \frac{-2}{2} & \frac{2}{2} \end{bmatrix} = \begin{bmatrix} 5 & 3 \\ 1 & -1 \end{bmatrix}$$

## Multiplication of Matrices

Two matrices A and B are conformable for product AB, if:

number of columns in A = number of rows in B

**Example:**

If  $A = \begin{bmatrix} 5 & 3 \\ 1 & -2 \end{bmatrix}$  and  $B = \begin{bmatrix} 6 \\ 4 \end{bmatrix}$ , then find AB and BA, if possible.


**Solution:**

**Finding AB**


As number of columns in A = 2 = number of rows in B. So, product AB is possible.

$$AB = \begin{bmatrix} 5 & 3 \\ 1 & -2 \end{bmatrix} \begin{bmatrix} 6 \\ 4 \end{bmatrix} = \begin{bmatrix} 5 \times 6 + 3 \times 4 \\ 1 \times 6 + (-2) \times 4 \end{bmatrix} = \begin{bmatrix} 30 + 12 \\ 6 - 8 \end{bmatrix} = \begin{bmatrix} 42 \\ -2 \end{bmatrix}.$$

Here



**Finding the sum of products of corresponding elements of R<sub>1</sub> of A and C<sub>1</sub> of B.**



**Finding the sum of products of corresponding elements of R<sub>2</sub> of A and C<sub>1</sub> of B.**

**Finding BA**

Since number of columns of B = 1, but number of rows in A = 2.

So, product BA is not possible.

### Key Fact:

- In multiplication, the first matrix contributes its number of rows and the second matrix contributes its number of columns, in the resultant matrix. i.e. If order of matrix A is m-by-n and order of matrix B is n-by-p, then order of AB is m-by-p.
- If A and B are two matrices which are conformable for product AB then B and A are not necessarily confirmable for product BA.

**Example:**

Check whether the following pairs of matrices are conformable for multiplication or not?

Mention order of product if possible.

(i)  $A = \begin{bmatrix} 5 \\ 3 \end{bmatrix}$  and  $B = \begin{bmatrix} 9 \\ 3 \end{bmatrix}$       (ii)  $E = \begin{bmatrix} 6 \\ 3 \end{bmatrix}^t$  and  $F = \begin{bmatrix} 6 \\ 3 \end{bmatrix}$

**Solution:**

(i)  $A = \begin{bmatrix} 5 \\ 3 \end{bmatrix}$  and  $B = \begin{bmatrix} 9 \\ 3 \end{bmatrix}$

Here, number of columns in  $B \neq$  number of rows in  $A$ , so product  $AB$  is not possible.

(ii)  $E = \begin{bmatrix} 6 \\ 3 \end{bmatrix}^t = [6 \ 3]$  and  $F = \begin{bmatrix} 6 \\ 3 \end{bmatrix}$

Here, number of columns of  $E =$  number of rows of  $F$ . So, product  $EF$  is possible.

Order of  $E$  is 1-by-2 and order of  $F$  is 2-by-1, so order of  $EF$  is 1-by-1.

**Example:** If  $A = \begin{bmatrix} 5 & 3 \\ 2 & 6 \end{bmatrix}$  and  $B = \begin{bmatrix} 7 & 2 \\ 4 & 9 \end{bmatrix}$  then find  $AB$  and  $BA$ . Also check whether  $AB = BA$  or not. What is concluded from the result?

**Solution:**  $AB = \begin{bmatrix} 5 & 3 \\ 2 & 6 \end{bmatrix} \times \begin{bmatrix} 7 & 2 \\ 4 & 9 \end{bmatrix}$

$$= \begin{bmatrix} (5 \times 7) + (3 \times 4) & (5 \times 2) + (3 \times 9) \\ (2 \times 7) + (6 \times 4) & (2 \times 2) + (6 \times 9) \end{bmatrix}$$

$$= \begin{bmatrix} 35 + 12 & 10 + 27 \\ 14 + 24 & 4 + 54 \end{bmatrix} = \begin{bmatrix} 47 & 37 \\ 38 & 58 \end{bmatrix}$$

$$BA = \begin{bmatrix} 7 & 2 \\ 4 & 9 \end{bmatrix} \times \begin{bmatrix} 5 & 3 \\ 2 & 6 \end{bmatrix}$$

$$= \begin{bmatrix} 7 \times 5 + 2 \times 2 & 7 \times 3 + 2 \times 6 \\ 4 \times 5 + 9 \times 2 & 4 \times 3 + 9 \times 6 \end{bmatrix}$$

$$= \begin{bmatrix} 35 + 4 & 21 + 12 \\ 20 + 18 & 12 + 54 \end{bmatrix} = \begin{bmatrix} 39 & 33 \\ 38 & 66 \end{bmatrix}$$

We see that  $AB \neq BA$ .

From this result it is concluded that:

Commutative property does not hold in matrix multiplication, in general.

**Key Fact:**

If  $A$  and  $B$  are diagonal matrices, then  $C = AB$  is diagonal.

**Associative Property of Matrix Multiplication**

Associative property of multiplication holds in matrices. If  $A$ ,  $B$ , and  $C$  are matrices and products  $AB$  and  $BC$  are possible, then:

$$(AB)C = A(BC)$$

**Example:** Verify Associative property of matrix multiplication for:

$$X = \begin{bmatrix} 5 & 3 \\ 2 & 6 \end{bmatrix}, \quad Y = \begin{bmatrix} 1 & -7 \\ 5 & 4 \end{bmatrix}, \quad Z = \begin{bmatrix} 6 \\ 4 \end{bmatrix}$$

**Solution:** Associative property of matrix multiplication is:  $(XY)Z = X(YZ)$

$$\text{L.H.S: } (XY)Z = \left( \left[ \begin{bmatrix} 5 & 3 \\ 2 & 6 \end{bmatrix} \times \begin{bmatrix} 1 & -7 \\ 5 & 4 \end{bmatrix} \right] \right) \times \begin{bmatrix} 6 \\ 4 \end{bmatrix} = \left[ \begin{bmatrix} 5 \times 1 + 3 \times 5 & 5 \times (-7) + 3 \times 4 \\ 2 \times 1 + 6 \times 5 & 2 \times (-7) + 6 \times 4 \end{bmatrix} \right] \begin{bmatrix} 6 \\ 4 \end{bmatrix}$$

$$= \begin{bmatrix} 20 & -23 \\ 32 & 10 \end{bmatrix} \begin{bmatrix} 6 \\ 4 \end{bmatrix} = \begin{bmatrix} 20 \times 6 - 23 \times 4 \\ 32 \times 6 + 10 \times 4 \end{bmatrix} = \begin{bmatrix} 120 - 92 \\ 192 + 40 \end{bmatrix} = \begin{bmatrix} 28 \\ 232 \end{bmatrix}$$

$$\text{R.H.S: } X(YZ) = \begin{bmatrix} 5 & 3 \\ 2 & 6 \end{bmatrix} \times \left( \begin{bmatrix} 1 & -7 \\ 5 & 4 \end{bmatrix} \times \begin{bmatrix} 6 \\ 4 \end{bmatrix} \right) = \begin{bmatrix} 5 & 3 \\ 2 & 6 \end{bmatrix} \left[ \begin{array}{l} 1 \times 6 + (-7) \times 4 \\ 5 \times 6 + 4 \times 4 \end{array} \right] \\ = \begin{bmatrix} 5 & 3 \\ 2 & 6 \end{bmatrix} \times \begin{bmatrix} 6 - 28 \\ 30 + 16 \end{bmatrix} = \begin{bmatrix} 5 & 3 \\ 2 & 6 \end{bmatrix} \times \begin{bmatrix} -22 \\ 46 \end{bmatrix} = \begin{bmatrix} 5 \times (-22) + 3 \times 46 \\ 2 \times (-22) + 6 \times 46 \end{bmatrix} = \begin{bmatrix} 28 \\ 232 \end{bmatrix}$$

As L.H.S = R.H.S, so associative property of matrix multiplication is verified.

### Distributive Property of Multiplication over Addition / Subtraction

- $A(B + C) = AB + AC$  (left distributive property of multiplication over addition)
- $A(B - C) = AB - AC$  (left distributive property of multiplication over subtraction)

**Example (a):** Verify  $A(B + C) = AB + AC$

$$A = \begin{bmatrix} 5 & 2 \end{bmatrix}, B = \begin{bmatrix} 6 \\ 4 \end{bmatrix} \text{ and } C = \begin{bmatrix} 1 \\ 2 \end{bmatrix}.$$

**(b):** Verify  $A(B - C) = AB - AC$  if  $A = \begin{bmatrix} 4 \\ 8 \end{bmatrix}$ ,  $B = [5]$ ,  $C = [7]$ .

**Solution:**

$$\text{(a) L.H.S: } A(B + C) = \begin{bmatrix} 5 & 2 \end{bmatrix} \left( \begin{bmatrix} 6 \\ 4 \end{bmatrix} + \begin{bmatrix} 1 \\ 2 \end{bmatrix} \right) = \begin{bmatrix} 5 & 2 \end{bmatrix} \begin{bmatrix} 6+1 \\ 4+2 \end{bmatrix} = \begin{bmatrix} 5 & 2 \end{bmatrix} \begin{bmatrix} 7 \\ 6 \end{bmatrix} \\ = [5 \times 7 + 2 \times 6] = [35 + 12] = [47]$$

$$\text{R.H.S: } AB + AC = \begin{bmatrix} 5 & 2 \end{bmatrix} \begin{bmatrix} 6 \\ 4 \end{bmatrix} + \begin{bmatrix} 5 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = [5 \times 6 + 2 \times 4] + [5 \times 1 + 2 \times 2] \\ = [30 + 8] + [5 + 4] = [38 + 9] = [47]$$

Comparing both the results, it is concluded that L.H.S = R.H.S

∴ Left distributive property of multiplication over addition is verified.

$$\text{(b) L.H.S: } A(B - C) = \begin{bmatrix} 4 \\ 8 \end{bmatrix} ([5] - [7]) = \begin{bmatrix} 4 \\ 8 \end{bmatrix} [5 - 7] = \begin{bmatrix} 4 \\ 8 \end{bmatrix} [-2] \\ = \begin{bmatrix} 4 \times (-2) \\ 8 \times (-2) \end{bmatrix} = \begin{bmatrix} -8 \\ -16 \end{bmatrix}$$

$$\text{R.H.S: } AB - AC = \begin{bmatrix} 4 \\ 8 \end{bmatrix} [5] - \begin{bmatrix} 4 \\ 8 \end{bmatrix} [7] = \begin{bmatrix} 4 \times 5 \\ 8 \times 5 \end{bmatrix} - \begin{bmatrix} 4 \times 7 \\ 8 \times 7 \end{bmatrix} \\ = \begin{bmatrix} 20 \\ 40 \end{bmatrix} - \begin{bmatrix} 28 \\ 56 \end{bmatrix} = \begin{bmatrix} -8 \\ -16 \end{bmatrix}$$

From both the results, it is concluded that L.H.S = R.H.S

∴ Left distributive property of multiplication over subtraction is verified.

#### Key Fact:

$(A + B)C = AC + BC$  (Right distributive property of multiplication over addition)

$(A - B)C = AC - BC$  (Right distributive property of multiplication over subtraction)

## Multiplicative Identity in Matrices

Multiplicative identity is such a matrix which causes no change in any matrix A when multiplied with A.

### Key Fact:

- Multiplicative identity matrices of orders 1, 2, 3, ...n are represented by  $I_1, I_2, I_3, \dots, I_n$  or simply by I.
- If I is multiplicative identity matrix of square matrix 'S', then  $IS = SI = S$ .
- $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix}, [1 \ 1]$  etc. are not multiplicative identities.
- Since multiplicative identity matrix can never be a rectangular matrix, so multiplicative identity matrices are only possible for square matrices.
- Commutative property of matrix multiplication can hold in particular cases:  
(i)  $A \times I = I \times A = A$  (if A and I are conformable for either multiplication.)  
(ii)  $B \times O = O \times B = O$  (if O and B are conformable for either multiplication.)

### Exercise 3.2

1. Find x, y, z, from the followings, if possible. Mention the reason if not possible.

$$(a) \begin{bmatrix} x & 9 \end{bmatrix} = \begin{bmatrix} 2 & y \end{bmatrix} \quad (b) \begin{bmatrix} 6 & 3 \\ 4 & x \end{bmatrix} = \begin{bmatrix} 7 & 3 \\ 4 & 0 \end{bmatrix} \quad (c) \begin{bmatrix} 5x \\ 2y \end{bmatrix} = \begin{bmatrix} -10 & 20 \end{bmatrix}$$

$$(d) -\begin{bmatrix} 2x & 3y \\ 4z & 10 \end{bmatrix} = \begin{bmatrix} 8 & 6 \\ 32 & -10 \end{bmatrix} \quad (e) \begin{bmatrix} x & -2y \\ 6 & x+y \end{bmatrix} = \begin{bmatrix} 3 & -6 \\ 6 & z \end{bmatrix} \quad (f) \begin{bmatrix} 5 \\ 6 \\ 7 \end{bmatrix} = \begin{bmatrix} 5 & 0 \\ x & 0 \\ 7 & 0 \end{bmatrix}$$

$$(g) \begin{bmatrix} x+y \\ x-y \end{bmatrix} = \begin{bmatrix} 11 \\ 1 \end{bmatrix} \quad (h) \begin{bmatrix} 5 & 10 \\ 15 & x \end{bmatrix} + \begin{bmatrix} 5 & y \\ 15 & 5 \end{bmatrix} = \begin{bmatrix} z & 15 \\ 30 & 7 \end{bmatrix}$$

$$(i) 5 \begin{bmatrix} x \\ 3y \end{bmatrix} - \begin{bmatrix} 36 \\ 26 \end{bmatrix} = 2 \begin{bmatrix} -2x \\ y \end{bmatrix} \quad (j) \begin{bmatrix} x & y & z \\ -2 & -4 & 5 \end{bmatrix} + 2 \times \begin{bmatrix} 5 & 3 & 2 \\ 1 & 6 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 8 & 11 \end{bmatrix}$$

2. Find the additive inverses of the following.

$$R = \begin{bmatrix} 5 & 0 & 3 \\ 7 & -9 & -1 \\ -8 & 5 & 6 \end{bmatrix}, S = \begin{bmatrix} -5 & 2 \\ 3 & -6 \\ -9 & 4 \end{bmatrix}, T = [5 \ -6 \ 1].$$

3. If  $A = \begin{bmatrix} 5 & 3 \\ -2 & 6 \end{bmatrix}, B = \begin{bmatrix} 10 & 8 \\ -8 & 6 \end{bmatrix}, C = \begin{bmatrix} 15 & 6 \\ 0 & -12 \end{bmatrix}$ , then find

$$(i) 2A + \frac{1}{2}B - \frac{1}{3}C \quad (ii) A - \frac{1}{2}B$$

4. If  $A = \begin{bmatrix} 1 & 2 \\ 3 & -3 \end{bmatrix}, B = \begin{bmatrix} 1 & -2 \\ 3 & 3 \end{bmatrix}, C = \begin{bmatrix} 2 & 0 \\ 6 & 0 \end{bmatrix}, D = \begin{bmatrix} 2 \\ 6 \end{bmatrix}, E = \begin{bmatrix} 2 & 2 \\ 6 & 6 \end{bmatrix}$   
Then check whether: (i)  $A + B = C$  (ii)  $C + D = E$  (iii)  $D + D = E$  (iv)  $E - D = C$

5. Taking matrices A and B from Q.3, verify commutative property of matrix addition.

6. Taking matrices A, B and C from Q. 3, verify associative property of matrix addition.

7. Taking matrices A and B from Q. 4, verify that  $A^t + B^t = (A + B)^t$ .

8. Find the matrix 'Z' from these equations.

$$(i) 4 \begin{bmatrix} 5 \\ 10 \end{bmatrix} - 5Z = \sqrt{5} \begin{bmatrix} \sqrt{45} \\ \sqrt{5} \end{bmatrix} \quad (ii) Z + \begin{bmatrix} 5 \\ -7 \end{bmatrix} = 3Z - \begin{bmatrix} 5 \\ 3 \end{bmatrix}.$$

9. (a) Mention the order of the indicated products where possible.

$$(i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \times [5 \ 3] \quad (ii) [5 \ 3] \times \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \quad (iii) \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \times \begin{bmatrix} 5 \\ 3 \end{bmatrix}$$

$$(iv) \begin{bmatrix} 5 \\ 3 \end{bmatrix} \times \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \quad (v) [5] \times \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \quad (vi) \begin{bmatrix} 1 & -2 \\ 3 & 4 \end{bmatrix} \times [5]$$

$$(vii) \begin{bmatrix} 1 \\ 3 \end{bmatrix} \times [5] \quad (viii) [5] \times \begin{bmatrix} 1 \\ 3 \end{bmatrix} \quad (ix) [5] \times [9 \ 3]$$

$$(x) [9 \ -3] \times [5] \quad (xi) [5] \times [10] \quad (xii) [5 \ 10] \times \begin{bmatrix} -2 \\ 3 \end{bmatrix}$$

$$(xiii) \begin{bmatrix} 2 \\ 3 \end{bmatrix} \times [5 \ 10] \quad (xiv) \begin{bmatrix} 5 & 2 \\ 3 & 4 \end{bmatrix} \times \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad (xv) \begin{bmatrix} 5 & 3 \\ 2 & 6 \end{bmatrix} \times [5 \ -2]$$

(b) Perform the indicated products in part (a) where possible.

10. If  $A = \begin{bmatrix} 1 & 0 \\ 2 & 3 \end{bmatrix}$ ,  $B = \begin{bmatrix} 2 & 1 \\ 0 & -3 \end{bmatrix}$  and  $C = \begin{bmatrix} 3 & 0 \\ -2 & 1 \end{bmatrix}$  then

- (i) Find  $AB$  and  $BA$ , check whether  $AB = BA$  or not?
- (ii) Find  $AC$  and  $CA$ , check whether  $AC = CA$  or not?
- (iii) Verify  $A(B + C) = AB + AC$
- (iv) Verify  $(A - B)C = AC - BC$
- (v) Verify the associative property of matrix multiplication.
- (vi) Find  $A^2$ ,  $B^2$ ,  $A + B$ ,  $A - B$ ,  $(A + B)(A - B)$ ,  $A^2 - B^2$ .
- (vii) Check whether  $(A + B) \times (A - B) = A^2 - B^2$  or not?
- (viii) Check whether  $(A - B) \times (A + B) = A^2 - B^2$  or not?
- (ix) Check whether  $(A + B) \times (A - B) = (A - B) \times (A + B)$  or not?
- (x) Verify that  $(AB)^t = B^t A^t$  and  $(A^t)^t = A$

## Multiplicative Inverse of a Matrix

Before we discuss inverse of a matrix, we need to examine the determinant of a matrix.

### Determinant of a Matrix

When a matrix has  $m$  rows and  $n$  columns, it is called an "m-by-n" matrix and is also called dimensions of a matrix. If a matrix has the same number of rows and columns, it is called a square matrix. With every square matrix is associated a number called its determinant, defined as follows for 2-by-2 matrices:

The determinant of its matrix  $\begin{bmatrix} a & c \\ b & d \end{bmatrix}$  is denoted  $\begin{vmatrix} a & c \\ b & d \end{vmatrix}$  and is defined as follows:

$$\begin{vmatrix} a & c \\ b & d \end{vmatrix} = ad - bc.$$

**Example:**

Find  $|A|$ , when  $A = \begin{vmatrix} 5 & 6 \\ 2 & 3 \end{vmatrix}$ .

$$\begin{aligned}\text{Det } A &= |A| = \begin{vmatrix} 5 & 6 \\ 2 & 3 \end{vmatrix} \\ &= (5 \times 3) - (6 \times 2) \\ &= 15 - 12 = 3\end{aligned}$$

**HISTORY:**

The theory of determinants is attributed to a German, Gottfried Wilhelm Leibiz. His work expanded upon the earlier work of Japanese mathematician Seki Kowa.

**Key Fact:**

- Every square matrix has a determinant either zero or nonzero.
- Determinants can be represented by  $D_1, D_2, D_3, \dots$
- The determinant of a matrix is not a matrix rather it is a number.
- While writing determinant of a matrix, the elements of the matrix should be written inside the vertical bars instead of brackets used for matrices.

**Example:** Evaluate the determinants

$$D_1 = \begin{vmatrix} -4 & 8 \\ 1 & -2 \end{vmatrix} = (-4 \times -2) - (8 \times 1) = 8 - 8 = 0$$

$$D_2 = \begin{vmatrix} \sqrt{5} & \sqrt{4} \\ \sqrt{1} & -\sqrt{5} \end{vmatrix} = (\sqrt{5} \times -\sqrt{5}) - (\sqrt{1} \times \sqrt{4}) = -(\sqrt{5})^2 - \sqrt{4} = -5 - 2 = -7$$

**Singular and Non-singular Matrices**

Given that A is a square matrix, then

A is singular matrix, if  $|A| = 0$

A is nonsingular matrix, if  $|A| \neq 0$

**Example:**

From the followings, check for the singular and non-singular matrices.

$$A = \begin{bmatrix} 5 & 10 \\ 3 & 6 \end{bmatrix}, B = \begin{bmatrix} 5 & 5 \\ 3 & \sqrt{5} \end{bmatrix}$$

$$\text{Solution: } |A| = \begin{vmatrix} 5 & 10 \\ 3 & 6 \end{vmatrix} = 5 \times 6 - 3 \times 10 = 30 - 30 = 0$$

As  $|A| = 0$ , therefore A is a singular matrix.

$$|B| = \begin{vmatrix} 7 & 5 \\ 3 & \sqrt{5} \end{vmatrix} = 7\sqrt{5} - 15 \neq 0.$$

As  $|B| \neq 0$ , therefore B is non-singular matrix.

**Check Point:**

Can you find determinant of a rectangular matrix?

**Adjoint of a Matrix**

Adjoint of a square matrix A (of order 2) is a square matrix obtained by:

- interchanging the positions of primary diagonal elements of A.
- changing the signs of secondary diagonal elements of A.

Adjoint of matrix A is represented by adj A.

$$\text{If } A = \begin{bmatrix} 5 & -3 \\ 4 & -2 \end{bmatrix}$$

Interchange the position  
of primary diagonal  
elements

$$\begin{bmatrix} -2 & -3 \\ 4 & 5 \end{bmatrix}$$

Change the signs of  
secondary diagonal  
elements

$$\begin{bmatrix} -2 & 3 \\ -4 & 5 \end{bmatrix} = \text{adj} A$$

### Finding Multiplicative Inverse of a Matrix

Any matrix  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ , will have an inverse  $A^{-1}$  if and only  $\begin{vmatrix} a & c \\ b & d \end{vmatrix} \neq 0$  mean A is non singular.

Then  $A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$ . We can also write:  $A^{-1} = \frac{1}{ad-bc} \text{Adjoint of } A$ .

**Example:** If  $A = \begin{bmatrix} 5 & 4 \\ 2 & 6 \end{bmatrix}$ , find  $A^{-1}$ .

**Solution:**  $A = \begin{bmatrix} 5 & 4 \\ 2 & 6 \end{bmatrix}$ , Compute the value of the determinant A.

$$|A| = \begin{vmatrix} 5 & 4 \\ 2 & 6 \end{vmatrix} = 5 \times 6 - 4 \times 2 = 30 - 8 = 22,$$

Since the determinant does not equal 0, therefore  $A^{-1}$  exists.

$A^{-1} = \frac{1}{22} \begin{bmatrix} 6 & -4 \\ -2 & 5 \end{bmatrix}$ , use formula:  $A^{-1} = \frac{1}{ad-bc} \text{Adjoint of } A$ .

This can be written as:  $A^{-1} = \begin{bmatrix} \frac{6}{22} & \frac{-4}{22} \\ \frac{-2}{22} & \frac{5}{22} \end{bmatrix} = \begin{bmatrix} \frac{3}{11} & \frac{-2}{11} \\ \frac{-1}{11} & \frac{5}{22} \end{bmatrix}$

#### Key Fact:

- Inverse of a square matrix, if exists, is unique and  $(A^{-1})^{-1} = A$ .
- A matrix A is invertible if its inverse exists.
- $A^{-1}A = I$ , I is identity matrix.

**Example:** If  $B = \begin{bmatrix} 5 & 2 \\ 6 & 3 \end{bmatrix}$ , find  $B^{-1}$  also verify that  $BB^{-1} = B^{-1}B = I$ .

**Solution:** If  $B = \begin{bmatrix} 5 & 2 \\ 6 & 3 \end{bmatrix}$ , then  $|B| = 5 \times 3 - 2 \times 6 = 15 - 12 = 3 \neq 0$ ,  $B^{-1}$  exists.

Adjoint of  $B = \begin{bmatrix} 3 & -2 \\ -6 & 5 \end{bmatrix}$ , apply formula:  $B^{-1} = \frac{1}{|B|} \text{Adjoint of } B$

$$B^{-1} = \frac{1}{3} \begin{bmatrix} 3 & -2 \\ -6 & 5 \end{bmatrix} = \begin{bmatrix} \frac{1}{3} & -\frac{2}{3} \\ -2 & \frac{5}{3} \end{bmatrix} = \begin{bmatrix} 1 & -\frac{2}{3} \\ -2 & \frac{5}{3} \end{bmatrix}$$

### Key Fact:

If A and B are square matrices, then  $|AB| = |A| \times |B|$ .

$$\text{Check: } B^{-1}B = \frac{1}{3} \begin{bmatrix} 3 & -2 \\ -6 & 5 \end{bmatrix} \begin{bmatrix} 5 & 2 \\ 6 & 3 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 15-12 & 6-6 \\ -30+30 & -12+15 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

$$\text{Similarly } BB^{-1} = \frac{1}{3} \begin{bmatrix} 5 & 2 \\ 6 & 3 \end{bmatrix} \begin{bmatrix} 3 & -2 \\ -6 & 5 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 15-12 & 6-6 \\ -30+30 & -12+15 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I$$

### Check Point:

Verify:  $(AB)^{-1} = B^{-1}A^{-1}$ , by taking A and B non singular matrix.

## Solution of Simultaneous Linear Equations

A large-scale use of matrices in almost every field is in the form of solution of simultaneous linear equations. While studying Economics, Statistics, Medical Sciences, Engineering etc, one has to find the solution of linear equations in two or more variables. However, our study at this level is confined to the solution of simultaneous linear equations only in two variables.

### Conversion of Matrix Equation into System of Linear Equations

A matrix equation  $\begin{bmatrix} 5 & 2 \\ 3 & 10 \end{bmatrix} \times \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9 \\ 23 \end{bmatrix}$ , can be written as  $\begin{bmatrix} 5x + 2y \\ 3x + 10y \end{bmatrix} = \begin{bmatrix} 9 \\ 23 \end{bmatrix}$ .

Now comparing the corresponding elements of equal matrices

$$5x + 2y = 9, 3x + 10y = 23$$

Which is a system of two linear equations in two variables x and y.

### Conversion of System of Linear Equations into Matrix Equations

By the converse process mentioned above, the system of linear equations can be rewritten in matrix equation:

$$\text{i.e. } 5x + 2y = 9, 3x + 10y = 23$$

writing in matrix form, we have

$$\begin{bmatrix} 5x & +2y \\ 3x & +10y \end{bmatrix} = \begin{bmatrix} 9 \\ 23 \end{bmatrix}$$

$$\text{or } \begin{bmatrix} 5 & 2 \\ 3 & 10 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 9 \\ 23 \end{bmatrix} \text{ or } AX = B \text{ is the required matrix equation.}$$

### Check Point:

Is  $AB = I = BA$  always true?

## Solution of System of two Linear Equations in Two Variables

We discuss here two methods for finding solution of liner equations in two variables.

- (a) Matrix inversion method
- (b) Cramer's rule

### (a) Matrix Inversion Method

Let us find the solution of:

$$5x + 4y = 14 ; \quad 3x + 7y = 13$$

This system of equations can be rewritten in matrix equation as follows:

$$\begin{bmatrix} 5 & 4 \\ 3 & 7 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 14 \\ 13 \end{bmatrix} \quad \dots \quad (\text{i})$$

where  $A = \begin{bmatrix} 5 & 4 \\ 3 & 7 \end{bmatrix}$  is a matrix of coefficients,  $X = \begin{bmatrix} x \\ y \end{bmatrix}$  is a matrix of variables and  $B = \begin{bmatrix} 14 \\ 13 \end{bmatrix}$  is a matrix of constants.

Now equation (i) can also be written as:

$$AX = B$$

$$A^{-1} A X = A^{-1} B \quad (\text{Pre multiplying with } A^{-1} \text{ if it exists.})$$

$$I X = A^{-1} B \quad (A^{-1} A = I)$$

$$X = A^{-1} B \quad \dots \quad (\text{ii})$$

This equation indicates that the values of the variables in the matrix  $X$  are equal to the corresponding elements in the matrix  $A^{-1} B$ . To find  $A^{-1} B$ , we need  $A^{-1}$ .

As,  $|A| = \begin{vmatrix} 5 & 4 \\ 3 & 7 \end{vmatrix} = 35 - 12 = 23 \neq 0$ , so inverse is possible.

$$\text{Also, } \text{adj } A = \begin{bmatrix} 7 & -4 \\ -3 & 5 \end{bmatrix}$$

$$\text{Now, } A^{-1} = \frac{1}{|A|} \times \text{adj } A = \frac{1}{23} \times \begin{bmatrix} 7 & -4 \\ -3 & 5 \end{bmatrix}$$

Substituting the values in equation (ii), we get:

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{23} \times \begin{bmatrix} 7 & -4 \\ -3 & 5 \end{bmatrix} \begin{bmatrix} 14 \\ 13 \end{bmatrix} = \frac{1}{23} \times \begin{bmatrix} 98 - 52 \\ -42 + 65 \end{bmatrix} = \begin{bmatrix} \frac{1}{23} \times 46 \\ \frac{1}{23} \times 23 \end{bmatrix}$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Comparing the elements of both matrices, we see that  $x = 2$  and  $y = 1$  solution set =  $\{(2, 1)\}$

#### Example:

Solve system of linear equation  $3y = 24 - 9x$ ,  $2y + 6x = 10$  (if possible) by matrix inversion method.

**Solution:** Writing the given system of linear equations in arranged form. i.e.

$$9x + 3y = 24, \quad 6x + 2y = 10.$$

Which can be written in matrix equation as:

$$\begin{bmatrix} 9 & 3 \\ 6 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 24 \\ 10 \end{bmatrix}$$

$$\text{or } AX = B \Rightarrow X = A^{-1} B$$

$$\text{Now } |A| = \begin{vmatrix} 9 & 3 \\ 6 & 2 \end{vmatrix} = 9 \times 2 - 6 \times 3 = 18 - 18 = 0$$

As the matrix A is a singular, so its multiplicative inverse does not exist and solution of given system of equation is not possible.

### (b) Cramer's Rule

This method is named after Gabriel Cramer (1704 -1752). This rule uses determinants to find the solution of system of linear equations.

Consider a system of linear equations as  $x + 2y = 6$ ,  $4x - 2y = 4$ .

Writing it in matrix equation, we get:

$$\begin{bmatrix} 1 & 2 \\ 4 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix} \quad \text{or} \quad A X = B$$

The special formulae used in this method are:  $x = \frac{D_x}{D}$  and  $y = \frac{D_y}{D}$

Where,  $D = |A| = \begin{vmatrix} 1 & 2 \\ 4 & -2 \end{vmatrix} = -2 - 8 = -10 \neq 0$ . (non singular)

Now

$$D_x = \begin{vmatrix} 6 & 2 \\ 4 & -2 \end{vmatrix} = -12 - 8 = -20$$

Original  $C_2$  of A  
Contributed by B

$$D_y = \begin{vmatrix} 1 & 6 \\ 4 & 4 \end{vmatrix} = 4 - 24 = -20$$

Original  $C_1$  of A  
Contributed by B

Using  $x = \frac{D_x}{D}$  and  $y = \frac{D_y}{D}$   
 $x = \frac{-20}{-10}$                                    $y = \frac{-20}{-10}$   
 $x = 2$      $y = 2$  and solution set =  $\{(2, 2)\}$

### Exercise 3.3

1. (a) Mention singular and nonsingular matrices from the followings.

$$A = \begin{bmatrix} 6 & 4 \\ 2 & 3 \end{bmatrix} \quad B = \begin{bmatrix} -5 & 7 \\ -9 & 4 \end{bmatrix}, \quad C = \begin{bmatrix} 9 & 18 \\ 2 & 4 \end{bmatrix}$$

$$D = \begin{bmatrix} a & b + 1/a \\ a & b \end{bmatrix} \quad E = \begin{bmatrix} a & b + c \\ a & d + c \end{bmatrix} \quad (\text{where } a \neq 0 \text{ but } b = d)$$

(b) If  $|P| = 9$  and  $P = \begin{bmatrix} 3 & 3 \\ 1 & 4k \end{bmatrix}$ , then find k.

(c) If  $|T| = 3$  and  $\text{adj } T = \begin{bmatrix} 5 & x \\ 3 & 2 \end{bmatrix}$ , then find x.

2. (a) Find the multiplicative inverses of these matrices if possible.

$$R = \begin{bmatrix} 4 & -1 \\ -6 & 2 \end{bmatrix}, \quad S = \begin{bmatrix} 5 \\ 3 \end{bmatrix}, \quad T = \begin{bmatrix} 25 & 2 \\ 50 & 4 \end{bmatrix}, \quad U = \begin{bmatrix} x & x+1 \\ y & y+1 \end{bmatrix}, \text{ if } x = y.$$

(b) If  $R = \begin{bmatrix} 4 & 1 \\ -6 & 2 \end{bmatrix}$  then verify that  $R R^{-1} = R^{-1} R = I$ .

3. If  $Y = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}$  and  $Z = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ , then verify that  $(YZ)^{-1} = Z^{-1}Y^{-1}$ .

4. What is relation among  $|A|$ ,  $|A^{-1}|$  and  $|\text{adj } A|$  if  $A = \begin{bmatrix} 5 & 7 \\ 2 & 3 \end{bmatrix}$ ?

5. Write these matrix equations into system of linear equations if possible.

(i)  $\begin{bmatrix} 4 & 2 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 6 \\ 4 \end{bmatrix}$       (ii)  $\begin{bmatrix} 5 & 0 \\ 0 & 4 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 10 \\ 20 \end{bmatrix}$

(iii)  $\begin{bmatrix} 0 \\ 3 \end{bmatrix} \times \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 12 \\ 24 \end{bmatrix}$       (iv)  $\begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} 5 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \end{bmatrix}$

6. (a) Write the systems of linear equations in matrix form.

(i)  $x + y = 2$       (ii)  $2x + y = 90$       (iii)  $y = 3$   
 $x - y = 4$        $5y - x = 10$        $x = 4$

(iv)  $\frac{5}{2}x - 3y = 1$ ,  $\frac{1}{2}y - 4x = 2$

(b) If matrix of coefficients of  $5x - 4y = 30$ ,  $10x - ky = 60$  is singular then find k.

7. (a) Use Matrix Inversion Method to solve the following systems of linear equations if possible.

(i)  $4x + 3y = -6$       (ii)  $-x + 2y = 1.5$   
 $x + 2y = 1$        $5x + 4y = 3$

(iii)  $2y = 10 - 16x$       (iv)  $9 - x = 7y$   
 $24x = 15 - 3y$        $14y + 2x = 18$

(b) Use Cramer's rule to solve following systems of linear equations if possible.

(i)  $2x + 3y = 5$       (ii)  $x = \frac{2}{3} - 2y$   
 $5x + 10y = 10$        $4y = 3 - 3x$   
 (iii)  $\frac{6}{10}x + \frac{8}{10}y = 20$       (iv)  $16x - 10 + 2y = 0$   
 $\frac{8}{10}x - \frac{6}{10}y = 10$        $15 - 3y - 24x = 0$

## Applications of Matrices

The most difficult part of solving a problem in algebra is almost always translating the problem situation to mathematical language. Once an equation is translated, the rest is usually straightforward. In this section, we study systems of equations and how to solve them using matrices. We can use matrices to help us solve problems that involve systems of equations. Using matrices often simplifies the process of solving these systems.

### **Example:**

Anas bought 3 cream puffs and 5 pringle packs paying Rs.650. Abid bought 4 cream puffs and 2 pringle packs paying Rs. 400. Find how much each item costs? (Hint: Either matrix inversion method or Cramer's rule can be used).

### **Solution:**

Let  $x$  be the cost of a cream puff and  $y$  be the cost of a pringle pack, then Anas's shopping is represented by  $3x + 5y = 650$  and Abid's shopping is represented by equation  $4x + 2y = 400$

Writing in matrix equation

$$\begin{bmatrix} 3 & 5 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 650 \\ 400 \end{bmatrix} \quad \text{or} \quad AX = B$$

Now by using Cramer's rule

$$D = |A| = \begin{vmatrix} 3 & 5 \\ 4 & 2 \end{vmatrix} = 6 - 20 = -14$$

$$D_x = \begin{vmatrix} 650 & 5 \\ 400 & 2 \end{vmatrix} = 1300 - 2000 = -700, D_y = \begin{vmatrix} 3 & 650 \\ 4 & 400 \end{vmatrix} = 1200 - 2600 = -1400$$

$$\text{Now, } x = \frac{D_x}{D} \quad \text{and} \quad y = \frac{D_y}{D}$$

$$x = \frac{-700}{-14} \quad \text{and} \quad y = \frac{-1400}{-14} \Rightarrow x = 50 \quad \text{and} \quad y = 100$$

So, each cream puff costs Rs. 50 and each pringle pack costs Rs. 100.

### **Example:**

Naveed is a chemist who is preparing an acid solution to be used as a cleaner for machine parts. The machine shops need several batches of 200 ml of solution at a 48% concentration. He only has 60% and 40% concentration solutions. The two solutions can be combined to make the 48% solution. How much of each solution should Naveed use to make 200 ml of solution?

### **Solution:**

Let  $x$  represent the amount of 60% solution and let  $y$  represent the amount of 40% solution.

$$x + y = 200 \quad \text{The total of two amount must be 200 ml.}$$

Now write an equation that represents the proportions of each solution needed.

$$60\% + 40\% = 48\%$$

$$60\% (x) + 40\% (y) = 48\% (x + y) \quad \text{Each part contribute to the total}$$

$$0.60(x) + 0.40(y) = 0.48(x + y) \quad \text{Multiply 100 to remove the decimals}$$

$$12x - 8y = 0 \quad \text{Write the equation in standard form}$$

**Step 1:** Write a system of equations. Then write the system as a matrix equation.

$x + y = 200$ ,  $12x - 8y = 0$  in matrix form:  $AX=B$ , and  $X=A^{-1}B$

$$\begin{bmatrix} 1 & 1 \\ 12 & -8 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 200 \\ 0 \end{bmatrix}, \text{ where } A = \begin{bmatrix} 1 & 1 \\ 12 & -8 \end{bmatrix}, X = \begin{bmatrix} x \\ y \end{bmatrix} \text{ and } B = \begin{bmatrix} 200 \\ 0 \end{bmatrix}$$

**Step 2:** To solve the matrix equation, first find the inverse of the matrix A.

$$A^{-1} = \frac{1}{-20} \begin{bmatrix} -8 & -1 \\ -12 & 1 \end{bmatrix}, \text{ where } |A| = -8 - 12 = -20 \text{ and Adjoint of } A = \begin{bmatrix} -8 & -1 \\ -12 & 1 \end{bmatrix}$$

$$X = \frac{1}{-20} \begin{bmatrix} -8 & -1 \\ -12 & 1 \end{bmatrix} \begin{bmatrix} 200 \\ 0 \end{bmatrix}, \text{ where } X = A^{-1}B$$

$$\begin{bmatrix} x \\ y \end{bmatrix} = \frac{1}{-20} \begin{bmatrix} -1600 \\ -2400 \end{bmatrix} = \begin{bmatrix} 80 \\ 120 \end{bmatrix}$$

We have  $x = 80$  and  $y = 120$ .

This means that 80 ml of the 60% solution is added to 120 ml of the 40% solution to make 200 ml of the 48% solution.

### Exercise 3.4

- From the two acute angles of a right triangle, one angle is 10 (degrees) less than the other. Find both the acute angles by using Cramer's rule.
- In a quiz, Abdullah correctly replied 15 out of 20 questions from literature and science, getting Rs.125. If every correct literature reply is credited Rs. 5 and every correct science reply is credited Rs. 10, then find the number of correct replies of literature and science separately.
- If Asad bought 3 notebooks and 2 books paying Rs. 600 and Iman bought 4 notebooks and 1 book paying Rs. 550, find the cost of each notebook and book? (Use Cramer's rule).
- Sum of ages of two sisters Sundas and Zenab is 22 years. If Sundas is 1 year older than twice the age of Zenab. Find the ages of both the sisters by using matrix inversion method.
- Two trucks have capacities of 10 tons and 12 tons. They made a total of 20 round trips to haul 226 tons of sand to the community park. How many round trips did each truck make?
- To make 20 kg of aluminum alloy with 70% aluminum, a metallurgist wants to use two metals with 55% and 80% aluminum content. How much of each metal should he use.
- Soybean meal contains 16% protein and corn meal contains 9% protein. How many pounds of each should be mixed together in order to get a 350-lb mixture that is 12% protein?
- The perimeter of a lot is 190 m. The width is one fourth of the length. Find the dimensions.

- Displaying information in the form of matrices of order 2.
- Calculating product of scalar quantity and a matrix.
- Solving situations involving sum, difference and product of two matrices.
- Evaluating the determinant and inverse of a matrix of order 2 by 2.
- Solving the simultaneous linear equations in two variables using matrix inversion method and Cramer's rule.
- Explaining with examples, how mathematics plays a key role in the development of new scientific theories and technologies.
- Applying concepts of matrices to real world problems

**MISCELLANEOUS EXERCISE-3**

1. Encircle the correct option in the following.

i. If  $\begin{bmatrix} 5 & 3 \\ 2 & 9 \end{bmatrix}^t = \begin{bmatrix} 5 & x \\ 3 & 2 \end{bmatrix}$ , then  $x =$

- (a) 6      (b) 4      (c) -6      (d) -4

ii. If  $I_3 = \begin{bmatrix} y & 0 & x \\ 0 & z & 0 \\ x & 0 & 1 \end{bmatrix}$  then

- (a)  $y = x = 1$     (b)  $x = z = 0$     (c)  $x = z = 1$     (d)  $y = z = 1, x = 0$

iii. Additive inverse of unit matrix of order 2, is

- (a)  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$     (b)  $\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$     (c)  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$     (d)  $\begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$

iv. Multiplicative inverse of a null matrix of order 2, is

- (a)  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$     (b)  $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$     (c)  $[0 \ 0]$     (d) impossible

v. A is a symmetric matrix if

- (a)  $A^t \neq A$     (b)  $(A^t)^t \neq -A$     (c)  $(A^t)^t = -A$     (d)  $(A^t)^t = A^t$

vi.  $\begin{bmatrix} 1 \\ 2 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \end{bmatrix} =$

- (a)  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$     (b)  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$     (c)  $\begin{bmatrix} \frac{1}{2} & 0 \\ 0 & 0 \end{bmatrix}$     (d) impossible

vii. Order of matrix A is 1-by-2 and order of matrix B is 2-by-3 then order of AB is

- (a) 1-by-3    (b) 3-by-1    (c) 2-by-2    (d) 3-by-2

viii. If  $AB = B$ , then  $A = \dots$

- (a) I    (b)  $A^{-1}$     (c) B    (d)  $B^{-1}$

ix.  $\begin{bmatrix} 15 \\ 25 \end{bmatrix} \times [3 \ 2] =$

- (a)  $[95]$     (b)  $\begin{bmatrix} 45 \\ 50 \end{bmatrix}$     (c)  $[45 \ 50]$     (d)  $\begin{bmatrix} 45 & 30 \\ 75 & 50 \end{bmatrix}$

x. If  $|T| = -1$  then  $T^{-1} =$

- (a)  $-T$  (b)  $\text{adj } T$  (c)  $-\text{adj } T$  (d)  $T$

xii. Matrix equation for  $y + x = 144$  and  $x + 2y = x + 13$  is

(a)  $\begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \times \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 144 \\ 13 \end{bmatrix}$  (b)  $\begin{bmatrix} 1 & 0 \\ 1 & 2 \end{bmatrix} \times \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 144 \\ 13 \end{bmatrix}$

(c)  $\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \times \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 144 \\ 13 \end{bmatrix}$  (d)  $\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \times \begin{bmatrix} y \\ x \end{bmatrix} = \begin{bmatrix} 144 \\ 13 \end{bmatrix}$

xiii. The matrix of coefficients for  $x - y = 3$ , is

- (a) [3] (b)  $\begin{bmatrix} 1 & -1 \end{bmatrix}$  (c)  $\begin{bmatrix} x \\ y \end{bmatrix}$  (d)  $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$

xiv.  $[1 \ 2][3] =$

- (a) [3 6] (b)  $\begin{bmatrix} 3 \\ 6 \end{bmatrix}$  (c) [5] (d) impossible

xv. If  $\begin{bmatrix} -3 & 5 \\ -3 & x-1 \end{bmatrix}$  is singular matrix, then  $x =$

- (a) 4 (b) 6 (c) -6 (d) -4

xvi. If  $\text{adj } A = \begin{bmatrix} 5 & 6 \\ 2 & 3 \end{bmatrix}$  then  $|A| =$

- (a)  $|\text{adj } A|$  (b)  $|A^t|$  (c)  $|-A|$  (d) all a, b, c

2. If  $P = \begin{bmatrix} 5 & -1 \\ 2 & -4 \end{bmatrix}$ , then show that  $PP^{-1} = P^{-1}P = I$ .

3. Solve:  $5x - 4 = y$ ;  $2y + 8 = 10x$ , by Matrix Inversion Method if possible.

4. Use Cramer's rule for solution of  $5x + 2y = 19$  and  $10x + 4y = 38$ , if possible.

5. The daily sum of labour and the material costs of Haani's and Massab's Home Industries are Rs.20,000 and Rs.40,000 respectively. If labour and the material rate, for both the Home Industries is same but Haani's Home Industry has 6 workers who make 80 items and Massab's Home Industry has 12 workers who make 160 items, find the labour and the material cost.

# 04

## Linear and Quadratic Inequalities

After studying this unit students will be able to:

- Solve linear inequalities in one variable.
- Solve quadratic inequalities in one variable.
- Solve two linear inequalities with two unknowns simultaneously.
- Interpret and identify the region in plane bounded by two linear inequalities in two unknowns.

You are checking a bag at an airport. Bags can weigh no more than 60 pounds. Your bag weighs 26.5 pounds. Find the possible weights 'w' (in pounds) that you can add to the bag.

Write a model:

Weight of bag + Weight you can add  $\leq$  Weight limit

$$26.5 + w \leq 60 \quad (\text{writing inequalities})$$

$$26.5 + w - 26.5 \leq 60 - 26.5 \quad (\text{subtracting 26.5 from both sides})$$

$$w \leq 33.5 \quad (\text{simplifying})$$

You can add less than 33.5 pounds.



## Inequalities in One Variable

Furqan and Qasim are the hockey players in the school hockey team. If we compare their scoring for the season, only one of the following statements will be true:

Furqan scored fewer goals than Qasim.

Furqan scored same number of goals as Qasim.

Furqan scored more goals than Qasim.

Let ' $a$ ' represents the number of goals of Furqan and ' $b$ ' represents the number of goals of Qasim.

We can compare the score using an inequality or an equation.


$$a < b, \quad a > b, \quad a = b$$

This is an example of trichotomy property.

### Graph of Inequality

On a number line, the graph of an inequality in one variable is the set of points that represents all solutions of the inequality. To graph an inequality in one variable, we use open circle for the symbols ' $<$ ' or ' $>$ ' and a closed circle for the symbols ' $\leq$ ' or ' $\geq$ '.

The graph of  $x < 1$ ,  $x = 1$  and  $x \geq 1$  respectively are:



**Example:** Solve  $9x + 6 < 8x - 2$ . Graph the solution set.


**Solution:** 
$$9x + 6 < 8x - 2$$

$$9x + 6 + (-8x) < 8x + (-8x) - 2 \quad (\text{Add } -8x \text{ to both sides})$$

$$x + 6 < -2$$

$$x + 6 + (-6) < -2 + (-6) \quad (\text{Add } -6 \text{ to both sides})$$

$$x < -8$$



Circle indicates that -8 is not included.

Any real number less than -8 is solution.



### Trichotomy Property

For any two real numbers  $a$  and  $b$ , exactly one of the following statements will be true:

$$a < b, \quad a > b, \quad a = b$$

**Check:** Substitute  $-8$  for  $x$  in  $9x+16 < 8x-2$ , the inequalities should not be true. Then substitute a number less than  $-8$ , the inequality should be true.

**Key Fact:**

(i) Addition and subtraction properties of inequalities.

For any real number  $a$ ,  $b$  and  $c$ :

- If  $a > b$ , then  $a+c > b+c$  and  $a-c > b-c$
- If  $a < b$ , then  $a+c < b+c$  and  $a-c < b-c$

(ii) Multiplication and division properties of inequalities.

- If  $c$  is positive and  $a < b$ , then  $ac < bc$  and  $\frac{a}{c} < \frac{b}{c}$

- If  $c$  is positive and  $a > b$ , then  $ac > bc$  and  $\frac{a}{c} > \frac{b}{c}$

- If  $c$  is negative and  $a < b$ , then  $ac > bc$  and  $\frac{a}{c} > \frac{b}{c}$

- If  $c$  is negative and  $a > b$ , then  $ac < bc$  and  $\frac{a}{c} < \frac{b}{c}$

**Example:** Solve  $6x - 7 \geq 2x + 17$ . Graph the solution.

**Solution:**  $6x - 7 \geq 2x + 17$ .


$$6x - 7 + 7 \geq 2x + 17 + 7 \quad (\text{adding } 7 \text{ on both sides})$$

$$6x \geq 2x + 24$$

$$6x - 2x \geq 2x - 2x + 24 \quad (\text{subtracting } 2x \text{ from both sides})$$

$$4x \geq 24 \quad (\text{dividing both sides by } 4)$$

$$x \geq 6$$



**Example:** Solve and graph the solution

$$10.7x - 6 + 5.5x \geq 3(5x - 3)$$

**Solution:**  $10.7x - 6 + 5.5x \geq 3(5x - 3)$

$$16.2x - 6 \geq 15x - 9$$

$$16.2x - 15x \geq -9 + 6$$

$$1.2x \geq -3$$

$$x \geq -2.5$$

**Check Point:**

Describe and correct the error.


a.  $17 - 3x \geq 56$

$$-3x \geq 39 \Rightarrow x \geq -13$$

b.  $-4(2x - 3) < 28$

$$-8x + 12 < 28$$

$$-8x < 16 \Rightarrow x > -2$$



## Solving Quadratic Inequalities

We will discuss the solution of quadratic inequalities in one variable. Quadratic inequalities can be derived from quadratic equations. The word ‘Quadratic’ comes from the word ‘Quadratum’ which means square in Latin. From this, we can define quadratic inequalities on second degree equation. The general form of a quadratic equation is  $ax^2 + bx + c = 0$ . Further if the quadratic polynomial  $ax^2 + bx + c$  is not equal to zero, then they are either  $ax^2 + bx + c > 0$  or  $ax^2 + bx + c < 0$  and are called quadratic inequalities. We explain quadratic inequalities with daily life example.

Consider a ball is thrown upward with an initial velocity of 64 feet per second.

The function  $h(t) = 64t - 16t^2$  gives the height of the ball after  $t$  seconds. During what time span is the ball in flight? We can find the times that the ball is in flight by solving the inequality

$$64t - 16t^2 > 0.$$

$$64t - 16t^2 > 0$$

$$-16t^2 + 64t > 0$$

$$-16t(t - 4) > 0$$



This inequality shows that the product of two numbers is positive if both of the numbers  $(-16t, t - 4)$  are positive or both are negative.

### Case I:

When both numbers are positive

$$-16t > 0 \text{ and } t - 4 > 0$$

$$t < 0 \text{ and } t > 4$$

Here,  $t < 0$  and at the same time  $t > 4$  is never true.

### Case II:

When both numbers are negative

$$-16t < 0 \text{ and } t - 4 < 0$$

$$t > 0 \text{ and } t < 4$$

Means:  $0 < t$  and  $t < 4$

$$0 < t < 4$$

The solution set is  $\{0 < t < 4\}$ . So, the ball is in flight between 0 and 4 seconds after it is thrown.

**Example:** Solve  $x^2 - 4x - 12 > 0$

**Solution:**  $x^2 - 4x - 12 > 0$

$$\begin{aligned} x^2 - 6x + 2x - 12 &> 0 \Rightarrow x(x - 6) + 2(x - 6) > 0 \\ \Rightarrow (x - 6)(x + 2) &> 0 \end{aligned}$$


Another way that you can solve quadratic inequalities is by using three test points.

First solve the equation  $(x - 6)(x + 2) = 0$

$$x - 6 = 0 \text{ or } x + 2 = 0$$

$$\Rightarrow x = 6, x = -2$$

These solutions are called critical points. The points  $-2$  and  $6$  separate the  $x$ -axis into three regions  $x < -2$ ,  $-2 < x < 6$ ,  $x > 6$ .




Choose a value from each of the parts and substitute it into  $(x - 6)(x + 2) > 0$ .

Construct a table to organize your results.

Part of x-axis	Point x	$(x - 6)(x + 2)$	Is $(x - 6)(x + 2) > 0$ ?
$x < -2$	-3	$(-3 - 6)(-3 + 2) = 9$	Yes
$2 < x < 6$	0	$(0 - 6)(0 + 2) = -12$	No
$x > 6$	8	$(8 - 6)(8 + 2) = 20$	Yes

The part of x-axis where points are solution of  $(x - 6)(x + 2) > 0$ , belongs to the solution set.

The solution set is  $\{x | x < -2 \text{ or } x > 6\}$ .



**Example:** Find the solution set of  $x^2 + 6x - 27 \leq 0$ .

$$\begin{aligned} \text{Solution: } x^2 + 6x - 27 \leq 0 &\Rightarrow x^2 + 9x - 3x - 27 \leq 0 \\ x(x+9) - 3(x+9) \leq 0 &\Rightarrow (x+9)(x-3) \leq 0 \end{aligned}$$

First solve the equation  $(x-3)(x+9)=0$


$$x-3=0 \text{ or } x+9=0 \Rightarrow x=3, x=-9$$

The point 3 and -9 are separate, its x-axis in three parts.

Make a table to organize results.

Part of x-axis	Point x	$(x-3)(x+9)$	Is $(x-3)(x+9) \leq 0$ ?
$x \geq 3$	4	$(4-3)(4+9)=13$	No
$-9 \leq x \leq 3$	0	$(0-3)(0+9)=-27$	Yes
$x \leq -9$	-10	$(-10-3)(-10+9)=12$	No

$$-9 \leq x \leq 3$$



The part of x-axis where points are solution to  $(x-3)(x+9) \leq 0$  belong to the solution set.

The solution set is  $\{x | -9 \leq x \leq 3\}$ .

### Exercise 4.1

Solve the inequality and graph the solution.

1.  $5x - 12 \leq 3x - 4$
2.  $1 - 8x \leq -4(2x - 1)$
3.  $\frac{-2}{3}x - 2 < \frac{1}{3}x + 8$
4.  $8 - \frac{4}{5}x > -14 + 2x$
5.  $-0.6(x - 5) \leq 15$
6.  $\frac{-1}{4}(x - 12) > -2$


Translate the phrase into an inequality. Then solve the inequality and graph the solution.

7. Four more than the product of 3 and  $x$  is less than 40.


8. Twice the sum of  $x$  and 8 appears less than or equal to -2.

Write and solve an inequality to find the possible values of

9. Area  $>$  81 square feet



10. Area  $\leq$  44 square centimeters



Solve each inequality

11.  $(x+1)(x-3) > 0$
12.  $(x-9)(x+1) < 0$
13.  $x^2 - x - 90 > 0$
14.  $x^2 + 4x - 21 < 0$
15.  $x^2 + 8x + 16 \geq 0$
16.  $9x^2 - 6x + 1 \leq 0$
17. Mr. Khalid has a field and want to make a rectangular garden with perimeter of 68ft. He would like the area of a garden to be at least 240 square feet. What would the width of the garden be?
18. Are -2, -1, 0, 1, 2 solution of the inequality  $2x^2 + 3x + 1 \leq 0$ .
19. The stopping distance  $d(x)$  (in meters) of a car traveling at  $x$  km/h is modeled by  $d(x) = 0.05x^2 + 0.2x$ . If the maximum stopping distance allowed in a school zone is 35 meters, find the speed range that ensures the stopping distance does not exceed this limit.
20. A rectangular garden is to be enclosed with a fence. The total length of the fence is at most 60 meters and the width  $x$  of the garden is twice its length. Find the possible dimensions of the garden.


## Solving System of Linear Inequalities

Mr. Waleed wants to rent a car for a business trip. A car rental company advertises that their rental rate is Rs.9000 plus Rs.60 per km. Mr. Waleed would like to compare this rate with the rates offered by other car rental agencies. First, he determines the equation containing the points that represent the relationship between the numbers of km ( $x$ ) and the total cost of the rental ( $y$ ). The initial cost of the car is Rs. 9000. Since this is the point where zero km are driven, it would be the  $y$ -intercept of the graph.

The slope would be the rate of change in the total cost. In this case, the rate is Rs.60 per km.

Since the slope is 60. Thus, an equation of the line is  $y = 60x + 9000$ .


km ( $x$ )	Cost ( $y$ )
0	9000
20	10200
40	11400
100	15000



The graph of  $y = 60x + 9000$  separates the coordinate plane into two regions.

The line is called boundary of the region. To graph an inequality, first you graph the boundary and then determine which region to shade. The graph  $y > 60x + 9000$  contains the points that are located above the boundary.

In that region, the value of the dependent variable  $y$  is greater than the value of  $60x + 9000$ . This graph represents car rental costs that are greater than those offered by car rental company A. For example, another company B charges 14000 for a car rental with 50km.




The point (50, 14000) lies above the boundary.

The graph of  $y > 60x + 9000$  contains points that are located below the boundary. In that region, the value of  $y$  is less than the value of  $60x + 9000$ .

The graph represents car rental costs that are less than those offered by company A.

For example, a company C charges 14000 For a car rental with 100km. the point (100, 14000) lies below the boundary.



- When graphing an inequality, the boundary you draw may be solid or dashed.
- If the inequality uses the symbol  $\leq$  or  $\geq$  which include equality then the boundary will be solid. Otherwise, it will be dashed.
- After graphing the boundary, you must determine which region is to be shaded. Test a point on one side of the line. If the ordered pair satisfies the inequality, that region contains solution of the inequality. If the ordered pair does not satisfy the inequality, the other region is the solution.

#### Key Fact:

A linear inequality in two variables: such as  $x - 3y < 6$ , is the result of replacing the “=” in a linear equation with  $<$ ,  $\leq$ ,  $>$  or  $\geq$ . A solution of an inequality in two variables  $x$  and  $y$  is an ordered pair  $(x, y)$  that produces a true statement when the values of  $x$  and  $y$  are satisfied into the inequality.

**Example:** Which ordered pair is not a solution of  $x - 3y \leq 6$ ?

- (0, 0)
- (6, -1)
- (10, 3)
- (-1, 2)

#### Solution:

Test (0, 0):  $x - 3y \leq 6$  (Write inequality)

$$0 - 3(0) \leq 6 \quad (\text{Substitute } 0 \text{ for } x \text{ and } 0 \text{ for } y.)$$

$0 \leq 6$  (True)

Test (6, -1):  $x - 3y \leq 6$  (Write inequality)

$$6 - 3(-1) \leq 6 \quad (\text{Substitute } 6 \text{ for } x \text{ and } -1 \text{ for } y.)$$


$9 \leq 6$  (False)

Similarly (10, 3) and (-1, 2) also true but (6, -1) is not a solution.

#### Feasible region

Inequalities represent regions graphically. Below are the rules in drawing feasible region:

$x \text{ or } y <$	$\geq \text{ or } \leq$
Dashed line	Solid line



**Example:** Graph the inequality  $y > 5x - 4$ .

**Solution:**

**Step I:** Graph the equation  $y = 5x - 4$ .


The inequality is  $>$ , use a dashed line.

**Step II:** Test  $(0, 0)$  in  $y > 5x - 4$ .

$$0 > 4(0) - 4$$

$$0 > -4 \text{ (True)}$$

**Step III:** Shade the half plane that contains  $(0, 0)$  because  $(0, 0)$  is a solution of the inequality.



**Example:** Graph the inequality  $x + 2y \leq 0$ .

**Solution:**

**Step I:** Graph the equation  $x + 2y = 0$ .


The inequality is  $\leq$ , use a solid line.

**Step II:** Test  $(1, 0)$  in  $x + 2y \leq 0$ .

$$1 + 2(0) \leq 0$$

$$1 \leq 0 \text{ (False)}$$

**Step III:** Shade the half plane that does not contain  $(1, 0)$  because  $(1, 0)$  is not a solution of the inequality.



**Example:** Graph the inequality  $y \geq -2$ .

**Solution:**


**Step I:** Graph the equation  $y = -2$ .

The inequality is  $\geq$ , use a solid line.

**Step II:** Test  $(0, 0)$  in  $y \geq -2$ .

$$0 \geq -2 \text{ (True)}$$

**Step III:** Shade the half plane that contains  $(0, 0)$  because  $(0, 0)$  is a solution of the inequality.



**Example:** Graph the inequality  $x < -1$ .

**Solution:**


**Step I:** Graph the equation  $x = -1$ .

The inequality is  $\geq$ , use a solid line.

**Step II:** Test  $(0, 0)$  in  $x < -1$

$$0 < -1 \text{ (False)}$$

**Step III:** Shade the half plane that does not contain  $(0, 0)$  because  $(0, 0)$  is not a solution of the inequality.



### Exercise 4.2

Tell whether the ordered pair is a solution of the inequality.

1.  $(0, 0); x + y < -4$

2.  $(-2, -3); y - x > -2$

3.  $(5, 2); 2x + 3y \geq 14$

4.  $(-1, 5); 4x - 7y > 28$

5.  $(-9, -7); y \leq 8$

6.  $(-4, 0); x \geq -3$

Graph the inequality.

7.  $x - y < -3$

8.  $-3y - 2x < 12$

9.  $x - y \geq 2$

10.  $2x + y \geq 8$

11.  $x - y \leq -11$

12.  $y < -5$

13.  $x \geq 4$

14.  $\frac{1}{2}(x + 2) + 3y < 8$

15. Can we use  $(0, 0)$  as a test point when graphing  $x + y > 0$ ? Explain with reason.

Write the verbal sentence as an inequality. Then graph the inequality:

16. Three less than  $x$  is greater than or equal to  $y$ .

17. The product of  $-2$  and  $y$  is less than or equal to the sum of  $x$  and  $6$ .

18. The sum of  $x$  and the product of  $4$  and  $y$  is less than  $-2$ .


# Absolute Valued Equations and Inequalities

## Equations with Absolute Value

Suppose you and your friend, each live in the same street as your school, but on opposite sides of the school. You each live 6 km from the school. What can you say about your trips to school each day? Do you each travel the same distance?

Consider placing both houses and the school on a number line with the school at the origin. Your home is located at 6, and your friend is located at  $-6$ .


Certainly,  $-6$  and 6 are quite different but they do have something in common. They have the same distance from 0 on the number line. This means that you and your friend travel the same distance, but in different direction, when you go to school.



We say that  $-6$  and 6 have the same absolute value. The absolute value of a number is the number of units it is from 0 in the number line.

We use the symbol  $|x|$  to represent the absolute value of a number  $x$ .

The absolute value of 6 is  $|6|=6$ , and absolute value of  $-6$  is  $|-6|=6$ .



In words: The definition states that the absolute value of a non-negative is the number itself and the absolute value of a negative number is the opposite of the number.

Since distance is always non-negative, we can think of a number's absolute value as its distance from zero on number line.

**Example:** Find the absolute value of 7 and  $-11$ .

**Solution:**  $|7|=7$        $|-11|=-(-11)=11$

**Example:** Find the absolute value of  $x - 5$ .

**Solution:** Let's make a list of possible cases.

If  $x$  is 5 or greater, then  $|x - 5| \geq 0$ ,

### Key Fact:

The absolute value of a number is always positive or zero.

$$|x - 5| = x - 5$$

If  $x$  is less than 5, then  $|x - 5| < 0$ ,

$$|x - 5| = -(x - 5) = 5 - x$$

**Example:** Find the solution set.

a.  $|2x + 5| = 11$

c.  $|2x - 3| = |x + 6|$

b.  $|3 - 2x| = -5$

d.  $2|x + 3| + 2 = 16$

**Solution:**

a.  $|2x + 5| = 11$

$2x + 5 = 11$

$2x = 11 - 5$

$2x = 6$

$x = 3$

S.S. = {3, -8}

c.  $|2x - 3| = |x + 6|$

$2x - 3 = x + 6$

$2x - x = 6 + 3$

$x = 9$

S.S. = {-1, 9}

b.  $|3 - 2x| = -5$

The absolute value principle reminds us that absolute value is always non-negative. The equation  $|3 - 2x| = -5$  has no solution. The solution set is  $\emptyset$ .

d.  $2|x + 3| + 2 = 16$

$2|x + 3| + 2 = 16$

$2|x + 3| = 14 - 2$

$|x + 3| = 7$

$x + 3 = 7 \quad \text{or} \quad x + 3 = -7$

$x = 4 \quad \text{or} \quad x = -10$

S.S. = {4, -10}

Student can verify the solution.

**Example:** A machine that fills the boxes of sugar is to fill the boxes with 14 kg of sugar. After the boxes are filled, another machine weighs the boxes. If the box is more than 0.3 kg above or below the desired weight, the box is rejected. What is the weight of the heaviest and the lightest box that the machine will pass?

**Solution:** Let  $w$  = the weight of the box, then:

$$|w - 14| = 0.3 \quad (\text{as per condition})$$

$$w - 14 = 0.3 \quad \text{or} \quad w - 14 = -0.3$$

$$w = 14.3 \quad \text{or} \quad w = 13.7$$


The heaviest weight allowed to pass is 14.3 kg. The lightest weight allowed to pass is 13.7 kg.

## Inequalities With Absolute Value

The absolute value of a number is its distance from 0 on the number line. We can use this idea to solve inequalities involving absolute value.

**Example:** Solve  $|x| < 5$ .

**Solution:**  $|x| < 5$  means that the distance between  $x$  and 0 is less than five units. To make  $|x| < 5$  true, we must substitute the values of  $x$  that are less than five units from 0.




All the numbers between  $-5$  and  $5$  are less than 5 units from zero.

The absolute solution set =  $\{|x|: -5 < x < 5\}$ .

**Example:** Solve  $|x| \geq 3$ . Graph the solution.

**Solution:** The solution of  $|x| \geq 3$  are those numbers whose distance from 0 is greater than or equal to 3. In other words, those numbers  $x$  such that  $x \leq -3$  or  $x \geq 3$ . The solution set is  $\{x|x \leq -3 \text{ or } x \geq 3\}$ .



**Example:** Solve and graph the solution.

a.  $|2x - 7| \geq 11$

b.  $|3x - 2| < 4$

**Solution:**

a.  $|2x - 7| \geq 11$  (Apply  $|x| \geq a: x \geq a$  or  $x \leq -a$ )


$$2x - 7 \geq 11 \quad \text{or} \quad 2x - 7 \leq -11$$

$$2x \geq 11 + 7 \quad \text{or} \quad 2x \leq -11 + 7$$

$$2x \geq 18 \quad \text{or} \quad 2x \leq -4$$

$$x \geq 9 \quad \text{or} \quad x \leq -2$$

$$\text{Solution set} = \{x|x \geq 9 \text{ or } x \leq -2\}$$



b.  $|3x - 2| < 4$  (Apply:  $|x| < a: -a < x < a$ )

$$-4 < 3x - 2 < 4$$


$$-4 < 3x - 2 \text{ and } 3x - 2 < 4$$

$$-4 + 2 < 3x \text{ and } 3x < 4 + 2$$

$$-2 < 3x \text{ and } 3x < 6$$

$$\frac{-2}{3} < x \text{ and } x < 2$$

$$\text{Solution set} = \left\{ x : \frac{-2}{3} < x < 2 \right\}$$



**Example:** Solve  $| -4x - 5 | + 3 < 9$ . Graph the solution.

**Solution:**  $| -4x - 5 | + 3 < 9 \Rightarrow | -4x - 5 | < 6$

$$-6 < -4x - 5 < 6$$

$$-6 < -4x - 5 \text{ and } -4x - 5 < 6$$

$$-1 < -4x \text{ and } -4x < 11$$


$$\frac{1}{4} > x \text{ and } x > \frac{-11}{4}$$

We can write as:  $-\frac{11}{4} < x < \frac{1}{4}$

$$\text{Solution set} = \left\{ x : -\frac{11}{4} < x < \frac{1}{4} \right\}$$

#### Key Fact:

- $|x| \leq a$  implies that  $-a \leq x \leq a$
- $|x| \geq a$  implies that  $x \leq -a$  or  $x \geq a$



**Example:**

You are buying a new calculator and find 10 models in a shop. The prices are Rs. 980, Rs. 570, Rs. 470, Rs. 820, Rs. 560, Rs. 770, Rs. 580, Rs. 390, Rs. 680 and Rs. 550.

- Find the mean of the calculator price.
- You are willing to pay the mean price with an absolute deviation of at most Rs. 100. How many of the calculator prices meet your condition?

**Solution:**

- Find the mean first:

$$\begin{aligned}\text{Mean} &= \frac{980+570+470+820+560+770+580+390+680+550}{10} \\ &= \frac{6380}{10} = 638\end{aligned}$$

- b. Write and solve an inequality.

An absolute deviation of at most Rs.100 from the mean 638, is given by the inequality.

$$|x - 638| \leq 100$$

$$-100 \leq x - 638 \leq 100$$

$$-100 \leq x - 638 \text{ and } x - 638 \leq 100$$

$$538 \leq x \text{ and } x \leq 738$$

We write as:  $538 \leq x \leq 738$

The price of calculator must be at least 538 and at most 738. Five prices meet your condition that is Rs. 570, Rs. 560, Rs. 580, Rs. 690 and Rs. 550.

### Exercise 4.3

Solve the absolute valued equations (1-6).

1.  $8|x - 3| = 88$

2.  $|2x + 9| = 30$

3.  $|x - 3| = -26$

4.  $3|x + 6| = 9x - 6$

5.  $|x - 3| = 2x$

6.  $\left|\frac{4-5x}{6}\right| = 7$

7. Explain, why the equation  $|3x - 6| + 7 = 4$  has no solution.

8. Before the start of a professional basketball game, a basketball must be inflated to an air pressure of 8 pounds per square inch (psi) (Absolute error is the absolute deviation). Find the minimum and maximum air pressure acceptable for the basketball.

Solve the inequality and graph the solution (9-14).

9.  $|9x - 1| \leq 10$

10.  $|2x - 7| < 1$

11.  $5\left|\frac{1}{2}x + 3\right| > 5$

12.  $3|14 - x| > 6$

13.  $\left|\frac{3x - 2}{5}\right| \leq 2$

14.  $\left|\frac{2-5x}{4}\right| \geq \frac{2}{3}$

15. An essay contest requires that essay entries consists of 500 words with an absolute deviation of at most 30 words. What are the possible number of words that the essay can have?

## Solving Systems of Linear Inequalities

A system of linear inequalities in two variables or simply a system of inequalities consists of two or more linear inequalities in the same variable. As:

$$\begin{aligned}x - y &> 9 \text{ Inequality I} \\2x + y &< 9 \text{ Inequality II}\end{aligned}\quad \left.\right\} A$$

A solution of a system of linear inequalities is an ordered pair that is a solution of each inequality in the system. For example,  $(7, -6)$  is a solution of the system A.

The graph of a system of linear inequalities is the graph of all solutions of the system.

**Example:** Graph the system of inequalities:

$$x + y > -2$$


$$3x - y \geq -6$$

**Solution:**

**Step I:** Graph both inequalities in the same coordinate plane.

**Step II:** Find the solution of each inequality. Preferably taking  $(0, 0)$  as test point to decide the solution region.

**Step III:** The graph of the system in the intersection of the two half planes. Which is shaded red.



**Check:** Choose a point in the region such as  $(0, 1)$ . To check this solution, substitute 0 for  $x$  and 1 for  $y$  into each inequality.

This dark shaded region is a solution of these inequalities.

Inequality I:

$$x + y > -2$$

$$0 + 1 > -2$$

$1 > -2$  True

Inequality II:

$$3x - y \geq -6$$

$$3(0) - 1 \geq -6$$

$-1 \geq -6$  True

### Key Fact:

- Graph each inequality in the cartesian plane.
- Find the intersection of the half planes.
- The graph of system is the intersection.

**Example:** Graph the system:


$$x + y \leq 4$$

$$x - y < 4$$

**Solution:**

To graph the inequality  $x + y \leq 4$ , we graph  $x + y = 4$  using a solid line. We then consider  $(0, 0)$  as a test point and find that it is a solution, we shade all points in that region.

Next, we graph  $x - y < 4$ . We graph  $x - y = 4$  as a dashed line and consider  $(0, 0)$  as a test point. Again  $(0, 0)$  is a solution, we shade that region. Finally, the solution set of the system is the region that is commonly shaded in both inequalities.



This dark shaded region is a solution of these inequalities.

**Example:** Graph the system of inequalities.

$$x \geq -2$$


$$x \leq 2$$

**Solution:**

Graph two inequalities in the same coordinate system.

$y = -2$  (The region is above the line)

$x = 2$  (The region is left of the line)



### Exercise 4.4

1. Describe the steps you would take to graph the system of inequalities shown.  
 $x - y < 6$ .....(Inequality I)  
 $y \geq 3$ .....(Inequality II)
2. Tell whether the ordered pair is a solution of the system of inequalities.
  - a.  $(3, 1), (0, 0), (-3, 1), (4, 1)$ ;  $x + y < 1, x - y < 2$
  - b.  $(1, -1), (4, 1), (2, 0), (3, 2)$ ;  $2x - y \leq 5, x + 2y > 2$

**3. Graph the system of inequalities.**

a.  $-x + y < 0$

$x + y > 3$

c.  $y - 2x \geq 3$

$y - 2x \leq 3$

e.  $x > -2$

$2x + y < 3$

$y \geq 2x = 1$

g.  $y < -x + 4$

b.  $x - y < 0$

$x + y < 1$

d.  $y \leq -2$

$x \geq 2$

f.  $x + y \leq 3$

$x - y \leq 4$

$x < 8$

h.  $x - 4y \leq -8$

**4. Does the system of inequalities have any solution?**

$x - y > 5$

$x - y < 1$

**5. Open-Ended Question:** Describe a real-world situation that can be modeled by a system of linear inequalities. Then write and graph the system of inequalities.

**I have Learnt**

- On a number line, the graph of an inequality in one variable is the set of points that represent all solutions of the inequality.
- To graph an inequality in one variable are an open circle for  $<$  or  $>$  and a closed circle for  $\leq$  or  $\geq$ .
- Dividing / multiplying each side of an inequality by a positive number produces an equivalent inequality.
- Dividing / multiplying each side of an inequality by a negative number and reversing the direction of the inequality symbol produces an equivalent inequality.
- A linear inequality in two variables, such as  $ax + by < c$  is the result of replacing the “=” sign in a linear equation with  $<$ ,  $\leq$ ,  $>$  or  $\geq$ .
- A solution of an inequality in two variables  $x$  and  $y$  is an ordered pair  $(x, y)$  that produces a true statement when the values of  $x$  and  $y$  are substituted into the inequality.
- The boundary line for an inequality in one variable is either vertical or horizontal.
- If an inequality has only the variable  $x$ , substitute the  $x$ -coordinate of the test point into the inequality.
- If an inequality has only the variable  $y$ , substitute the  $y$ -coordinate of the test point into the inequality.
- A solution of a system of linear inequalities is an ordered pair that is a solution of each inequality in the system.
- The graph of a system of linear inequalities is the graph of all solutions of the system.

### MISCELLANEOUS EXERCISE-4

**1.** Tick the correct option.

- The solution of the inequality  $6x - 7 \geq 2x + 17$  is:  
 a.  $x < 6$       b.  $x \leq 6$       c.  $x > 6$       d.  $x \geq 6$
- The solution of the inequality  $x < 3$  for  $x \in \mathbb{N}$  is:  
 a.  $\{0, 1, 2\}$       b.  $\{-2, -2, 0, 1, 2\}$       c.  $\{1, 2\}$       d.  $\{1\}$
- In general, we use a test point when graphing the inequality:  
 a.  $(-1, -1)$       b.  $(0, 0)$       c.  $(2, 2)$       d.  $(-3, -3)$
- The solution of the  $\frac{x}{-2} < 3 - x$  is:  
 a.  $x < 2$       b.  $x < 6$       c.  $x > 2$       d.  $x < -2$
- Which ordered pair is a solution of the inequality  $4x - y \geq 3$ ?  
 a.  $(0, 0)$       b.  $(-1, 2)$       c.  $(1, 1)$       d.  $(0, -2)$
- Which ordered pair is a solution of the system  $2x - y \leq 5$  and  $x + 2y > 2$ ?  
 a.  $(1, -1)$       b.  $(4, 1)$       c.  $(2, 0)$       d.  $(3, 2)$
- The solution for  $-5x \geq -80$  is:  
 a.  $\{x | x < 16\}$       b.  $\{x | x \leq 16\}$       c.  $\{x | x > 16\}$       d.  $\{x | x \geq 16\}$
- $(7, -2)$  is a solution of the inequality;  
 a.  $x - y < -4$       b.  $2x + y < 10$       c.  $x + 10y < 1$       d.  $-x - y > -3$

Solve the inequalities, if possible, graph the solution.

2.  $\frac{2}{3}x - 4 \geq 1$

4.  $7(x - 1) > -8 + 7x$

Graph on a plane.

6.  $y > 2x$

7.  $y < 3x$

Graph the system of inequalities:

10.  $3x + 4y \geq 12$

$5x + 6y \leq 30$

12.  $8x + 5y \leq 40$

$x \geq 0$

3.  $1 - 3x \leq -14 + 2x$

5.  $-3(2x - 1) \geq 1 - 8x$

8.  $y \geq -5$

9.  $x \leq 6$

11.  $y - x \geq 1$

$y - 4 \leq 4$

13.  $2x + y \leq 12$

$y \geq 0$

Write a system of inequalities for the shaded region described.

14. The shaded region is a rectangle with vertices  $(2, 1)$ ,  $(2, 4)$ ,  $(6, 4)$  and  $(6, 1)$ .

15. The shaded region is a triangle with vertices at  $(3, 0)$ ,  $(3, 2)$  and  $(0, -2)$ .

# 05

## Algebraic Fractions

**After studying this unit students will be able to:**

- Describe rational expression.
- Factorize and simplify rational expression.
- Demonstrate manipulation of algebraic fractions.
- Perform operations on rational expressions.
- Solve rational equations.
- Apply the concept of rational equations to real world problems.

Use of letters to represent an unknown quantity was introduced by "Rene Descartes", a French Mathematician, in 1637. Today 'x' is used by most nations as the standard letter for a single unknown. In fact x-rays were so named because the scientists who discovered them did not know what they were and thus labeled them the 'unknown rays' or x-rays. Algebra is basically an extension of Arithmetic.



X-ray

A factory worker wants to work overtime to fulfill his basic necessities of life. The factory owner has decided to give him Rs. 60 per hour for working extra. But sometimes he does not have enough time to work overtime daily. So he cannot find his accurate monthly income for working extra. If we represent overtime of the worker in hours for a month by an unknown quantity say 't', then we can say that his salary for working overtime will be '60t'.



# Algebraic Expressions

A statement in which variables or constants or both are connected by arithmetic operations (i.e.  $+$ ,  $-$ ,  $\times$ ,  $\div$ ) is called an algebraic expression.

For example,

$$\frac{-5x^2 + 4}{4}, \quad 3(a+b) - 4, \quad 0, \quad -5$$

$$z - \sqrt{2} t, \quad \frac{1}{x}, \quad \frac{x}{y+1}, \quad \sqrt{b^2 - 4ac} \text{ etc.}$$

## HISTORY:

Muhammad Bin Musa Al-Khawarizimi was the first Muslim mathematician who introduced Algebra and wrote a book entitled Hisab-Al-Jabr Wal Muqabala in 820 A.D. He was known as 'Father of Algebra'.

## Key Fact:

An algebraic expression may contain:

- Numbers
- Signs of operations ( $+$ ,  $-$ ,  $\times$ ,  $\div$ )
- Variables ( $a, b, c, \dots, x, y, z$ )
- Grouping symbols  $-$ ,  $( )$ ,  $\{ \}$ ,  $[ ]$

## Kinds of Algebraic Expressions

Algebraic expressions are of three kinds.

- Polynomial Expressions
- Rational Expressions
- Irrational Expressions

### (a) Polynomial Expressions (Polynomials)


Polynomials are algebraic expressions consisting of one or more terms in which exponents of the variables involved are non-negative (either zero or positive) integers.

For example:

$$0, \quad -2, \quad \frac{3}{4}x + \frac{3}{4}y^2z, \quad -\sqrt{\frac{3}{9}}y^3, \quad \sqrt{2}x^4 - \pi x^2 - \sqrt{10} \text{ etc.}$$

The expressions  $x^{-3}$ ,  $y^2 + \frac{1}{y^2}$ ,  $\sqrt[3]{y^4}$ ,  $2y^{\frac{1}{2}}$  are not polynomials because their exponents are not non negative integers (whole numbers).

### Classification of Polynomials



### (a) Types of Polynomials w.r.t. Terms

- A polynomial having one term is called a monomial. e.g., 3,  $x$ , 0 etc.
- A polynomial having two terms is called a binomial.

e.g.  $a + b$ ,  $2x + y$ ,  $7ab - \frac{1}{3}$ ,  $x^3 + 2\frac{1}{4}$  etc.

- A polynomial having three terms is called a trinomial.

e.g.  $x^2 + 2xy + y^2$ ,  $\frac{1}{8}x^3 + \frac{3}{4}x^2 - 9$  etc.

All other polynomials having more than three non-zero terms are called polynomials of four terms, five terms, six terms and so on.

#### Key Fact:

When a polynomial is written in descending order, then the first coefficient is called the leading coefficient. e.g., in  $p(x) = 3x^4 - 2x^2 + 1$ , the leading coefficient is 3. Here  $p(x)$  means that the variable in the polynomial is ' $x$ '.

### (b) Types of Polynomials w.r.t. Degree

#### ➤ Zero polynomial or no degree polynomial:

'0' is called a polynomial of no degree. Also,  $0x^3 + 0x$  is a no degree polynomial, because coefficients are always zero in zero polynomial.

#### ➤ Constant polynomial:

A polynomial having degree zero is called a constant polynomial. e.g.  $2, -5, \frac{1}{2}, \sqrt{5}$  are all constant polynomials.

#### ➤ Linear polynomial:

A polynomial having degree one is called a linear polynomial. e.g.,  $x, 2x - y, -7xy^0$  etc.

#### ➤ Quadratic polynomial:

A polynomial having degree two is called a quadratic

polynomial. e.g.,  $2x^2 + 7, ax + 2xy + 3, -\frac{3}{4}xyz^0$  etc.

#### ➤ Cubic polynomial:

A polynomial having degree three is called a cubic polynomial. e.g.  $9x^3 - 7x + 5, 9xzy, 3x^2y - \frac{3}{4}z$  etc.

All other polynomials have no specific name w.r.t. degree but simply, we call them polynomials of degree four, degree five and so on.

### (c) Types of Polynomials w.r.t. Coefficients

Polynomials can also be classified according to the coefficients of the variables involved in each term of them. e.g.,  $-3x^2 + 4x - 1$  is a polynomial with integers co-efficients because  $-3, 4, -1 \in \mathbb{Z}$ .

Similarly,  $\frac{1}{2}y^3 - \pi x^2 y + \frac{22}{7}$  is a polynomial with real numbers because  $\frac{1}{2}, -\pi, \frac{22}{7} \in R$ .

Just like this we can also check the coefficients in any polynomial and classify it accordingly.

#### (d) Types of Polynomials w.r.t. Variables

Classification of polynomials is also possible with respect to the variables involved in them. e.g.,  $x^2 y - 7$  is a polynomial in two variables  $x, y$ . We can express it as  $P(x, y) = x^2 y - 7$ .

Similarly,  $6x^2yz + 12xy^6z^6 + 9z^4$  is a polynomial in three variables  $x, y, z$ . We may write it as  $Q(x, y, z) = 6x^2yz + 12xy^6z^6 + 9z^4$ .

#### Recall

The highest exponent of the variable involved in a polynomial is called its degree. If more than one variable are being multiplied in terms of a polynomial. Then the degree of that polynomial is the sum of exponent of all the variables present in a term.

#### (b) Rational Expression

An algebraic expression of the form  $\frac{P(x)}{Q(x)}$ , where  $P(x)$  and  $Q(x)$  are polynomials and  $Q(x) \neq 0$

(i.e. it is not a zero polynomial) is called a rational expression.

For example:  $1\frac{1}{2}, \frac{3}{4x^2}, \frac{2x-1}{x^2+3}, \frac{2x+4}{x^2+5x+6}, \frac{x^2+x}{x^3+x}, 5$  etc.



Memory Plus:

#### Reduction / Simplification of Rational Expressions

In algebraic rational expressions, we can multiply or divide the numerator and denominator by a same non-zero expression.

Rational expression are the expression in which exponent of variable are integers.

$$\text{e.g. } \frac{x^2+x}{x^3+x} = \frac{(x^2+x) \div x}{(x^3+x) \div x} = \frac{x+1}{x^2+1}$$

Hence,  $\frac{x+1}{x^2+1}$  is the lowest or reduced form of  $\frac{x^2+x}{x^3+x}$ .

Reduction of a rational expression is done on the same rules as in reduction of a rational number. To reduce a rational expression to its lowest form, we have to

- Factorize the numerator and denominator completely.
- Divide the numerator and denominator by their common factors.
- Rewrite the remaining terms as a fraction, computing all products. This will be the reduced form.

**Example:** Reduce each rational expression to its lowest form.

(a)  $\frac{x^2 y^3}{x^5 y^3}$

(b)  $\frac{b-a}{a^3 - b^3}$

**Solution:** (i) By using laws of exponents, we have

$$\frac{x^2 y^3}{x^5 y^3} = \frac{y^{3-3}}{x^{5-2}}$$

$$= \frac{y^0}{x^3} = \frac{1}{x^3}$$

(ii) Since,  $(a^3 - b^3) = (a - b)(a^2 + ab + b^2)$

So, 
$$\begin{aligned}\frac{b-a}{a^3-b^3} &= \frac{(b-a)}{(a-b)(a^2+ab+b^2)} \\ &= \frac{-1}{(a-b)(a^2+ab+b^2)} \\ &= \frac{-1}{a^2+ab+b^2}\end{aligned}$$

### (c) Irrational Expression

An algebraic expression which cannot be expressed in the form  $\frac{P(x)}{Q(x)}$ , where  $P(x)$  and  $Q(x)$  are polynomials but  $Q$  is not a zero polynomial, is called an irrational expression.

For example:  $\frac{x}{\sqrt{y}}$ ,  $x^{\frac{3}{2}}y - 7$ ,  $\sqrt{x} - \frac{1}{\sqrt[3]{y}}$ ,  $\sqrt[5]{x^2 + y^2}$ ,  $24x^{\frac{3}{2}}y^{-2} + \frac{9}{y^2} - 7$  etc.

## Evaluation of Algebraic Expressions

The process of replacing the variables by numbers in an algebraic expression is called 'substitution'. The process of getting a numerical answer by substituting numbers for each variable is called 'evaluation'.

**Example:** Find the value of  $\frac{3}{4}x^2 - \frac{2}{3}x + 1$  at  $x = \frac{-1}{2}$ .

**Solution:**  $\frac{3}{4}x^2 - \frac{2}{3}x + 1$

By substituting the value of  $x$ , we have

$$= \frac{3}{4} \left( \frac{-1}{2} \right)^2 - \frac{2}{3} \left( \frac{-1}{2} \right) + 1$$

$$= \frac{3}{4} \left( \frac{1}{4} \right) - \frac{2}{3} \left( \frac{-1}{2} \right) + 1$$

### Check Point:

What will be the degree of polynomial  $\frac{x^3 y^2 - \sqrt{3} y^2 z^3 x}{xy^2}$  after simplification?

### Memory Plus:

Irrational expressions are those expression in which exponent of variable are non-integers.

$$= \frac{3}{16} + \frac{1}{3} + 1$$

$$= \frac{9+16+48}{48} = \frac{73}{48} = 1\frac{25}{48}$$

**Example:** If  $p = -5$  and  $q = 2$ , find the value of  $p^2q + pq^2 + 2p \cdot pq$

**Solution:**  $p^2q + pq^2 + 2p \cdot pq$

$$\begin{aligned} &= \underbrace{p^2q + pq^2 + 2p^2q}_{\leftarrow \text{simplify the given expression}} \\ &= 3p^2q + pq^2 \quad \leftarrow \text{combine like terms to reduce number of terms} \\ &= 3(-5)^2(2) + (-5)(2)^2 \leftarrow \text{substitute the given values} \\ &= 3(25)(2) + (-5)(4) \quad \leftarrow \text{simplify exponents} \\ &= 150 - 20 = 130 \end{aligned}$$

**Example:** Find the value of the expression

$$3.1x^4 - 0.2x^3y^2 + 0.3x^2y^2 + 1.1xy^3 - y^4 \text{ at } x = 1, y = -1.$$

**Solution:**  $3.1x^4 - 0.2x^3y^2 + 0.3x^2y^2 + 1.1xy^3 - y^4$

$$\begin{aligned} &= 3.1(1)^4 - 0.2(1)^3(-1)^2 + 0.3(1)^2(-1)^2 + 1.1(1)(-1)^3 - (-1)^4 \\ &= 3.1 - 0.2(1)(+1) + 0.3(1)(+1) + 1.1(1)(-1) - (+1) \\ &= 3.1 - 0.2 + 0.3 - 1.1 - 1 \\ &= 3.4 - 2.3 = 1.1 \end{aligned}$$

### Exercise 5.1

1. Reduced the following expressions to their lowest form.


<b>(i)</b>	$\frac{15ax^3y^2}{25a^2xy^6}$	<b>(ii)</b>	$\frac{38k^2p^3m^4}{57k^3pm^2}$	<b>(iii)</b>	$\frac{mn^4pq}{m^2n^3p^4}$
<b>(iv)</b>	$\frac{3abc}{15a^2b^2c}$	<b>(v)</b>	$\frac{46l^3m^4n^5}{69l^2m^3n^4}$	<b>(vi)</b>	$\frac{x-3}{3-x}$
<b>(vii)</b>	$\frac{x^2-81}{x+9}$	<b>(viii)</b>	$\frac{(r+3)(r+4)}{r^2-16}$		

2. Evaluate the following expressions for the given value of each variable.

<b>(i)</b> $3(r^2 - s^2)$ , if $r = 2, s = -1$	<b>(ii)</b> $\frac{1}{2}mv^2$ at $m = 18.75$ and $v = 5.6$
<b>(iii)</b> $\sqrt{2gs}$ when $g = 32.2$ and $s = 144.9$	

- (iv)  $3x - y + \frac{1}{z}$  if  $x = \frac{-1}{2}$ ,  $y = 3$ ,  $z = \frac{-1}{3}$
- (v)  $0.1d^2 + 0.01d + 1$  if  $d = -0.2$  (vi)  $\frac{4}{7}b^3 - 3\frac{1}{2}b^2 + b - 3$  if  $b = \frac{1}{2}$
3. If  $n$ th triangular number is represented by  $T(n) = \frac{n(n+1)}{2}$ , then find  $100^{\text{th}}$  triangular number.
4. If  $P(x) = x^2 + 2x - 15$ ,  $D(x) = x - 3$  and  $Q(x) = x + 5$ , show that  $\frac{P(2)}{Q(2)} = D(2)$ .
5. If  $g(x) = \frac{1}{2x^3} + \frac{x}{2} + 2$ , find  $g\left(-\frac{1}{3}\right)$ .
6. The volume of a basketball (sphere) is approximately  $38808 \text{ cm}^3$ . The radius 'r' of the ball is given by

$$r = \sqrt[3]{\frac{3v}{4\pi}}, \text{ where } v \text{ is its volume.}$$



Determine the radius of that ball. (take  $\pi = \frac{22}{7}$ )

## Basic Operations on Algebraic Fractions

Operations with rational numbers and rational algebraic expressions are very similar. A rational number can be expressed as the quotient of two integers. A rational algebraic expression can be expressed as the quotient of two polynomials. In either case, the denominator can never be 0.

Rational Numbers:  $\frac{3}{11}, \frac{27}{100}, \frac{-7}{25}, \frac{61}{-89}$

Rational Algebraic Expressions:  $\frac{5}{x}, \frac{y}{10}, \frac{2x}{x-5}, \frac{y^2-36}{y+6}, \frac{x^2-6x+5}{\frac{5x}{x+1}}$

If P and Q are any two algebraic expressions then  $\frac{P}{Q}$  is called an algebraic fraction,

where  $Q \neq 0$  e.g.,  $\frac{a-b}{a^2+2ab+b^2}$

To write a fraction in simplest form, we divide both the numerator and denominator by their greatest common factor (GCD).

## Addition of Algebraic Expressions

Algebraic expressions can be added as explained in the following examples.

**Example:** Add the following expressions.

$$(i) \quad \frac{x^2 - 1}{2}, \quad \frac{4(2x^2 - x + 1)}{3} \quad (ii) \quad 4x + 7, \quad \frac{(x+3)(x-1)}{x-2}$$

**Solution:**

$$\begin{aligned}(i) \quad & \frac{x^2 - 1}{2} + \frac{4(2x^2 - x + 1)}{3} \\&= \frac{3(x^2 - 1) + 8(2x^2 - x + 1)}{6} \\&= \frac{3x^2 - 3 + 16x^2 - 8x + 8}{6} \\&= \frac{19x^2 - 8x + 5}{6}\end{aligned}$$

$$\begin{aligned}(ii) \quad & 4x + 7 + \frac{(x+3)(x-1)}{x-2} \\&= \frac{(4x+7)(x-2) + (x+3)(x-1)}{x-2} \\&= \frac{4x^2 + 7x - 8x - 14 + x^2 + 3x - x - 3}{x-2} \\&= \frac{5x^2 + x - 17}{x-2}\end{aligned}$$

### Key Fact:

For all rational expressions

$$\frac{a}{b} \text{ and } \frac{c}{d}, b \neq 0, d \neq 0 :$$

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd}, \quad \frac{a}{b} - \frac{c}{d} = \frac{ad - bc}{bd}$$

## Subtraction of Algebraic Expressions

In subtraction of algebraic expressions, we have to change the signs of the expression which is being subtracted and all other steps are the same as in addition of algebraic expressions.

**Example:** Subtract  $\frac{3x^2 - 7x + 4}{x^2 - 4}$  from  $\frac{3x - 1}{x + 2}$ .

$$\begin{aligned}\text{Solution: } & \frac{3x - 1}{x + 2} - \frac{3x^2 - 7x + 4}{x^2 - 4} \\&= \frac{3x - 1}{x + 2} - \frac{3x^2 - 7x + 4}{(x+2)(x-2)} \quad \leftarrow \text{By using the formula } a^2 - b^2 = (a+b)(a-b) \\&= \frac{(3x-1)(x-2) - (3x^2 - 7x + 4)}{(x+2)(x-2)} \\&= \frac{3x^2 - 7x + 2 - 3x^2 + 7x - 4}{(x+2)(x-2)} = \frac{-2}{(x+2)(x-2)} = \frac{-2}{x^2 - 4}\end{aligned}$$

## Multiplication of Algebraic Expressions

Multiplication of algebraic expression can be explained with the help of following examples.

**Example:** Find the product of  $\frac{x^2 - 9}{x^2 + 2x}$  and  $\frac{x}{x+3}$ .

**Solution:** 
$$\begin{aligned} & \frac{x^2 - 9}{x^2 + 2x} \times \frac{x}{x+3} \\ &= \frac{(x+3)(x-3)}{x(x+2)} \times \frac{x}{x+3} = \frac{x-3}{x+2} \end{aligned}$$

**Example:** Multiply  $x^2 - x - 20$  by  $\frac{3}{x^3 + 4x^2}$ .

**Solution:** 
$$\begin{aligned} & (x^2 - x - 20) \times \frac{3}{x^3 + 4x^2} \\ &= (x^2 - 5x + 4x - 20) \times \frac{3}{x^2(x+4)} \\ &= (x-5)(x+4) \times \frac{3}{x^2(x+4)} = \frac{3x-15}{x^2} \end{aligned}$$

### Key Fact:

For all rational expressions

$\frac{a}{b}$  and  $\frac{c}{d}$ ,  $b \neq 0$ ,  $d \neq 0$ :

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{ac}{bd}, \quad \frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c} = \frac{ad}{bc}$$

## Division of Algebraic Expressions

The process of dividing any two algebraic expressions is explained with the help of following examples.

**Example:** Divide  $\frac{2x^2 - 11x - 21}{x^2 - 49}$  by  $\frac{2x+3}{x+7}$ .

**Solution:** 
$$\begin{aligned} & \frac{2x^2 - 11x - 21}{x^2 - 49} \div \frac{2x+3}{x+7} \\ &= \frac{2x^2 - 14x - 3x - 21}{(x)^2 - (7)^2} \times \frac{x+7}{2x+3} \\ &= \frac{2x(x-7) + 3(x-7)}{(x+7)(x-7)} \times \frac{x+7}{2x+3} \\ &= \frac{(x-7)(2x+3)}{(x+7)(x-7)} \times \frac{x+7}{2x+3} = 1 \end{aligned}$$

## Complex Algebraic Expressions

For simplifying the complex algebraic expression involving addition, subtraction, multiplication and division, we apply the BODMAS rule similar to arithmetic expression.

**Example:** Simplify.

$$\frac{3x+2}{x+2} + \frac{x-2}{2x+10} \div \frac{2x^2-4}{x^2+2x-15}$$

**Solution:**

$$\begin{aligned} & \frac{3x+2}{x+2} + \frac{x-2}{2x+10} \times \frac{x^2+2x-15}{2x^2-8} \\ &= \frac{3x+2}{x+2} + \frac{x-2}{2(x+5)} \times \frac{x^2+5x-3x-15}{2(x^2-4)} \\ &= \frac{3x+2}{x+2} + \frac{x-2}{2(x+5)} \times \frac{x(x+5)-3(x+5)}{2(x+2)(x-2)} \\ &= \frac{3x+2}{x+2} + \frac{x-2}{2(x+5)} \times \frac{(x+5)(x-3)}{2(x+2)(x-2)} \\ &= \frac{3x+2}{x+2} + \frac{x-3}{4(x+2)} \\ &= \frac{4(3x+2) + (x-3)}{4(x+2)} = \frac{12x+8+x-3}{4(x+2)} = \frac{13x+5}{4x+8} \end{aligned}$$

**Example:** Simplify the following.

$$\left[ \frac{a}{(a+b)^2 - 2ab} \div \frac{(a+b)^3 - 3ab(a+b)}{a^4 - b^4} \right] \div \frac{(a+b)^2 - 4ab}{(a+b)^2 - 3ab}$$

**Solution:**

$$\begin{aligned} & \left[ \frac{a}{(a+b)^2 - 2ab} \div \frac{(a+b)^3 - 3ab(a+b)}{a^4 - b^4} \right] \div \frac{(a+b)^2 - 4ab}{(a+b)^2 - 3ab} \\ &= \left[ \frac{a}{(a+b)^2 - 2ab} \times \frac{a^4 - b^4}{(a+b)^3 - 3ab(a+b)} \right] \div \frac{(a+b)^2 - 4ab}{(a+b)^2 - 3ab} \\ &= \left[ \frac{a}{a^2 + b^2 + 2ab - 2ab} \times \frac{(a^2 - b^2)(a^2 + b^2)}{a^3 + b^3 + 3ab(a+b) - 3ab(a+b)} \right] \div \frac{(a+b)^2 - 4ab}{(a+b)^2 - 3ab} \\ &= \left[ \frac{a}{a^2 + b^2} \times \frac{(a^2 - b^2)(a^2 + b^2)}{a^3 + b^3} \right] \div \frac{(a+b)^2 - 4ab}{(a+b)^2 - 3ab} \\ &= \left[ \frac{a}{a^2 + b^2} \times \frac{(a+b)(a-b)(a^2 + b^2)}{(a+b)(a^2 - ab + b^2)} \right] \div \frac{(a+b)^2 - 4ab}{(a+b)^2 - 3ab} \\ &= \left[ \frac{a(a-b)}{a^2 - ab + b^2} \div \frac{(a+b)^2 - 4ab}{(a+b)^2 - 3ab} \right] \end{aligned}$$

$$\begin{aligned}
 &= \left[ \frac{a(a-b)}{a^2-ab+b^2} \times \frac{(a+b)^2 - 3ab}{(a+b)^2 - 4ab} \right] \\
 &= \left[ \frac{a(a-b)}{a^2-ab+b^2} \times \frac{a^2 + 2ab + b^2 - 3ab}{a^2 + 2ab + b^2 - 4ab} \right] \\
 &= \left[ \frac{a(a-b)}{a^2-ab+b^2} \times \frac{a^2 + b^2 - ab}{a^2 + b^2 - 2ab} \right] \\
 &= \left[ \frac{a(a-b)}{a^2 - 2ab + b^2} \right] \\
 &= \left[ \frac{a(a-b)}{(a-b)^2} \right] \\
 &= \frac{a}{a-b}
 \end{aligned}$$

Another method for simplifying a complex rational expression involves first adding or subtracting, as necessary, to get one rational expression in the numerator and one rational expression in the denominator. The problem is thereby simplified to one involving the division of two expressions.

**Example:** Simplify.  $\frac{\frac{3}{x} - \frac{2}{x^2}}{\frac{3}{x-2} + \frac{1}{x^2}}$

**Solution:** 
$$\frac{\frac{3}{x} - \frac{2}{x^2}}{\frac{3}{x-2} + \frac{1}{x^2}} = \frac{\frac{3}{x} - \frac{2}{x^2}}{\frac{3x^2 + x - 2}{x^2(x-2)}}, \text{ taking LCM}$$

$$= \frac{3x-2}{x^2} \times \frac{x^2(x-2)}{3x^2+x-2} \text{ to divide, multiply, by the reciprocal of the divisor.}$$

$$= \frac{3x-2}{x^2} \times \frac{x^2(x-2)}{3x^2+3x-2x-2} = \frac{(3x-2)}{x^2} \times \frac{x^2(x-2)}{(3x-2)(x+1)} \text{ factoring}$$

$$= \frac{x-2}{x+1} \text{ removing a common factor from numerator and denominator.}$$

## Exercise 5.2

**1.** Add the following expressions.

(i)  $\frac{x}{2}, \frac{x}{5}$       (ii)  $\frac{x-2}{2}, \frac{x+10}{9}$       (iii)  $\frac{4+x}{4}, \frac{x-1}{7}, \frac{5x}{2}$

(iv)  $\frac{3x}{x+5}, \frac{10}{5x+25}$       (v)  $\frac{24x}{6x-18}, \frac{3(1+x)}{x-3}$

**2.** Subtract.

(i)  $\frac{23-x}{5}$  from 7      (ii)  $\frac{6(x-8)}{7}$  from  $\frac{5(x-7)}{3}$

(iii)  $2x^2 - 2x + 1$  from  $\frac{x+1}{x}$ .

**3.** Divide the first expression by the second.

(i)  $x^4 - 10x^2 + 9, x^2 - 2x - 3$       (ii)  $x^3 - 3x^2y + 3xy^2 - y^3, x - y$

(iii)  $\frac{4x^2 - 16}{5x}, \frac{2x+4}{15}$       (iv)  $\frac{x^2 + 5x}{x-3}, \frac{x^2 - 25}{x-3}$

**4.** Simplify the following.

(i)  $\frac{x}{2} + \frac{x}{3} - \frac{x}{4} + \frac{x}{5}$       (ii)  $\frac{1}{2}\left(4 - \frac{x}{3}\right) - \frac{5}{6} + \frac{1}{3}\left(11 - \frac{x}{2}\right)$

(iii)  $\frac{2}{x+1} + \frac{x}{x-1} - \frac{x+2}{x-1}$       (iv)  $\frac{x^2 - 25}{5} - \frac{x}{4} \div \frac{3x}{20}$

(v)  $\frac{45a^2b^3c^4}{27x^4y^3z} \times \frac{243xy^2z^3}{180a^2bc^3}$       (vi)  $\frac{m^2}{8n} \times \frac{36p^3q^2}{81mn} \div \frac{15mpx^5}{270n^2x^3y}$

(vii)  $3x \div \frac{3x^2 - 27}{x+3} + \frac{1}{x-3}$       (viii)  $\frac{5x+5}{3(2x-1)} + \frac{6-2x}{2(1-2x)}$

(ix)  $\frac{2a}{2a-3} - \frac{5}{6a+9} - \frac{4(3a+2)}{3(4a^2-9)}$       (x)  $\frac{5}{5+x-18x^2} - \frac{2}{2+5x+2x^2}$

(xi)  $\frac{m+3}{24m} - \frac{m+1}{24m} + \frac{3m-1}{6m^2+18m} \div \frac{12m-4}{m+3}$

(xii)  $\frac{1-p^2}{1+q} \times \frac{1-q^2}{p+p^2} \times \left(1 + \frac{p}{1-p}\right)$       (xiii)  $\left(1 - \frac{x}{1+\frac{x}{1-x}}\right) \div (1+x^3)$

## Solving Rational Equations

An equation that contains one or more rational expressions is called a rational equation. Here are some examples:

$$\frac{2}{5} - \frac{x}{4} = \frac{1}{x}, \quad \frac{x-4}{x+1} - \frac{6}{x-2} = \frac{7}{x+1}, \quad x^2 + \frac{5}{x} = 9$$

Equations of this type occur frequently in applications. It is easiest to solve a rational equation if the fractions are eliminated. This can be done by multiplying each side of the equation by the least common denominator (LCD). Remember that when you multiply each side by the LCD, each term on each side must be multiplied by the LCD.

**Example:** Solve  $\frac{3}{x-7} + 1 = \frac{8}{x^2 - 9x + 14}$ . Verify the solution.

**Solution:**  $\frac{3}{x-7} + 1 = \frac{8}{x^2 - 9x + 14}$

$$(x-2)(x-7) \cdot \frac{3}{x-7} + (x-2)(x-7) \cdot 1 = (x-2)(x-7) \cdot \frac{8}{(x-2)(x-7)}$$
 Multiply by LCD  $(x-2)(x-7)$ .

$$3(x-2) + (x^2 - 9x + 14) = 8 \text{ Simplify.}$$

$$x^2 - 6x + 8 = 8 \text{ Combine the like terms.}$$

$$x^2 - 6x = 0 \text{ Simplify.}$$

$$x(x-6) = 0$$

$x = 0$ , or  $x = 6$ . The solutions are 0 and 6.

**Check:** if  $x = 0$ ,  $\frac{3}{0-7} + 1 = \frac{8}{0^2 - 9.0 + 14}$ , we have:  $\frac{4}{7} = \frac{4}{7}$ , True.

if  $x = 6$ ,  $\frac{3}{6-7} + 1 = \frac{8}{6^2 - 9.6 + 14}$ , we have:  $-2 = -2$ , True.

$$S.S = \{0, 6\}$$

**Note:** Remember that a rational expression is undefined when the value for variables results in a denominator of zero. When solving rational equations, we check that for solution denominator should be non-zero. If denominator is zero in any value of  $x$ , these values be excluded from list of solutions.

**Example:** Solve  $\frac{x}{x-2} + \frac{1}{5} = \frac{2}{x-2}$ .

**Solution:**  $\frac{x}{x-2} + \frac{1}{5} = \frac{2}{x-2}$ .

$$5(x-2) \cdot \frac{x}{x-2} + 5(x-2) \cdot \frac{1}{5} = 5(x-2) \cdot \frac{2}{x-2}$$
 Multiply by LCD  $5(x-2)$ .

$$5x + x - 2 = 10 \text{ Simplify.}$$

$$6x - 2 = 10 \text{ Combine the like terms.}$$

$$x = 2$$

The solution appears to be 2, but the expression  $\frac{x}{x-2} + \frac{1}{5} = \frac{2}{x-2}$  are undefined when  $x=2$ .

We see that 2 is extraneous solution.

There is no solution.

**Example:** If a certain number is added 5 times the reciprocal of 2 more than that number, the result is 4.

**Solution:** Suppose a number is  $x$ . According to condition, added to 5 times the reciprocal of 2 more than that number, the result is 4:

$$x + 5\left(\frac{1}{x+2}\right) = 4 \quad \text{As per statement, written in mathematical form}$$

$$(x+2).x + (x+2).5\left(\frac{1}{x+2}\right) = (x+2).4 \quad \text{Multiply by LCD on both sides.}$$

$$x^2 + 2x + 5 = 4x + 8 \quad \text{Simplify}$$

$$x^2 - 2x - 3 = 0 \quad \text{Collecting all like terms on one side}$$

$$(x-3)(x+1) = 0 \quad \text{Factoring}$$

$x = 3$  or  $x = -1$ , the solution can be checked.

The solution can be verified:

$$x + 5\left(\frac{1}{x+2}\right) = 4 \quad (\text{As per statement, written in mathematical form})$$

$$(x+2).x + (x+2).5\left(\frac{1}{x+2}\right) = (x+2).4 \quad (\text{Multiply by LCD on both sides.})$$

$$x^2 + 2x + 5 = 4x + 8 \quad (\text{Simplify})$$

$$x^2 - 2x - 3 = 0 \quad (\text{Collecting all like terms on one side})$$

$$(x-3)(x+1) = 0 \quad (\text{Factoring})$$

$x = 3$  or  $x = -1$ , the solution can be checked.

**Result:** The required number is  $-1$  or  $3$ .

**Example:** A car travels 300 km in the same time that a freight train travels 200 km. The speed of the car is 20 km/h more than the speed of the train. Find the speed of the car and the speed of the train.

**Solution:** We know that distances both vehicles traveled. We also observe that both vehicles traveled the same amount of time and that the speed of the car is 20 km/h faster than that of the train.

Suppose the distance is  $S$  and speed is  $V$ , we know that  $S=VT$ , where  $T$  is time. If  $V$  be the speed of the train, then car moves 20 km/h faster than the speed of the train. So, the speed of

the car is  $V+20$ . We can equate the formula for the two vehicles in terms of  $T$ . We have:

$$T = \frac{S}{V}$$

car's time = train's time

$$\frac{S}{V} = \frac{300}{V+20} = \frac{200}{V}$$

$$\frac{300}{V+20} = \frac{200}{V} \quad (\text{Solve for } V)$$

$$(V+20)V \cdot \frac{300}{V+20} = V(V+20) \cdot \frac{200}{V} \quad (\text{Multiply each side by LCD})$$

$$300V = 200(V+20) \quad (\text{Simplify})$$

$$100V = 4000$$

$$V = 40$$

The speed of the train is 40 km/h and the speed of the car is 60 km/h.

### Exercise 5.3

Solve and check your solutions (1-7).

$$1. \frac{6x}{x-11} + 1 = \frac{3}{x-11}$$

$$2. \frac{2y}{y+3} = \frac{-4}{y-7}$$

$$3. \frac{x+7}{x+4} - 1 = \frac{x+10}{2x+8}$$

$$4. \frac{3y}{y+1} = \frac{12}{y^2-1} + \frac{y+4}{y+1}$$

$$5. x + \frac{5}{x} = -6$$

$$6. \frac{y+2}{y^2+6y-7} = \frac{8}{y^2+3y-4}$$

$$7. \frac{5}{y+1} + \frac{3y+5}{y^2+4y+3} = \frac{2}{y+3}$$

8. Kaleem can mow a lawn in 4 hours. Moiz can mow the same lawn in 5 hours.

How long would it take both of them, working together, to mow the lawn.

9. You have an 8-pint mixture of paint that is made up of equal amounts of yellow paint and blue paint. To create a certain shade of green, you need a paint mixture that is 80 % yellow. How many pints of yellow paint do you need to add to the mixture?
10. Waqar takes 9 hours longer to build a wall than it takes Wasi. If they work together, they can build the wall in 20 hours. How long would it take each, working alone, to build the wall?

- Describing rational expression.
  - Factorizing and simplifying rational expression.
  - Demonstrating manipulation of algebraic fractions.
  - Performing operations on rational expressions.
  - Solving rational equations.
  - Applying the concept of rational equations to real world problems.

## MISCELLANEOUS EXERCISE-5

- vii. If  $y = 2 - \frac{1}{y}$ , then the value of  $y^2 + \frac{1}{y^2}$  is  
 (a) 4      (b) zero      (c) not possible      (d) 2
- viii. Simplified from of  $\frac{(a+b)^2 - (a-b)^2}{8ab}$  is  
 (a)  $\frac{2(a^2 + b^2)}{8ab}$       (b) 2      (c)  $\frac{a^2 + b^2}{+ ab}$       (d)  $\frac{1}{2}$
- ix. Difference of the sum of a and b from the product of a and b is  
 (a)  $ab - a - b$       (b)  $a + b - ab$       (c)  $2ab - b$       (d) none
- x. 
$$\frac{x^3y^3 + y^3z^3 + z^3x^3}{x^3y^3z^3} =$$
  
 (a)  $x^3 + y^3 + z^3$       (b)  $x^6 + y^6 + z^6$   
 (c)  $\frac{1}{x^3} + \frac{1}{y^3} + \frac{1}{z^3}$       (d)  $\frac{1}{x^6} + \frac{1}{y^6} + \frac{1}{z^6}$
- xi. Leading coefficient in  $\frac{x^2}{2} - \frac{1}{8} - \frac{x^4}{4} + \frac{x^3}{7}$  is  
 (a)  $\frac{1}{2}$       (b)  $\frac{1}{4}$       (c)  $\frac{1}{7}$       (d)  $-\frac{1}{4}$
- xii. Coefficients in the polynomial  $\sqrt{16}x^2y - \frac{1}{2}y^3 + \frac{22}{7}z$  are the elements of the set of:  
 (a) Integers      (b) Irrational numbers  
 (c) Odd numbers      (d) Rational numbers
- xiii. The degree of the quotient in  $(x-y)^3 \div (x-y)^2$  will be  
 (a) 3      (b) 2      (c) 1      (d) no
2. Find the value of the circumference of a circle whose radius is 12.5 cm.  
 (Take  $\pi = \frac{22}{7}$ )
3. The surface S of a sphere of radius 'r' is given by the formula  $S = 4 \times \frac{22}{7} r^2$ , find  
 (i) The surface of a sphere whose radius is 1.4 inches;  
 (ii) The radius of a sphere whose surface is  $38\frac{1}{2}$  square feet.


4. In a right-angled triangle, if  $a$  and  $b$  denote the lengths of the sides containing the right angle and  $c$  denotes the length of the hypotenuse, it is known that  $c^2 = a^2 + b^2$ . By substitution find which of the following sets of numbers represent the sides of right-angle triangle.
- (i) 7, 24, 25      (ii) 1.6, 6.3, 6.5  
(iii) 12, 35, 36      (iv) 3, 4, 5
5. If  $a = 3$ ,  $b = 4$  and  $c = 1$ , find the value of:  $\sqrt{2ab + 4ac} + \sqrt{9b} + \frac{2abc}{3}$
6. Subtract the sum of:  $2x^3 - 3x + 4$  and  $-3x^2 + 2x - 7$  from  $4x^3 - 3x^2 + x - 6 - \{2x^3 - (x - 6)\}$
7. Divide the product of  $x - 2$ ,  $x + 3$  and  $2x - 7$  by the sum of  $3(x^2 - 2x - 2)$  and  $5x - x^2 - 15$ .
8. Simplify the following.
- (i)  $\frac{x}{x+2} - \frac{5x+3}{x-2} + \frac{1}{2}$       (ii)  $\frac{x}{x^2-y^2} \times \frac{x^2+2xy+y^2}{x+y} \div \frac{3x}{x-y}$
9. Solve.
- (i)  $\frac{12}{x^2-16} - \frac{24}{x-4} = 3$       (ii)  $\frac{y}{2y-6} - \frac{3}{x^2-6x+9} = \frac{y-2}{3y-9}$
10. When checking a possible solution of a rational equation, is it necessary to check that the solution does not make any denominator equal 0? Why or why not?

# Functions And Graphs

After studying this unit students will be able to:

- Recognize function as a rule of correspondence and find domain and range of a function.
- Identify types of functions (into, onto, one-to-one, injective, surjective and bijective) by using Venn diagrams.
- Determine value of a function and perform operations on functions.
- Find inverse of a function.
- Formulate composite functions as:  $fg(x) = f(g(x))$ .
- Identify, sketch and interpret graphs of constant function, identity function, linear and non-linear function, absolute valued function and function of the type  $y = x^n$  where  $n$  is a + ve or - ve integer and  $x > 0$
- Determine gradient of the curve through tangent line.
- Solve a system of one linear and one quadratic equation graphically.
- Solve real life problems related to functions.

Every machine needs some input to perform its functions. When we go shopping, we pay our bill at the counter. The cashier inputs the prices of all the items purchased into the calculating machine. The machine performs its function and processes out a bill. This bill is the output produced by the machine. It means every machine needs some input to give some output, i.e; every output is related to some input.



Our calculator is also a machine, we input some values in it and it gives us the result after performing some process.

Cartesian product of any two non-empty sets A and B denoted by  $A \times B$  (read as A cross B) is a set containing all ordered pairs  $(x, y)$  such that  $x \in A$  and  $y \in B$ . We can write it as:

$$A \times B = \{(x, y) \mid x \in A \wedge y \in B\}$$

For example, if  $A = \{a, b\}$  and  $B = \{2\}$ , then

$$A \times B = \{(a, 2), (b, 2)\}$$

All the subsets of  $A \times B$  are:

$$R_1 = \{\}, R_2 = \{(a, 2)\}, R_3 = \{(b, 2)\} \text{ and } R_4 = \{(a, 2), (b, 2)\}$$

Any subset of the cartesian product  $A \times B$  is called a binary relation (or simply a relation) from set A to set B. Basically, a binary relation relates an element of one set to an element of the other set.

All the above four subsets are binary relations from set A to set B.

Domain of any relation R denoted by  $\text{dom}(R)$  is a set containing first element of each ordered pair in the relation.  $\text{Dom}(R_4) = \{a, b\}$

Range of a relation R denoted by  $\text{ran}(R)$  is a set containing second element of each ordered pair in the relation.  $\text{Ran}(R_4) = \{2\}$

In  $R_4$ , 2 is image of a and b but a and b are preimages of 2

#### Key Fact:

- If  $n(A) = p$  and  $n(B) = q$ , then  $n(A \times B) = pq$
- Number of subsets of  $A \times B = 2^{n(A \times B)} = 2^{pq}$

## Function

A function f from set A to set B is a relation, rule or mapping which maps each element of set A to a unique element of set B such that domain of  $f = A$  and there is no repetition in the first element of ordered pairs in f.

Mathematically, it is written as  $f : A \rightarrow B$  and is read as f is a function from A to B.

In the above example in which  $A = \{a, b\}$  and  $B = \{2\}$ ,

Consider  $R_4 = \{(a, 2), (b, 2)\}$ ,

which is shown in arrow diagram at right.


Domain of  $R_4 = \{a, b\} = A$

There is no repetition in the first element of ordered pairs in  $R_4$ .

Therefore,  $R_4$  is a function.

The first element of each ordered pair in  $R_4$  is called pre-image while its corresponding second element is called image of the first element.

In  $(a, 2)$ , a is the pre-image of 2 and 2 is the image of a.



#### Check Point:

$R_1, R_2$  and  $R_3$  are not functions. Justify.

### Domain of a Function

A set containing first element of each ordered pair in the function is called its domain.

### Range of a Function

A set containing second element of each ordered pair in the function is called its range.

#### Key Fact:

If f is a function from set A to set B, i.e.,  $f : A \rightarrow B$ , then B is called co-domain of f and range of f is a subset of co-domain.

**Example:** If  $A$  = Set of first three natural numbers

$B$  = Set of prime factors of 6







Is  $f = \{(1, 2), (2, 3), (3, 2)\}$  a function from  $A$  to  $B$ ?

**Solution:**  $A = \{1, 2, 3\}$ ,  $B = \{2, 3\}$

Here  $f$  is a subset of  $A \times B$  and  $\text{dom } f = \{1, 2, 3\} = A$ .

Also, there is no repetition in the first element of the ordered pairs in  $f$ , therefore  $f$  is a function.

**Example:** Which of the following relations indicated by the given figures are functions?



**Solution:** Here  $f$ ,  $g$  and  $h$  are functions as the domain of each of them is equal to their corresponding first set and no first element is repeated in their respective mapping.

But in mapping  $t$ ,  $\text{dom}(t) \neq P$  ( $\because$  2 has no image in set  $Q$ ) and there is repetition of 0 ( $\because$  it is mapped to two elements 7 and 9 of  $Q$ ), therefore  $t$  is not a function.

In mapping  $p$ ,  $\text{dom}(p) \neq L$  ( $\because$  b has no image in set  $M$ ), so  $p$  is not a function.

In mapping  $q$ , although  $\text{dom}(q) = S$  but b is repeated ( $\because$  it is the pre-image of two elements 2 and 3 of  $T$ ), therefore  $q$  is not a function.

## Types of Functions

Range of a function is required for identifying the type of a function.

For any two non-empty sets  $A$  and  $B$ , these are the five types of functions:


### Into Function

A function  $f : A \rightarrow B$  will be into if there is atleast one element in set  $B$  which is not an image of any element of set  $A$ .

i.e.,  $\text{range}(f) \subset B$ .

In the adjoining arrow diagram,  $f = \{(a, 1), (b, 4), (c, 3)\}$  is an into function.

**Key Fact:**  
In into function,  $\text{range } f \neq B$ .



## Onto Function

A function  $f : A \rightarrow B$  will be onto if every element of set B is an image of atleast one element of set A. i.e,  $\text{range}(f) = B$ .


In the adjoining arrow diagram,  $f = \{(a, 1), (b, 2), (c, 2)\}$  is an onto function.

An onto function is also called a surjective function.

## One-to-One Function

A function  $f : A \rightarrow B$  will be one-to-one if distinct elements of set A have distinct images in set B.

In the adjoining arrow diagram,  $f = \{(a, 2), (b, 4), (c, 1)\}$  is a one-to-one function.



### Points To Ponder!

- We can write one-to-one function as one-one or simply 1-1 function.
- There should be no repetition in the second element of the ordered pairs in a 1-1 function.
- Range of a one-one function may not be equal to set B.

## Injective Function

A function which is into as well as 1-1 is called an injective function.


In the above arrow diagram,  $f = \{(a, 2), (b, 4), (c, 1)\}$  is an injective function.

## Bijective Function

A function which is both onto and 1-1 is called a bijective function.

In the arrow diagram at right,

$f = \{(a, 2), (b, 1), (c, 3)\}$  is a bijective function.



Bijective function shows one-one correspondence between the elements of two sets.

### Example:

If  $A = \{2, 3, 5\}$  and  $B = \{a, e, i, o, u\}$ , which of the following relations from A to B are functions?

Also tell the type of function.

- (i)  $R_1 = \{(2, i), (2, o), (5, u), (3, e)\}$       (ii)  $R_2 = \{(3, i), (2, o), (5, i)\}$   
(iii)  $R_3 = \{(2, u), (5, i), (3, o)\}$

Is it possible to have a function from A onto B?

### Solution:

- (i)  $R_1 = \{(2, i), (2, o), (5, u), (3, e)\}$

$\text{Dom}(R_1) = \{2, 3, 5\} = A$ , but there is repetition in the first element of first two ordered pairs, therefore  $R_1$  is not a function.

(II)  $R_2 = \{(3, i), (2, o), (5, i)\}$

$\text{Dom}(R_2) = \{2, 3, 5\} = A$  and there is no repetition in the first element of the ordered pairs in  $R_2$ , therefore  $R_2$  is a function.  $\text{Range}(R_2) = \{i, o\} \subset B$  which implies that  $R_2$  is an into function.

(III)  $R_3 = \{(2, u), (5, i), (3, o)\}$

$\text{Dom}(R_3) = \{2, 3, 5\} = A$  and there is no repetition in the first element of the ordered pairs in  $R_3$ , therefore  $R_3$  is a function.

$\text{Range}(R_3) = \{i, o, u\} \subset B$  which implies that  $R_3$  is an into function. Also, there is no repetition in the second element of the ordered pairs in  $R_3$ , so  $R_3$  is one-one function.

Hence,  $R_3$  is an injective function.

Also number of elements in set  $B$  is more than number of elements in set  $A$ , therefore  $A$  onto  $B$  function is not possible.

## Evaluating a Function

A function is usually written as  $y = f(x)$ , in which the values of  $y$  (dependent variable) depends upon the values of  $x$  (independent variable) and is read as 'y is a function of  $x$ ' or simply  $y$  is equal to  $f$  of  $x$ .

Evaluation of a function is a process of finding the value of dependent variable by substituting any specific value of the independent variable.

**Example:** Find the value of  $f(x) = -2x^3 + x^2$  at:

(I)  $f(-1)$

(II)  $f(t)$

(III)  $f\left(\frac{-1}{2}\right)$

(IV)  $f(x+2)$

**Solution:** (I) Substituting  $x = -1$  in  $f(x) = -2x^3 + x^2$ , we have

$$f(-1) = -2(-1)^3 + (-1)^2 = -2(-1) + 1 = 3$$

(II) Replacing  $x$  by  $t$  in  $f(x)$ , we have

$$f(t) = -2(t)^3 + (t)^2 = -2t^3 + t^2$$

(III) Substituting  $x = \frac{-1}{2}$  in  $f(x)$ , we have

$$f\left(\frac{-1}{2}\right) = -2\left(\frac{-1}{2}\right)^3 + \left(\frac{-1}{2}\right)^2 = -2\left(\frac{-1}{8}\right) + \frac{1}{4} = \frac{1}{2}$$

(IV) Replacing  $x$  by  $x+2$  in  $f(x)$ , we have

$$\begin{aligned}f(x+2) &= -2(x+2)^3 + (x+2)^2 = -2x^3 - 16 - 12x^2 - 24x + x^2 + 4x + 4 \\&= -2(x^3 + 8 + 6x^2 + 12x) + (x^2 + 4x + 4) \\&= -2x^3 - 11x^2 - 20x - 12\end{aligned}$$

**Example:** The value of a photocopy machine  $t$  years after its purchase is given by

$$V(t) = 9500 - 860t \text{ dollars.}, \text{Find:}$$

- (i)  $V(5)$  and tell what does  $V(5)$  mean?
- (ii) Find  $t$  when  $V(t) = 6920$  and explain what it represents.

**Solution:**

(i)  $V(t) = 9500 - 860t$

$$V(5) = 9500 - 860 \times 5 = 5200 \text{ dollars}$$


It means the value of machine after 5 years will be 5200 dollars.

(ii) When  $V(t) = 6920$ , then  $6920 = 9500 - 860t$

$$\Rightarrow 860t = 9500 - 6920 = 2580$$

$$\Rightarrow t = \frac{2580}{860} = 3$$

Hence,  $t = 3$ , which tells us that the value of photocopy machine will be 6920 dollars after 3 years.






### Exercise 6.1

1. If  $A = \{2, 4, 6, 8\}$  and  $B = \{1, 3, 5\}$ , which of the following relations are functions from  $A$  to  $B$ ?

- (i)  $R_1 = \{(2, 3), (6, 5), (8, 3), (4, 1)\}$
- (ii)  $R_2 = \{(2, 3), (6, 1), (8, 3), (6, 5)\}$
- (iii)  $R_3 = \{(2, 3), (6, 5), (8, 3)\}$
- (iv)  $R_4 = \{(2, 3), (6, 3), (8, 3), (4, 1)\}$

2. Identify the functions from the following diagrams. Also write their type.



3. If  $f(x) = x^2 - \frac{1}{2}x + 3$ , evaluate the following.

(i)  $f(2)$       (ii)  $f(-1)$       (iii)  $f(\frac{2}{3})$       (iv)  $f(t + 1)$

4. If  $X = \text{Set of prime factors of } 6$  and  $Y = \text{Set of first three non-negative integers}$ , check whether the function defined by  $f : X \rightarrow Y = \{(x, y) | x - y = 1\}$  is injective or bijective.

- A function  $f$  is defined by  $f(x) = t - ax$ , find the values of  $a$  and  $t$  if 0 and 2 are the images of 1 and  $-1$  respectively.
- A function  $f$  is defined by  $f(x) = mx + c$ , is  $f$  bijective? Justify your answer.

## Operations on Functions

Operations on functions are the ways of combining functions to create new functions. Some common operations on any two functions  $f(x)$  and  $g(x)$  are:

- (i) **Addition of functions:**  $(f + g)(x) = f(x) + g(x)$ .

For example, if  $f(x) = x^2$  and  $g(x) = 2x$ , then

$$(f + g)(x) = f(x) + g(x) = x^2 + 2x$$

- (ii) **Subtraction of functions:**  $(f - g)(x) = f(x) - g(x)$ .

For example, if  $f(x) = 2x^2 + 3x$  and  $g(x) = x^2 - 2x$ , then,

$$\begin{aligned} (f - g)(x) &= f(x) - g(x) = (2x^2 + 3x) - (x^2 - 2x) \\ &= 2x^2 + 3x - x^2 + 2x = x^2 + 5x \end{aligned}$$

- (iii) **Multiplication of functions:**  $(f \times g)(x) = f(x) \times g(x)$ .

For example, if  $f(x) = x^2$  and  $g(x) = -2x$ , then  $(f \times g)(x) = f(x) \times g(x)$   
 $= x^2 \times (-2x) = -2x^3$

- (iv) **Division of functions:**  $(f \div g)(x) = f(x) \div g(x)$ , where  $g(x) \neq 0$

For example, if  $f(x) = x^2$  and  $g(x) = 2x$ , then  $(f \div g)(x) = f(x) \div g(x)$   
 $= x^2 \div (2x) = \frac{x^2}{2x} = \frac{x}{2}$

- (v) **Composition of functions:** The composition of  $f(x)$  and  $g(x)$  is a function denoted by  $fog$  and is defined as:

$$(fog)(x) = f(g(x))$$

For example, if  $f(x) = x^2$  and  $g(x) = 2x - 1$ , then:

$$(fog)(x) = f(g(x)) = f(2x - 1) = (2x - 1)^2 = 4x^2 - 4x + 1$$

In other words, the output of inner function  $g(x)$  will become the input of the outer function  $f(x)$ .

In this example, we first applied the function  $g$  to  $x$ , which resulted in  $2x - 1$ , then applied the function  $f$  to the result which gave us  $(2x - 1)^2$ .

Furthermore, for finding  $(gof)(x)$ ,  $f(x)$  will be the input of the outer function  $g(x)$ .

So,  $(gof)(x) = g(f(x)) = g(x^2) = 2x^2 - 1$

### Key Fact:

- $(fog)(x)$  and  $(gof)(x)$  can simply be written as  $fg(x)$  and  $gf(x)$  respectively.
- $f(f(x)) = f^2(x)$  and  $f(f^2(x)) = f^3(x)$

**Example:** Aahil wants to have a new bike which costs 24000 rupees but he has only a saving of 12000 rupees. If his parents have agreed to give him additional 1000 rupees per week for doing chores. However, he also spends 500 rupees per week on snacks. How much will Aahil have as his

saving after 10 weeks. Also, tell in how many weeks he will be able to save the money for getting a new bike?

**Solution:** Aahil's saving function:  $S(x) = 12000 + 1000x$ , where  $x$  represents number of weeks  
His spending function:  $T(x) = 500x$

Total money with Aahil after  $x$  weeks =  $S(x) - T(x)$

$$= 12000 + 1000x - 500x = 12000 + 500x$$

After 10 weeks, saving with Aahil =  $12000 + 500(10) = 17000$  rupees


$$\text{Now, } 24000 = 12000 + 500x \Rightarrow 24000 - 12000 = 500x \Rightarrow x = \frac{12000}{500} = 24$$

Hence, he can save required amount in 24 weeks.

## Inverse of a Function

The inverse of any function  $f(x)$  is a function denoted by  $f^{-1}(x)$  which reverses the effect of  $f(x)$  and it undoes what  $f(x)$  does. If  $f: X \rightarrow Y$ , then  $f^{-1}: Y \rightarrow X$ , therefore inverse of only that function exists which is bijective (both one-one and onto).

Thus, if  $y = f(x) \forall x \in X$ , then  $x = f^{-1}(y) \forall y \in Y$ . It can be illustrated in the following diagram as :



It is clear from above diagram that domain of  $f$  = range of  $f^{-1}$  and range of  $f$  = domain of  $f^{-1}$ .

### Finding the Inverse of an Algebraic Function

If  $f(x)$  is any algebraic function, then its inverse  $f^{-1}(x)$  involves the following steps:

- I. Write  $y = f(x)$
- II. Find the value of  $x$  in terms of  $y$  from the equation of step I, means write  $x$  in terms of  $y$ .
- III. Replace  $x$  by  $f^{-1}(y)$  in the resulting equation of step II.
- IV. Replace each  $y$  by  $x$  in the resulting equation of step III to get  $f^{-1}(x)$ .

The result obtained in step IV can be verified by showing  $f^{-1}(f(x)) = x$ .

**Example:** If  $f: R \rightarrow R$  is defined by  $f(x) = 3x + \frac{1}{2}$ , find  $f^{-1}(x)$ .

**Solution:** Writing  $f(x) = y$ , we have,

$$y = 3x + \frac{1}{2}$$

$$3x = y - \frac{1}{2} = \frac{2y-1}{2} \Rightarrow x = \frac{2y-1}{6} \quad (\text{Isolating } x)$$

$$\text{So, } f^{-1}(y) = \frac{2y-1}{6}$$

$$\text{Hence, } f^{-1}(x) = \frac{2x-1}{6} \quad (\text{Replacing each } y \text{ by } x)$$

### Exercise 6.2

1. If  $f(x) = 4(x - 1)$  and  $g(x) = x^2 - 2x + 1$ , find  
(i)  $(f+g)(x)$     (ii)  $(f-g)(x)$     (iii)  $(f \times g)(x)$     (iv)  $(f \div g)(x)$
2. If  $f(x) = 4x$  and  $g(x) = x + 1$ , find,  
(i)  $(fog)(x)$     (ii)  $(gof)(x)$     (iii)  $(fof)(x)$     (iv)  $(gog)(x)$
3. Find  $(fog)(x)$  and  $(gof)(x)$ , if  
(i)  $f(x) = 3 - 2x$  and  $g(x) = x + 1$   
(ii)  $f(x) = \frac{2}{x}$  and  $g(x) = \frac{2x}{x-1}$   
(iii)  $f(x) = 3x$  and  $g(x) = \frac{2}{\sqrt{x-1}}$   
(iv)  $f(x) = x^2 - 1$  and  $g(x) = \sqrt{x-1}$
4. If  $f(x) = x^2$  and  $g(x) = 2x + 1$ , find the value of  $x$  if:  
(i)  $(fog)(x) = (gof)(x)$     (ii)  $f(x) = g(x)$     (iii)  $(gof)(x) = 9$
5. Find the inverse of the following functions.  
(i)  $f(x) = 2x - 1$     (ii)  $g(x) = \frac{2}{x-3}$ , where  $x \neq 3$   
(iii)  $f(x) = \sqrt{x+5}$ , where  $x \geq -5$     (iv)  $g(x) = (x-3)^2$ , where  $x \geq 3$
6. If  $f(x) = \frac{3}{x-5}$  and  $g(x) = x + \frac{3}{2}$ , find the value of  $f^{-1}(1)$  and  $g^{-1}\left(\frac{1}{2}\right)$ .

## Graphs of Some Algebraic Functions

### Polynomial Functions

A function  $f : R \rightarrow R$  defined as  $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots + a_{n-1}x^{n-1} + a_nx^n$ , where  $a_0, a_1, a_2, a_3, \dots, a_{n-1}, a_n$  are all real numbers and  $n$  (called the degree of polynomial) is a non-negative integer is called a polynomial function.

For example,  $f(x) = 2x^3 - x^2 + x$  is a polynomial function of degree 3 having leading coefficient 2.

#### Key Fact:

- The highest exponent of the variable involved in a polynomial is called its degree.
- The coefficient of highest degree term in a polynomial is called the leading coefficient.


### (i) Zero Polynomial Function

A polynomial function of no degree is called a zero polynomial function. This function is of the form:

$y = f(x) = 0$  in which  $a_0, a_1, a_2, a_3, \dots, a_{n-1}, a_n$  are all zero.

The graph of  $y = 0$  is a straight line that lies on the x-axis indicating that the value of  $y$  is always zero regardless of the value of  $x$ . It passes through origin and extends infinitely in both positive and negative x-directions.

The graph of  $y = 0$  is shown in the given coordinate plane.



### Do You Know?

A vertical line intersects the graph of a function only at one point. If it intersects the graph at more than a point, then that graph will not be the graph of a function. This is called vertical line test.

### (ii) Constant Function


A polynomial function of degree zero is called a constant function.

This function is of the form,

$$y = f(x) = a = ax^0,$$

where  $a$  is any non-zero constant.

For example,  $y = 3$ ,  $y = \frac{-1}{2}$  etc.



The graph of  $y = 1$  (shown in the given graph) is a straight line parallel to x-axis.

### (iii) Linear Function

A polynomial function of degree one is called a linear function. This function is of the form,  $y = f(x) = ax + b$ , where  $a, b \in R$  and  $a \neq 0$ . For example,  $y = 2x + 3$ ,  $y = 2x$  etc.

The graph of a linear equation is a straight line. For sketching the graph of a linear function, we first find two points which satisfy the given equation. It is convenient to find x-intercept and y-intercept for getting two points.


To calculate x-intercept, substitute  $y = 0$  in  $y = 2x + 3$  and to calculate y-intercept, substitute  $x = 0$  in the equation.

**x-intercept:** Put  $y = 0$ ,  $0 = 2x + 3$

$$\Rightarrow x = -\frac{3}{2} = -1.5$$

So, the point where the graph intersects x-axis is  $(-1.5, 0)$ .

**y-intercept:** Put  $x = 0$ ,  $y = 3$



The point where the graph intersects y-axis is (0, 3).

From the graph, it is clear that this line does not pass through origin. We can find the coordinates of any point on the graph just by taking its distance from the axis.


#### (iv) Identity Function

An identity function can be represented as:

$y = f(x) = x$ , means it has the same value as the input.


For example,  $f(2) = 2$ ,  $f(-3) = -3$  etc.

It is a linear function and its graph is a straight line bisecting first and third quadrants passing through origin.



#### (v) Absolute Valued Function

The absolute value of a real number is its distance from 0 on the number line.



For example, the absolute value of 5 written as  $|5| = 5$  and of  $-5$  written as  $|-5| = 5$ .



The absolute value of a real number  $x$  denoted by  $|x|$  is defined as:


$$|x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$$

The function defined by  $f(x) = |x|$ , is called the absolute valued function or the modulus function.

Its graph is V-shaped with vertex at origin.

It bisects first quadrant for  $x > 0$  and bisects second quadrant for  $x < 0$ .

Its domain is set of all real numbers but range is set of non-negative real numbers.



Absolute valued functions are widely used in many real life problems including distance, displacement, magnitude and to calculate energy wave etc.

#### Example:

The profit (in rupees) of a company is given by the function  $P(x) = |2x - 100|$ , where  $x$  is the number of units sold. If 120 units are sold, what is the profit?

**Solution:**

Profit function,  $P(x) = |2x - 100|$

Substituting  $x = 120$ ,  $P(120) = |2 \times 120 - 100| = 140$

Thus, the company makes a profit of 140 rupees.

**Example:**

A wave's energy ( $E$ ) is given by the function  $E(x) = |3\sin(x)|$ , where  $x$  is the wave's phase angle in radians. If the phase angle is  $30^\circ$ , what is the wave's energy?

**Solution:**

First convert phase angle to radians as under:

$$\text{Phase angle} = 30^\circ \times \frac{\pi}{180^\circ} \text{ radians} = \frac{\pi}{6} \text{ radians}$$

Now substitute  $x = \frac{\pi}{6}$  in the given function,


$$E\left(\frac{\pi}{6}\right) = \left|3\sin\left(\frac{\pi}{6}\right)\right| = \left|3 \times \frac{1}{2}\right| = \frac{3}{2}$$

Hence, the wave's energy is  $\frac{3}{2}$  units.

**(vi) Graphs of the functions of the form  $y = nx$ , where  $n$  is an Integer.**

For the graph of the function  $y = nx$ ;  
 $n$  is a + ve or - ve integer and  $x > 0$ ,  
we take different integral values of  $n$ .


The graph for  $n = -2, -1, 1, 4$  is drawn  
at right.

**Points To Ponder!**

From the graph, we observe that:

- All these lines passes through origin.
- For positive value of  $n$ , the line passes through first and third quadrant.
- For negative value of  $n$ , the line passes through second and fourth quadrant.

Similarly, we can sketch the linear graphs for fractional values of  $n$  as drawn here.




## Identifying, Sketching and Interpreting Graphs of Some Non-Linear Functions

### (i) Quadratic Function

A polynomial function of degree two is called a quadratic function. This function is of the form,  $f(x) = ax^2 + bx + c$ ;  $a, b, c \in R$  and  $a \neq 0$ .

For example,  $y = x^2 - 2x - 3$ ,  $y = -4x^2 + 1$  are quadratic functions.

The graph of a quadratic function is a U-shaped curve called parabola that opens either upward (if  $a > 0$ ) or downward (if  $a < 0$ ).



The vertical line passing through the vertex of parabola is called 'axis of symmetry' as the graph is symmetrical about y-axis.

In order to sketch the graph of such quadratic functions, follow the following steps.

**Step-1:** Identify the shape of the graph whether it opens upward or downward.

**Step-2:** Find x-intercept and y-intercept.

**Step-3:** Find the vertex of the parabola.

For finding vertex of the parabola, we proceed as follows:

$$\begin{aligned} y &= ax^2 + bx + c \\ &= a(x^2 + \frac{b}{a}x) + c \end{aligned}$$

(Isolate the constant term and take coefficient of  $x^2$  as common from first two terms)

$$= a \left( x^2 + \frac{b}{a}x + \left(\frac{b}{2a}\right)^2 - \left(\frac{b}{2a}\right)^2 \right) + c$$

(Adding and subtracting square of half of the coefficient of  $x$  to complete the square)

$$= a \left( x + \frac{b}{2a} \right)^2 - a \left( \frac{b}{2a} \right)^2 + c$$

$$= a \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a} + c$$

$$= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a} \quad \text{-----} \rightarrow (\text{i})$$

Comparing (i) with

$$y = a(x - h)^2 + k \quad \text{-----} \rightarrow (\text{ii}) \text{ (general form of the curve)}$$

We have,

$$h = \frac{-b}{2a} \text{ and } k = c - \frac{b^2}{4a}$$

So, the graph of the curve,  $y = ax^2 + bx + c$  is a parabola with vertex at  $(h, k)$ . The point  $(h, k)$  is the turning point of the graph. The curve is symmetrical about the line  $x = h$ .

To sketch the graph of parabola, we draw its vertex, x- and y-intercepts and some other points to get a smooth curve.

### Example:

Sketch the graph of  $y = x^2 - 2x - 3$ .

### Solution:

$$y = x^2 - 2x - 3 \quad \text{-----} \rightarrow (\text{i})$$

Comparing (i) with  $y = ax^2 + bx + c$ , we have  $a = 1, b = -2$  and  $c = -3$

Here,  $a = 1 > 0$ , so the curve will open upward.

$$h = \frac{-b}{2a} = \frac{-(-2)}{2(1)} = 1 \text{ and}$$

$$k = c - \frac{b^2}{4a} = -3 - \frac{(-2)^2}{4(1)} = -4$$

So, the vertex of parabola is  $(h, k) = (1, -4)$ .

For y-intercept, put  $x = 0, y = -3$

For x-intercept, put  $y = 0$ .

$$0 = x^2 - 2x - 3$$

$$0 = x^2 - 3x + x - 3$$

$$0 = x(x - 3) + 1(x - 3)$$

$$0 = (x - 3)(x + 1)$$


This implies, either  $x - 3 = 0$  or  $x + 1 = 0$

$$\Rightarrow x = 3 \text{ or } x = -1$$

So, the graph will intersect x-axis at  $(3, 0)$  and  $(-1, 0)$ . The parabola is symmetrical about the vertical line  $x = -1$  through the turning point.

### Enlighten Yourself!

If the graph of a function intersects x-axis at point  $(a, 0)$ , then x-coordinate 'a' of the point is called x-intercept and y-coordinate of that point will always be zero. Similarly y-coordinate 'b' of  $(0, b)$  is called y-intercept in which x-coordinate is always zero.



## (ii) Cubic Function

From bottom left to top right


A polynomial function of degree three is called a cubic function. This function is of the form:

$$f(x) = ax^3 + bx^2 + cx + d ; a, b, c, d \in R \text{ and } a \neq 0.$$

We will discuss here the graph of the functions  $y = ax^3$ .

To sketch the graph of  $y = x^3$  (when  $a = 1$ ), we need the shape,  $x$  and  $y$  intercepts and turning point.

Here leading coefficient is 1 which is positive which shows that the graph will concave up in the first quadrant and concave down in the third quadrant. For each point  $(x, y)$  on the graph, the point  $(-x, -y)$  is also on the graph. At  $x = 0, y = 0$ , so this graph intersects both the axis at origin and is symmetric about origin as shown in the figure. Also origin is the turning point of this graph. The domain and range of this function is the set of all real numbers.



From top left to bottom right


### Example:

Sketch the graph of  $y = -2x^3$ .

**Solution:**  $y = -2x^3$  -----> (i)

Here, the leading coefficient is  $-2$  (negative), which shows that the graph will concave up in the second quadrant and concave down in the fourth quadrant.

At  $x = 0, y = 0$ , so the graph passes through origin which is the turning point of the graph. For each point  $(-x, y)$  on the graph, the point  $(x, -y)$  is also on the graph, so this graph is symmetric about origin. The domain and range of this function is the set of all real numbers.




### (iii) Square Root Function

The function defined by  $f(x) = \sqrt{x}$ , where  $x \geq 0$  is called a square root function.

The graph of this function will remain in first quadrant and increases with increase in the value of  $x$  as shown in the given graph.

The domain and range of this function is the set of non-negative real numbers.



### (iv) Reciprocal Function

The reciprocal function is of the form  $f(x) = \frac{k}{x}$ ,

where  $x \neq 0$  and  $k$  (a constant)  $\neq 0$ . We will discuss here simple example of reciprocal function  $f(x) = \frac{1}{x}$ . The graph of reciprocal function shows how the function's output decreases as the input increases illustrating inverse relationships.

#### Enlighten Yourself!

- The reciprocal of a very large number is a very small number.
- The reciprocal of a very small number is a very large number.
- The reciprocal of zero is undefined.

First we draw the graph of the original function  $y = x$  whose reciprocal is to be drawn.


The graph of  $y = x$  is a straight line passing through origin bisecting first and third quadrants.

The graph of  $y = \frac{1}{x}$  is a curve which is undefined at  $x = 0$  and is symmetrical about the line  $y = x$ . As we increase the value of  $x$ , the value of the function decreases and vice versa.

This graph never touches coordinate axis.

$y \rightarrow \infty$  as  $x \rightarrow 0$ , but it will never touch y-axis and  $x \rightarrow \infty$  as  $y \rightarrow 0$ , but it will never touch x-axis.

The domain of this curve is all real numbers except zero.



### Have You Noticed?

- The graph of  $f(x) = \frac{1}{x}$  has two branches.
- The graph of  $y = \frac{1}{x}$  exists in the first and third quadrants only.
- As  $x \rightarrow \infty$ ,  $f(x) \rightarrow 0$  from above.  
As  $x \rightarrow -\infty$ ,  $f(x) \rightarrow 0$  from below.  
As  $x \rightarrow 0$  from the right,  $f(x) \rightarrow \infty$ .  
As  $x \rightarrow 0$  from the left,  $f(x) \rightarrow -\infty$ .

### Enlighten Yourself!

The symbol  $\infty$  was first used by John Wallis in 1655 which means unbouded.

### Example:


Sketch the graph of  $y = \frac{1}{x-2}$ .

### Solution

$$\text{y-intercept: } y = \frac{-1}{2} = -0.5$$

As  $x \rightarrow \infty$ ,  $y \rightarrow 0$

As  $x \rightarrow -\infty$ ,  $y \rightarrow 0$



### Example:

Saim can jog 4 miles downhill in the same time which he takes to jog 2 miles uphill. He takes downhill 3mph faster than he jogs uphill. Find his jogging rate uphill and downhill.

### Solution:

Let his uphill rate =  $x$

Then, his downhill rate =  $x + 3$


$$\text{Uphill time} = \frac{2}{x}$$

$$\text{Downhill time} = \frac{4}{x+3}$$

According to the given condition,  
downhill time = uphill time

$$\begin{aligned}\Rightarrow \frac{4}{x+3} &= \frac{2}{x} \\ \Rightarrow 4x &= 2(x+3) \\ \Rightarrow 4x &= 2x + 6 \Rightarrow 4x - 2x = 6 \Rightarrow x = 3\end{aligned}$$

Thus, his uphill rate = 3mph and his downhill rate =  $3 + 3 = 6$  mph



### (v) Exponential Function

The most simple exponential function is of the form  $f(x) = a^x$ , where  $a > 0$  and  $a \neq 1$ .

For example,  $y = 2^x$  is an exponential function.

For sketching its graph, we find:

**y-intercept:**  $y = 1$

**Asymptote:** Horizontal asymptote,  $y = 0$

As  $x$  becomes large and negative, the graph of  $y = 2^x$  becomes closer to  $x$ -axis but never quite reaches it.


We can say that as  $x$  approaches  $-\infty$ ,  $y$  approaches 0.

Similarly, as  $x$  becomes large and positive,  $y$  also

becomes large and positive, so we can say that as  $x$  approaches  $\infty$ ,  $y$  also approaches  $\infty$ . The simple exponential functions have smooth increasing or decreasing graphs. All exponential graphs are similar in shape and have a horizontal asymptote.


The exponential functions are used for the quantities which increase or decrease exponentially.

For example, population of animals, people and bacteria usually grow in an exponential way.



### (vi) The Function $y = x^n$ , where $n$ is +ve or -ve Integer and $x > 0$

For the graph of the function  $y = x^n$ , where  $n$  is a + ve or - ve integer and  $x > 0$ , we take different integral values of  $n$ . The graphs for  $n = -2, -1, 0, 1, 2$  are shown in the given graph.



#### Enlighten Yourself!

The graph of  $y = x^n$  (where  $n$  is +ve or -ve integer and  $x > 0$ ):

- Is a horizontal line at  $n = 0$ .
- Bisects first quadrant at  $n = 1$ .
- Will remain in first quadrant.
- Passes through  $(1, 1)$ .

## Solution of a Linear and a Quadratic Equation Graphically

As we know that the graph of a linear equation is a straight line and of a quadratic equation is a curve. The point of intersection of these two graphs is a point where both the graphs intersect each other.

**Example:**

Solve  $f(x) = 2x + 1$  and  $g(x) = x^2 - x + 3$  graphically.

**Solution:**

Table of values for  $f(x) = 2x + 1$

$x$	-2	-1	0	1	3
$f(x)$	-3	-1	1	3	7

For the graph of  $g(x) = x^2 - x + 3$ , we compare it with  $y = ax^2 + bx + c$ , we have:

$$a = 1, b = -1 \text{ and } c = 3.$$

Here,  $a = 1 > 0$ , so the curve will open upward.

$$\text{Also } h = \frac{-b}{2a} = \frac{-(-1)}{2(1)} = \frac{1}{2} = 0.5$$

$$\text{and } k = c - \frac{b^2}{4a} = 3 - \frac{(-1)^2}{4(1)} = 2.75$$

So, the vertex of parabola is:

$$(h, k) = (0.5, 2.75).$$


Table of values for  $g(x) = x^2 - x + 3$

$x$	-1	0	0.5	1	2	3
$g(x)$	5	3	2.75	3	7	9

Both the graphs intersect each other at:

$$(1, 3) \text{ and } (2, 5).$$

Hence, solution set =  $\{(1, 3) \text{ and } (2, 5)\}$



## Determining the Gradient of Curves through Tangents

The gradient of a curve at a given point is the slope of the tangent line drawn to the curve at that point. It is obtained by drawing a tangent to the curve at that point. The tangent to a curve is a line which touches the curve only at one point externally. The slope of tangent to the curve is its gradient. The slope of the tangent =  $\frac{\text{change in } y}{\text{change in } x}$  ( $\frac{\text{rise}}{\text{run}}$ ).

So, the gradient of the curve =  $\frac{\text{change in } y}{\text{change in } x} = \frac{y_2 - y_1}{x_2 - x_1}$


### **Example:**

Find the gradient to the curve  $y = x^2$  drawn below at:

- (i)  $x = 1$       (ii)  $0$       (iii)  $-1.4$ .

**Caution!**  
Make sure that the angle of tangent line with the curve below and above the given point must be same. The gradient of the curve can also be found by taking any two points on the tangent line.

**Solution:** The graph of the given curve is drawn here.



- (i) For finding gradient, we first draw a tangent to the curve with equal space on either side at  $x = 1$ . Then draw right triangle by taking the point at which  $x = 1$  and any other point on the tangent line as shown in the graph.

The gradient of the tangent =  $\frac{y_2 - y_1}{x_2 - x_1} = \frac{\text{change in } y}{\text{change in } x} = \frac{3-1}{2-1} = 2$

Hence, the gradient of the curve at  $x = 1$  is 2.

- (ii) The tangent at  $x = 0$  is a horizontal line, so its slope is zero. Hence, the gradient of the curve at  $x = 0$  is 0.
- (iii) Similarly by drawing a tangent to the curve at  $x = -1.4$ , we have a right triangle as shown in the graph. By taking the point  $(-1.4, 2)$  and any other point  $(-0.8, 0.4)$  on the tangent line, we have,

$$\begin{aligned}\text{The gradient of the tangent} &= \frac{y_2 - y_1}{x_2 - x_1} \\ &= \frac{0.4 - 2}{-0.8 + 1.4} = \frac{-1.6}{0.6} \approx -2.7\end{aligned}$$

#### Key Fact:

The slope of the vertical line is undefined.

#### Yo Be Noted!


- If the tangent line comes down from left to right, the slope of tangent line will be negative.
- If the tangent line goes up from left to right, the slope of tangent line will be positive.

#### Example:

Estimate gradient to the given curve at  $(1, 2)$ .

**Solution:** We first draw a tangent to the curve with equal space on either side at  $(1, 2)$ . Then draw right triangle at the point at  $(1, 2)$  and any other point on the tangent line as shown in the graph.

$$\begin{aligned}\text{The gradient of the tangent} &= \frac{y_2 - y_1}{x_2 - x_1} \\ &= \frac{\text{change in } y}{\text{change in } x} \\ &= \frac{4 - 2}{2.5 - 1} = 1.33\end{aligned}$$



Hence, the gradient of the curve at  $(1, 2)$  is estimated as 1.33.

#### Real Life Problems Related to Functions and Graphs

Graphs and functions are essential tools in various aspects of daily life. They have wide applications often unconsciously in solving many real life problems including distance and time, supply and demand, temperature and time, population growth, health and fitness, weather forecasting, financial planning, time management, medical research, environmental monitoring, fitness tracking etc. By applying graphs and functions to daily life problems, we can develop a more analytical approach to decision making and problem solving.

Here are some problems involving functions and their graphs.


**Example:** A company's cost function is  $C(x) = 2x + 300$ , where  $x$  is the number of units produced and cost is given in dollars. Graph the cost function and find the cost of producing 150 units.

**Solution:**

Here,  $C(x) = 2x + 300$

Table of values

$x$	0	50	100	150
$C(x)$	300	400	500	600



From the graph, cost of producing 150 units is 600 dollars.

**Example:** A researcher studies the spread of a disease and finds that the number of infected people grows according to the function  $f(t) = 10(1.2)^t$ , where  $t$  is time in days. Determine the number of people to become infected in next 6 days.


**Solution:**

Here,  $f(t) = 20(1.2)^t$






Table of values

$t$	0	2	4	6
$f(t)$	20	29	41	60

From the graph, number of people to be infected in next 6 days = 60



1. Identify the graphs of linear and non-linear functions from the following? Also indicate the graphs which are graphs of functions by applying vertical line test.



2. Sketch the graph of the following functions and identify the graphs which are linear.

(i)  $y = 3x + 2$       (ii)  $y = x$       (iii)  $y = x^2$

3. Sketch the graph of the following functions.

(i)  $y = 3^x$       (ii)  $y = |x|$       (iii)  $y = x^3$   
 (iv)  $y = \sqrt{x+2}$       (v)  $y = \frac{1}{x-1}$

4. Check whether the curve (parabola) drawn for the following functions will open upwards or downwards. Also find x-intercept, y-intercept and sketch the graph of the given functions.

(i)  $f(x) = (x - 1)^2$       (ii)  $f(x) = -x^2 + 2$   
 (iii)  $f(x) = 3 - (x + 2)^2$       (iv)  $f(x) = -x^2 + 2x + 3$

5. Find the values of h and k if  $y = hx^2 + 2x + k$  cuts x-axis and y-axis at  $(-3, 0)$  and  $(0, 2)$  respectively.

6. Solve the following equations graphically.

(i)  $f(x) = x + 2$ ,  $g(x) = x^2 - 3x + 2$       (ii)  $f(x) = 2x + 5$ ,  $g(x) = 2x^2 + 1$

7. If demand function of a product is  $D(x) = 100 - 5x$  and supply function is  $S(x) = x - 200$ . Draw graphs of both functions on same graph by using same scale and find the value of  $x$  for which the supply will become equal to demand?

8. The growth of the population of a town is given by the function,  $P(t) = 50,000(1.05)^t$ , where  $t$  is the time in years. Draw the graph of this function and find the population growth over 5 years.

9. Draw gradient of the curve  $y = \frac{1}{2}x^2$  by drawing tangent line at  $x = 2$ .

10. A logistics company charges a base fee of 50 dollars and additional charges of 0.5 dollars per mile for transporting goods. If a shipment travels 250 miles, what is the total cost?

### I have Learnt

- Cartesian product of any two non-empty sets A and B is a set containing all ordered pairs  $(x, y)$  such that  $x \in A$  and  $y \in B$ .
  - A function  $f$  from set A to set B is a relation, rule or mapping which maps each element of set A to a unique element of set B.
    - A function  $f : A \rightarrow B$  will be into if  $\text{range}(f) \subset B$ .
    - A function  $f : A \rightarrow B$  will be onto if  $\text{range}(f) = B$ .
    - A function  $f : A \rightarrow B$  will be one-one if unique element of A have unique image in B.
    - A function  $f : A \rightarrow B$  will be injective if it is both into and one-one.
    - A function  $f : A \rightarrow B$  will be bijective if it is both onto and one-one.
  - Every bijective function has one-one correspondence between its elements.
  - The process of finding the value of dependent variable by substituting any specific value of the independent variable is called evaluation of a function.
  - For any two functions  $f$  and  $g$ :
    - $(f + g)(x) = f(x) + g(x)$ ,  $(f - g)(x) = f(x) - g(x)$
    - $(f \times g)(x) = f(x) \times g(x)$ ,  $(f \div g)(x) = f(x) \div g(x)$
  - The composition of any two functions,  $f(x)$  and  $g(x)$  is denoted by  $(f \circ g)(x) = f(g(x))$ .
  - If  $y = f(x) \forall x \in X$ , then  $x = f^{-1}(y) \forall y \in Y$ .
  - The domain of  $f(x) = \text{range of } f^{-1}(x)$  and range of  $f(x) = \text{domain of } f^{-1}(x)$ .
  - A polynomial function of no degree is called a zero polynomial function.
  - A polynomial function of degree zero is called a constant function.
  - A polynomial function of degree one is called a linear function.
  - A polynomial function of degree two is called a quadratic function.
  - The graph of the curve,  $y = ax^2 + bx + c$  is a parabola with vertex at  $(h, k)$ , where  $h = \frac{-b}{2a}$  and  $k = c - \frac{b^2}{4a}$ . It will open upward when  $a > 0$  and open downward when  $a < 0$ .
  - A polynomial function of degree three is called a cubic function.
  - The function of the form,  $f(x) = \sqrt{x}$ , where  $x \geq 0$  is called a square root function.
  - The function of the form,  $f(x) = |x|$  is called the absolute valued function.
  - The function defined by  $f(x) = \frac{1}{x}$ , where  $x \neq 0$  is called a reciprocal function.
  - The exponential function is of the form  $f(x) = a^x$ , where  $a > 0$  and  $a \neq 1$ .
  - The point of intersection of two graphs is called their solution point.

## MISCELLANEOUS EXERCISE-6

1. Encircle the correct option in each of the following.

i. If  $A = \{-2, 0, 2\}$  and  $B = \{0, 2\}$  and  $f: A \rightarrow B$  is defined as  $f = \{(-2, 2), (0, 0), (2, 0)\}$   
 what type of function is  $f$ ?

(a) into      (b) onto      (c) injective      (d) bijective



# 07

## Vectors in Plane

After studying this unit students will be able to:

- Introduce rectangular coordinate system in a plane.
- Represent vector as a directed line segment.
- Express a vector in terms of two non-zero and non-parallel coplanar vector.
- Express a vector in terms of position vector.
- Express translation by a vector.
- Find the magnitude of a vector.
- Add and subtract vectors.
- Multiply a vector by a scalar.
- Solve geometrical problems involving the use of a vector.
- Apply concepts of vectors in geometrical problems such as parallel and perpendicular lines in geometrical shapes, vector projectile motion, crosswinds aviation, military usage, designing roller coasters.

Vectors are fundamental mathematical concept that have a wide range of applications in the various fields from the physics and engineering to the computer graphics and navigation systems. Vectors are utilized to assist in the localization of people, places and things, and to describe things that are acting in response to an external force being applied to them.

### Do you know?

- Navigating by air and by boat is generally done using vectors.
- Planes are given a vector to travel, and they use their speed to determine how far they need to go before turning or landing. Flight plans are made using a series of vectors.
- Sports instructions are based on using vectors.
- Vectors are used in electrical engineering for analyzing and designing circuits, signals, and electromagnetic systems.



# Introduction of Vectors

There are two main types of physical quantities used in Physics, Mathematics and Engineering. They are known as, ‘scalars’ and ‘vectors’


## Scalar

A physical quantity that possesses magnitude only, is called a scalar quantity. It can be completely specified by a number along with unit. For example, mass, time, speed, density, work, distance, length, perimeter, area, volume, etc.

## Vector

A physical quantity that possesses both magnitude and direction, is called a vector quantity. For example, weight, displacement, momentum, force, velocity, acceleration, electric field etc.

### Scalar and Vector




### Class Activity:

Look at the table below and categorize the quantities as scalar or vector.

S. No	Quantity	Category
1	7 m	
2	40 m/sec, West	
3	144 square metre	
4	32 °C	
5	98 Newtons	
6	6540 Calories	
7	5 km, East	


### Geometrical Representation of a Vector

Geometrically a vector is represented by a line segment with an arrow head at its one end (the ray). The length of the line segment describes the magnitude and the arrow head indicates the direction of the vector.



In the figure, the end A of the ray AB is called the tail or the initial point of the vector and the end B is called the terminal point with the arrow head denoted by  $\overrightarrow{AB}$ .

Usually, the vectors are denoted by bold face letters  $a, b, c$  or  $\vec{a}, \vec{b}, \vec{c}$  or  $\underline{a}, \underline{b}, \underline{c}$  etc.



### Position Vector

The vector used to specify the position of a point P with respect to origin O is called position vector of P. The tail of this vector is at origin and head at the point P. Thus  $\overrightarrow{OP}$  is the position vector of point P with respect to O.

### Result:

If  $\vec{a}$  and  $\vec{b}$  are position vectors of points A and B respectively, then  $\overrightarrow{AB} = \vec{b} - \vec{a}$ .

### Proof:

If  $\vec{a}$  and  $\vec{b}$  are position vectors of points A and B respectively, then:



$$\overrightarrow{OA} = \vec{a} \text{ and } \overrightarrow{OB} = \vec{b}$$

Using triangle law of addition, we have:

$$\overrightarrow{OA} + \overrightarrow{AB} = \overrightarrow{OB}$$

$$\Rightarrow \overrightarrow{AB} = \overrightarrow{OB} - \overrightarrow{OA}$$


$$\Rightarrow \overrightarrow{AB} = \vec{b} - \vec{a}$$



### Rectangular or Cartesian Coordinate System

A rectangular or Cartesian coordinate system consists of two number lines which are mutually perpendicular.

Their point of intersection is called the origin, usually denoted by the letter 'O'. Horizontal line is called x-axis and vertical line is called y-axis. Ordered pairs of numbers can be represented on a Cartesian plane, which is sometimes called the Coordinate plane or xy-plane. Each point in the plane corresponds to an ordered pair  $(x, y)$  of real numbers  $x$  and  $y$ , called the coordinates of the point.



#### Key Fact:

Cartesian plane has got this name after the French mathematician Rene Descartes (1596 – 1650). He developed the idea of the coordinate plane and the correspondence of ordered pairs of numbers to points in the coordinate plane. Each point in the plane can correspond to only one ordered pair and vice versa.

# Vector in a Plane

Let  $\mathbb{R}$  be the set of real numbers, then the Cartesian plane is defined as:


$$\mathbb{R}^2 = \{(x, y) : x, y \in \mathbb{R}\}$$

Any point  $P(x, y)$  is uniquely determined by its coordinates  $x$  and  $y$ . For a given vector  $\vec{u}$  in the plane, there exists a unique point  $P(x, y)$  in the plane such that:

$$\vec{u} = \overrightarrow{OP}$$

For the point  $P(x, y)$ , a unique ordered pair  $[x, y]$  is used for the vector  $\vec{u}$  called position vector of  $P(x, y)$ . Thus:

$$\vec{u} = \overrightarrow{OP} = [x, y]$$




## Negative of a Vector

A vector having the same magnitude but opposite in direction of a given vector  $\vec{b}$  is called the negative of  $\vec{b}$  and is denoted by  $-\vec{b}$ .

If  $\vec{u} = [x, y]$ , then negative vector of  $\vec{u}$  is defined as:

$$-\vec{u} = [-x, -y]$$



## Example:

$P(3, -4)$  is a point in the plane. Find the position vector  $\overrightarrow{OP}$  and then its negative vector.

### Solution:

Given is  $P(3, -4)$  in the plane.

Position vector of  $P = \overrightarrow{OP} = [3, -4]$

Negative vector of  $\overrightarrow{OP} = -\overrightarrow{OP} = -[3, -4] = [-3, 4]$

## Zero or Null Vector

If the initial and terminal points of a vector coincide then the vector has zero length.

This vector is called zero vector and is denoted by  $\vec{0}$ . The zero vector has no direction. So, it can be assigned as convenient direction according to the situation. The zero vector is denoted by:


$$\vec{0} = [0, 0]$$

## Magnitude (Norm) of a Vector

Given the vector  $\vec{u} = \overrightarrow{AB}$ . The magnitude or the length or the norm of the vector  $\vec{u}$  is denoted by  $|\vec{u}| = |\overrightarrow{AB}|$ .

If  $\vec{u} = [x, y]$ , then:

$$|\vec{u}| = \sqrt{x^2 + y^2}$$



### Key Fact:

- (i) If  $\vec{u}$  is any vector, then  $|\vec{u}| \geq 0$ .
- (ii)  $|\vec{u}| = 0$  if and only if  $\vec{u} = \vec{0}$ .
- (iii)  $|c\vec{u}| = |c||\vec{u}|$  where  $c$  is a scalar.
- (iv)  $|-u| = |\vec{u}|$

## Unit Vector

A vector which is in the direction of a non-zero vector  $\vec{u}$  and has magnitude 1 is called a unit vector of  $\vec{u}$  and is denoted by  $\hat{u}$ . If  $\vec{u}$  is non-zero vector of arbitrary length  $|\vec{u}|$  then:

$$\hat{u} = \frac{\vec{u}}{|\vec{u}|} \Rightarrow \vec{u} = |\vec{u}| \hat{u}$$

This mean any vector  $\vec{u}$  can be constructed by multiplying the magnitude of the vector to its unit vector. The process of finding the unit vector of a vector  $\vec{u}$  is called normalizing the vector  $\vec{u}$ .

If  $\vec{u} = [x, y]$ , then its unit vector is written as:

$$\hat{u} = \frac{[x, y]}{\sqrt{x^2+y^2}} = \left[ \frac{x}{\sqrt{x^2+y^2}}, \frac{y}{\sqrt{x^2+y^2}} \right]$$

### Example:

Find magnitude of the vector  $\vec{u} = [-5, 12]$ . Also find unit vector of  $\vec{u}$ .

**Solution:**  $\vec{u} = [-5, 12]$

Here,  $x = -5, y = 12$

Magnitude of  $\vec{u} = |\vec{u}| = \sqrt{x^2 + y^2} = \sqrt{(-5)^2 + (12)^2} = \sqrt{25 + 144} = \sqrt{169} = 13$  units

Unit vector of  $\vec{u} = \hat{u} = \frac{\vec{u}}{|\vec{u}|} = \frac{[-5, 12]}{13} = \left[ \frac{-5}{13}, \frac{12}{13} \right]$

### Scalar Multiplication

If  $\lambda$  is a non-zero scalar and  $\vec{a}$  is a non-zero vector then the scalar multiple  $\lambda\vec{a}$  is a vector whose magnitude is  $\lambda$  times the magnitude  $\vec{a}$ .

- $\vec{a}$  and  $\lambda\vec{a}$  have the same direction if  $\lambda > 0$ .
- $\vec{a}$  and  $\lambda\vec{a}$  have the opposite direction if  $\lambda < 0$ .
- If  $\lambda\vec{a} = 0$  then either  $\lambda = 0$  or  $\vec{a} = 0$ .
- If  $\lambda = 0$ , then  $\lambda\vec{a}$  is a zero vector.

Note: If  $\vec{u} = [x, y]$  then product of  $\lambda$  with  $\vec{u}$  is  $\lambda\vec{u} = \lambda[x, y] = [\lambda x, \lambda y]$ .


For example, if  $\vec{u} = [-6, 4]$  and  $\lambda = 3$ , then:

$$3\vec{u} = 3[-6, 4] = [-18, 12]$$

### Example:

Using graph paper draw the vectors: (i)  $-\vec{u}$  (ii)  $3\vec{u}$  (iii)  $1.5\vec{u}$ , when  $\vec{u}$  is given.

### Solution:




**Example:**

In the figure, direction of  $\vec{p}$  is 3 square units to the right and 4 square units up. Find the relation of other vectors with  $\vec{p}$ .

**Solution:**

- (i)  $\vec{q} = 2\vec{p}$
- (ii)  $\vec{r} = -1.5\vec{p}$
- (iii)  $\vec{s} = -0.5\vec{p}$
- (iv)  $\vec{t} = 0.5\vec{p}$
- (v)  $\vec{u} = -\vec{p}$

**Translation by Vector**

Translation by a vector is a process that moves a point or figure in space by a given vector. It is a type of transformation that slides a figure without rotating it, so the shape, size, and orientation of the figure do not change.

**Geometrically**

Place the tail of the arrow at the point, and the translated point is at the head of the arrow.

**Algebraically**

Represent the points and the vector by their coordinates. The translation vector's coordinates indicate how far to move the points along each coordinate axis.

For example, the vector  $[a, b]$  moves points  $a$  units horizontally and  $b$  units vertically.


For example, a translation vector that moves a figure 4 units right and 2 units down can be represented mathematically as  $[4, -2]$ , or graphically as shown in the adjoining figure. It doesn't matter where the vector is positioned in the plane. In this figure, the vector starts at  $(-2, 2)$  and ends at  $(2, 0)$ . But the initial point and terminal point of the vector is irrelevant. What matters is the length of the vector and the direction in which it points, so all you have to look at is how many units the vector moves in the y-direction and how many units the vector moves in the x-direction.

**Example:**

Use the translation vector  $[2, -1]$  to find the new coordinates of triangle ABC when:  
 $A = (-3, 2)$ ,  $B = (-1, 0)$  and  $C = (-2, -1)$

**Solution:**

The vector indicates a translation of 2 units to the right and 1 unit down. We can therefore add 2 to all of the x-values and subtract 1 from all of the y-values to find the vertices of the image.



First let's write down the coordinates of the pre-image, triangle ABC.

$$A = (-3, 2), B = (-1, 0), C = (-2, -1)$$



Now we can make the calculations to translate each vertex.

$$A' = (-3 + 2, 2 - 1) = (-1, 1)$$

$$B' = (-1 + 2, 0 - 1) = (1, -1)$$

$$C' = (-2 + 2, -1 - 1) = (0, -2)$$

The image after the translation is shown in the adjoining figure.



## Parallel Vectors

Two non-zero vectors  $\vec{u}$  and  $\vec{v}$  are said to be parallel if  $\vec{u} = \lambda\vec{v}$ , where  $\lambda$  is a scalar. If  $\lambda > 0$  then both vectors have the same direction and if  $\lambda < 0$  then both are in the opposite direction.

The vectors which are in the opposite direction are known as anti-parallel vectors.

In component form, when  $\vec{u} = [x_1, y_1]$  and  $\vec{v} = [x_2, y_2]$  are parallel vectors, then:

$$\begin{aligned}\vec{u} = \lambda\vec{v} &\Rightarrow [x_1, y_1] = \lambda[x_2, y_2] \\ \Rightarrow [x_1, y_1] &= [\lambda x_2, \lambda y_2] \Rightarrow x_1 = \lambda x_2 \text{ and } y_1 = \lambda y_2\end{aligned}$$

### Example:

Prove that  $\vec{a} = [-12, -15]$  and  $\vec{b} = [4, 5]$  are parallel vectors.

**Solution:** Given that  $\vec{a} = [-12, -15]$  and  $\vec{b} = [4, 5]$

$$\vec{a} = [-12, -15] = -3[4, 5] = -3\vec{b}$$

As  $\vec{a}$  is scalar multiple of  $\vec{b}$ , therefore  $\vec{a} \parallel \vec{b}$ .


### Equal Vectors

Two vectors  $\vec{u}$  and  $\vec{v}$  are said to be equal if both have the same magnitude and direction.

If vectors  $\vec{u}$  and  $\vec{v}$  are equal then we write  $\vec{u} = \vec{v}$ .

If  $\vec{u} = [x_1, y_1]$  and  $\vec{v} = [x_2, y_2]$ , then  $\vec{u} = \vec{v}$  implies that:

$$[x_1, y_1] = [x_2, y_2] \Rightarrow x_1 = x_2 \text{ and } y_1 = y_2$$



### Key Fact:

- It is not necessary for the equal vectors to have the same position.
- Geometrically two vectors are equal if they are translation of one another.
- If in the relation  $\vec{a} = \lambda\vec{b}$  and  $\lambda = 1$  then  $\vec{a} = \vec{b}$ .
- Equal vectors are also parallel.

**Example:**


If  $\vec{p} = [-c + 4, -2]$  and  $\vec{q} = [4, 2d]$  are equal vectors, find the values of  $c$  and  $d$ .

**Solution:**

$$\begin{aligned}\text{Given that, } \vec{p} = \vec{q} &\Rightarrow [-c + 4, -2] = [4, 2d] \\ &\Rightarrow -c + 4 = 4, -2 = 2d \\ &\Rightarrow -c = 4 - 4, -2 \div 2 = d \\ &\Rightarrow c = 0, d = 1\end{aligned}$$

**Check Point:**

Are the vectors equal? If not why?

**Addition of Vectors****Head to Tail Rule**

To add non-zero vectors  $\vec{p}$  and  $\vec{q}$  join the tail of the second vector  $\vec{q}$  with the head of the first vector  $\vec{p}$ . Now the vector obtained by joining the tail of the first vector to the head of the second vector is the vector  $\vec{p} + \vec{q}$  called the resultant vector of  $\vec{p}$  and  $\vec{q}$ .



This method for the addition of two vectors is called head to tail rule of addition. Since  $\vec{p}, \vec{q}$  and  $\vec{p} + \vec{q}$  are along the sides of a triangle XYZ, so this method of addition of vectors is also known as triangle law of addition.

**Parallelogram Law of Addition**

Consider any parallelogram PQRS. Let  $\overrightarrow{PQ} = \vec{p}$  and  $\overrightarrow{QR} = \vec{q}$ .


Since the vector  $\overrightarrow{QR}$  has the same magnitude and direction as that of  $\overrightarrow{PS}$ . Similarly,  $\overrightarrow{SR}$  has the same magnitude and direction as that of  $\overrightarrow{PQ}$ . Therefore:

$$\overrightarrow{PQ} = \overrightarrow{SR} \text{ and } \overrightarrow{QR} = \overrightarrow{PS}$$

Using triangle law of addition, we have:

$$\overrightarrow{PQ} + \overrightarrow{QR} = \overrightarrow{PR}$$

$$\Rightarrow \vec{p} + \vec{q} = \overrightarrow{PR}$$



Showing that the diagonal vector  $\overrightarrow{PR}$  of the parallelogram is the sum of the vectors of  $\vec{p}$  and  $\vec{q}$ . This is known as parallelogram law of addition. Furthermore:

$$\text{From } \Delta PQR, \quad \overrightarrow{PR} = \vec{p} + \vec{q} \quad (\text{i})$$

$$\text{From } \Delta PSR, \quad \overrightarrow{PR} = \vec{q} + \vec{p} \quad (\text{ii})$$

$$\therefore \vec{p} + \vec{q} = \vec{q} + \vec{p} \quad [\text{from (i) and (ii)}]$$

This shows that vector addition is commutative.

**Key Fact:**

For any vector  $\vec{a}$ :

- $\vec{a} + \vec{0} = \vec{0} + \vec{a} = \vec{a}$
- $\vec{a} + (-\vec{a}) = (-\vec{a}) + \vec{a} = \vec{0}$


## Polygon Law of Vector Addition

The process for the addition of vectors can be extended to any number of vectors. For instance, let we have four vectors  $\vec{a}, \vec{b}, \vec{c}, \vec{d}$  and we need to find their sum.

From the figure,

$$\overrightarrow{OA} = \vec{a}, \overrightarrow{AB} = \vec{b}, \overrightarrow{BC} = \vec{c}, \overrightarrow{CD} = \vec{d}$$

$$\begin{aligned}\text{Now } \vec{a} + \vec{b} + \vec{c} + \vec{d} &= \overrightarrow{OA} + \overrightarrow{AB} + \overrightarrow{BC} + \overrightarrow{CD} \\ &= (\overrightarrow{OA} + \overrightarrow{AB}) + \overrightarrow{BC} + \overrightarrow{CD} \\ &= \overrightarrow{OB} + \overrightarrow{BC} + \overrightarrow{CD} \quad (\because \overrightarrow{OA} + \overrightarrow{AB} = \overrightarrow{OB}) \\ &= (\overrightarrow{OB} + \overrightarrow{BC}) + \overrightarrow{CD} \\ &= \overrightarrow{OC} + \overrightarrow{CD} \quad (\because \overrightarrow{OB} + \overrightarrow{BC} = \overrightarrow{OC}) \\ &= \overrightarrow{OD}\end{aligned}$$



Here,  $\overrightarrow{OD}$  is the sum of all these four vectors. The same method is adopted to find the sum of any number of vectors. This method is called polygon law of addition of vectors.

**Explanation:** If  $\vec{u} = [x_1, y_1]$  and  $\vec{v} = [x_2, y_2]$ ,  $\vec{w} = [x_3, y_3]$ , then:

$$\vec{u} + \vec{v} + \vec{w} = [x_1, y_1] + [x_2, y_2] + [x_3, y_3] = [x_1 + x_2 + x_3, y_1 + y_2 + y_3]$$

**Example:**

If  $\vec{a} = [4, 6]$  and  $\vec{b} = [-3, 1]$ ,  $\vec{c} = [2, -2]$  then find  $\vec{a} + 2\vec{b} + \vec{c}$ . Also find  $|\vec{a} + 2\vec{b} + \vec{c}|$ .

**Solution:**

Given that,  $\vec{a} = [4, 6]$ ,  $\vec{b} = [-3, 1]$  and  $\vec{c} = [2, -2]$

$$\begin{aligned}\text{Now, } \vec{a} + 2\vec{b} + \vec{c} &= [4, 6] + 2[-3, 1] + [2, -2] = [4, 6] + [-6, 2] + [2, -2] \\ &= [4 - 6 + 2, 6 + 2 - 2] = [0, 6]\end{aligned}$$

$$|\vec{a} + 2\vec{b} + \vec{c}| = \sqrt{0^2 + 6^2} = \sqrt{36} = 6 \text{ units}$$

## Subtraction of Two Vectors

To subtract a vector, form the other we find the negative vector of the vector to be subtracted and then add it to the other vector.


Consider two non-zero vectors  $\vec{u}$  and  $\vec{v}$  then:

$$\vec{u} - \vec{v} = \vec{u} + (-\vec{v}).$$

From the figure,  $\overrightarrow{AB} = \vec{u}$  and  $\overrightarrow{BC} = -\vec{v}$ ; then

$$\overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC} \quad (\text{head to tail rule of addition})$$


$$\vec{u} + (-\vec{v}) = \vec{u} - \vec{v} = \overrightarrow{AC}$$



Thus,  $\overrightarrow{AC}$  is the vector which represents  $\vec{u} - \vec{v}$ .

In component form, if  $\vec{u} = [x_1, y_1]$  and  $\vec{v} = [x_2, y_2]$ , then:

$$\begin{aligned}\vec{u} - \vec{v} &= [x_1, y_1] - [x_2, y_2] \\ &= [x_1 - x_2, y_1 - y_2]\end{aligned}$$



**Example:**

If  $\vec{a} = [7, 10]$  and  $\vec{b} = [-3, 10]$ , then find  $\vec{a} - \vec{b}$  and a unit vector along  $\vec{a} - \vec{b}$ .

**Solution:**

Given that,  $\vec{a} = [7, 10]$  and  $\vec{b} = [-3, 10]$

$$\text{Now, } \vec{a} - \vec{b} = [7, 10] - [-3, 10] = [7 + 3, 10 - 10] = [10, 0]$$


$$|\vec{a} - \vec{b}| = \sqrt{10^2 + 0^2} = \sqrt{100} = 10 \text{ units}$$

$$\text{Unit vector along } \vec{a} - \vec{b} = \frac{\vec{a} - \vec{b}}{|\vec{a} - \vec{b}|} = \frac{[10, 0]}{10} = \left[ \frac{10}{10}, \frac{0}{10} \right] = [1, 0]$$

**Example:**

Draw vectors  $\vec{a} = 4$  units east and  $\vec{b} = 3$  units north on graph paper. Then, draw the following vectors on the graph paper.

- (i)  $\vec{a} + \vec{b}$  (ii)  $2\vec{a} - \vec{b}$  (iii)  $2\vec{a} + 1.5\vec{b}$  (iv)  $(\vec{a} + \vec{b}) + (2\vec{a} - \vec{b})$

**Solution:****Representation of a Vector in Cartesian Plane**

We denote the unit vector along  $x$ -axis by  $\hat{i}$  and along  $y$ -axis by  $\hat{j}$ .

Let us consider a point  $P(x, y)$  in the Cartesian plane.

We want to find the position vector  $\overrightarrow{OP}$  in terms of unit vectors  $\hat{i}$  and  $\hat{j}$ .

Draw perpendicular from  $P$  on  $x$ -axis meeting the  $x$ -axis at point  $A$ . As we know that for any vector  $\vec{u}$ :

$$\hat{u} = \frac{\vec{u}}{|\vec{u}|} \text{ or } \vec{u} = |\vec{u}| \hat{u}$$

i.e., Every vector is equal to the product of magnitude with its unit vector. So,


$$\overrightarrow{OA} = |\overrightarrow{OA}| \hat{i} = x \hat{i}$$

$$\overrightarrow{AP} = |\overrightarrow{AP}| \hat{j} = y \hat{j}$$

$$\text{Now, } \overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$$

$$\Rightarrow \overrightarrow{OP} = x \hat{i} + y \hat{j}$$

$$\Rightarrow \vec{r} = x \hat{i} + y \hat{j}$$



This is the position vector of the point  $P(x, y)$  lying in the Cartesian plane.

**Check Point:**

If  $\vec{a} = [3, 4]$  and  $\vec{b} = [-3, 4]$ , then find:

$$5\vec{a} + \vec{b} \text{ and } 3\vec{a} - 2\vec{b}$$

## Components of Vector When its Tail is not at Origin

Consider a vector  $\overrightarrow{PQ}$  with end points  $P(x_1, y_1)$  and  $Q(x_2, y_2)$ .

Draw perpendiculars from P and Q on  $x$ -axis meeting at S and T respectively.

Now from figure:

$$|\overrightarrow{PR}| = |\overrightarrow{ST}| = |\overrightarrow{OT}| - |\overrightarrow{OS}| = x_2 - x_1$$

$$|\overrightarrow{RQ}| = |\overrightarrow{TQ}| - |\overrightarrow{TR}|$$

$$= |\overrightarrow{TQ}| - |\overrightarrow{SP}| = y_2 - y_1$$


$$\therefore \overrightarrow{PR} = |\overrightarrow{PR}| \hat{i} = (x_2 - x_1) \hat{i}$$

$$\text{and } \overrightarrow{RQ} = |\overrightarrow{RQ}| \hat{j} = (y_2 - y_1) \hat{j}$$

$$\text{As, } \overrightarrow{PQ} = \overrightarrow{PR} + \overrightarrow{RQ}$$

Therefore:

$$\overrightarrow{PQ} = (x_2 - x_1) \hat{i} + (y_2 - y_1) \hat{j}$$



### Key Fact :

- In the relation,  $\vec{r} = x\hat{i} + y\hat{j}$ :  
 $x$  is called  $x$ -component of  $\vec{r}$  and  $y$  is called  $y$ -component of  $\vec{r}$ .
- $x\hat{i}$  and  $y\hat{j}$  are known as parts of vector  $\vec{r}$  in the direction of  $\hat{i}$  and  $\hat{j}$  respectively.
- When the tail of a vector is not at origin then it is not a position vector.

## Rectangular Components of a Vector

Consider a vector  $\overrightarrow{OP} = \vec{F}$  making angle  $\theta$  with  $x$ -axis as shown in the adjoining figure. We want to find its components that are perpendicular to each other.

If we draw perpendicular AP on  $x$ -axis, then from figure:

$$\vec{F} = \overrightarrow{F_x} + \overrightarrow{F_y} = F_x \hat{i} + F_y \hat{j} \quad \dots \dots \quad (\text{i})$$

Here,  $\overrightarrow{F_x}$  and  $\overrightarrow{F_y}$  are rectangular components of  $\vec{F}$  while  $F_x$  and  $F_y$  are magnitudes of  $\overrightarrow{F_x}$  and  $\overrightarrow{F_y}$  respectively.


Now if the magnitude of  $\vec{F}$  is given, then we can find  $F_x$  and  $F_y$  using trigonometric ratios.

From right triangle OAP:

$$\frac{F_x}{|\vec{F}|} = \cos \theta \quad \text{and} \quad \frac{F_y}{|\vec{F}|} = \sin \theta$$

$$\Rightarrow F_x = |\vec{F}| \cos \theta \quad \dots \dots \quad (\text{ii})$$

$$F_y = |\vec{F}| \sin \theta \quad \dots \dots \quad (\text{iii})$$



### Key Fact :

$\overrightarrow{F_x}$  and  $\overrightarrow{F_y}$  are called horizontal and vertical components of  $\vec{F}$  respectively.

Equations (ii) and (iii) are used to find the rectangular components of a given vector making angle  $\theta$  with  $x$ -axis.

Conversely, if the rectangular components of  $\vec{F}_x$  and  $\vec{F}_y$  of a vector are given then we can find the magnitude and angle of resultant vector by using following relations.

$$|\vec{F}| = \sqrt{(F_x)^2 + (F_y)^2} \quad \dots \dots \text{(iv)} \quad \text{and} \quad \theta = \tan^{-1} \left( \frac{F_y}{F_x} \right) \quad \dots \dots \text{(v)}$$

### Example:

A ball is thrown with an initial velocity of 25 m/s at an angle of  $45^\circ$  with the horizontal. Find the horizontal and vertical components of the velocity.

### Solution:

Let  $V$  represents the velocity.

Horizontal component of velocity is given as:

$$V_x = V \cos 45^\circ = 25 \times 0.707 = 17.68 \text{ m/s}$$

Vertical component of velocity is given as:

$$V_y = V \sin 45^\circ = 25 \times 0.707 = 17.68 \text{ m/s}$$

### Example:

Two forces  $F_1$  and  $F_2$  with magnitudes 30 N and 40 N, respectively, act on an object at a point A as shown in the figure. Find the resultant force  $F$  acting at A and its magnitude.

### Solution:

First, we write  $F_1$  and  $F_2$  in component form:

$$\begin{aligned} F_1 &= F_1 \cos 45^\circ \hat{i} + F_1 \sin 45^\circ \hat{j} \\ &= 30 \times 0.707 \hat{i} + 30 \times 0.707 \hat{j} = 21.21 \hat{i} + 21.21 \hat{j} \end{aligned}$$

$$\begin{aligned} F_2 &= F_2 \cos 150^\circ \hat{i} + F_2 \sin 150^\circ \hat{j} \\ &= 40 \times (-0.866) \hat{i} + 40 \times 0.5 \hat{j} = -34.64 \hat{i} + 20 \hat{j} \end{aligned}$$

So, the resultant force  $F$  is:

$$\begin{aligned} F &= F_1 + F_2 = (21.21 \hat{i} + 21.21 \hat{j}) + (-34.64 \hat{i} + 20 \hat{j}) = (21.21 - 34.64) \hat{i} + (21.21 + 20) \hat{j} \\ &= -13.43 \hat{i} + 41.21 \hat{j} \end{aligned}$$

$$\begin{aligned} \text{Magnitude of resultant force} &= |F| = \sqrt{(-13.43)^2 + (41.21)^2} = \sqrt{180.36 + 1,698.26} \\ &= \sqrt{1,878.62} = 43.34 \text{ units} \end{aligned}$$

## Concept Related to Vectors Using Analytical Representation

### Magnitude of a Vector

In Cartesian plane a vector  $\vec{u}$  is expressed as:

$$\vec{u} = x\hat{i} + y\hat{j}$$

The magnitude of vector  $\vec{u}$  is defined as:

$$|\vec{u}| = \sqrt{x^2 + y^2}$$

$$|\vec{u}| = \sqrt{(x\text{-component})^2 + (y\text{-component})^2}$$

### Equal Vectors

Two vectors are said to be equal if and only if they have same  $x$  and  $y$  components.

If  $\vec{u} = x_1\hat{i} + y_1\hat{j}$  and  $\vec{v} = x_2\hat{i} + y_2\hat{j}$  then  $\vec{u} = \vec{v}$  if  $x_1 = x_2$  and  $y_1 = y_2$ .

### Negative of a Vector

If  $\vec{p} = x\hat{i} + y\hat{j}$  then negative of  $\vec{p}$  is:

$$-\vec{p} = (-1)\vec{p} = (-1)(x\hat{i} + y\hat{j}) = (-x)\hat{i} + (-y)\hat{j}$$

### Unit Vector

If  $\vec{w} = x\hat{i} + y\hat{j}$  then  $\vec{w}$  is called a unit vector if:

$$|\vec{w}| = 1 \Rightarrow \sqrt{x^2 + y^2} = 1 \quad \text{or} \quad x^2 + y^2 = 1$$

### Parallel Vectors

The vectors  $\vec{u} = x_1\hat{i} + y_1\hat{j}$  and  $\vec{v} = x_2\hat{i} + y_2\hat{j}$  are said to be parallel if for some scalar  $\lambda$  we have:

$$\vec{u} = \lambda\vec{v}$$

$$\Rightarrow (x_1\hat{i} + y_1\hat{j}) = \lambda(x_2\hat{i} + y_2\hat{j})$$

$$\Rightarrow x_1\hat{i} + y_1\hat{j} = \lambda x_2\hat{i} + \lambda y_2\hat{j}$$

Comparing the components, we have:

$$x_1 = \lambda x_2 \text{ and } y_1 = \lambda y_2$$

$$\Rightarrow \frac{x_1}{x_2} = \lambda \text{ and } \frac{y_1}{y_2} = \lambda$$

$$\text{or } \frac{x_1}{x_2} = \frac{y_1}{y_2} = \lambda$$

Which is the condition for the two vectors to be parallel. If  $\lambda$  is positive then  $\vec{u}$  and  $\vec{v}$  have the same direction and if  $\lambda$  is negative then  $\vec{u}$  and  $\vec{v}$  are in the opposite direction.

### Addition and Subtraction of Vectors

If  $\vec{u} = x_1\hat{i} + y_1\hat{j}$  and  $\vec{v} = x_2\hat{i} + y_2\hat{j}$  are any two vectors then:

$$(i) \quad \vec{u} + \vec{v} = (x_1\hat{i} + y_1\hat{j}) + (x_2\hat{i} + y_2\hat{j}) = (x_1 + x_2)\hat{i} + (y_1 + y_2)\hat{j}$$

$$(ii) \quad \vec{u} - \vec{v} = (x_1\hat{i} + y_1\hat{j}) - (x_2\hat{i} + y_2\hat{j}) = (x_1 - x_2)\hat{i} + (y_1 - y_2)\hat{j}$$

### Unit Vector in the Direction of another Vector

Let we have a non-zero vector  $\vec{u} = x\hat{i} + y\hat{j}$

$$\therefore |\vec{u}| = \sqrt{x^2 + y^2}$$

Now the unit vector in the direction of  $\vec{u}$  is

$$\hat{u} = \frac{\vec{u}}{|\vec{u}|} = \frac{x\hat{i} + y\hat{j}}{\sqrt{x^2 + y^2}}$$

$$\hat{u} = \left( \frac{x}{\sqrt{x^2 + y^2}} \right) \hat{i} + \left( \frac{y}{\sqrt{x^2 + y^2}} \right) \hat{j}$$

### Example:

If  $\vec{a} = 2\hat{i} - \hat{j}$  then find unit vector along the vector  $\vec{a}$ .

#### Key Fact:

A vector having both  $x$ -component and  $y$ -component zero is called a null vector denoted by  $\vec{0}$ .

$$\vec{0} = 0\hat{i} + 0\hat{j}$$

#### Key Fact:

If  $\lambda$  is a scalar and  $\vec{u} = x\hat{i} + y\hat{j}$  then product of  $\lambda$  with  $\vec{u}$  is  
 $\lambda\vec{u} = \lambda(x\hat{i} + y\hat{j}) = \lambda x\hat{i} + \lambda y\hat{j}$

**Solution:**

$$|\vec{a}| = \sqrt{2^2 + (-1)^2} = \sqrt{4 + 1} = \sqrt{5}$$

The unit vector in the direction of  $\vec{a}$  is:

$$\hat{a} = \frac{\vec{a}}{|\vec{a}|} = \frac{2\hat{i} - \hat{j}}{\sqrt{5}} = \frac{2}{\sqrt{5}}\hat{i} + \frac{-1}{\sqrt{5}}\hat{j}$$

**Example:** If  $\vec{u} = 5\hat{i} + \hat{j}$  and  $\vec{v} = 2\hat{i} - 9\hat{j}$ , then find (i)  $2\vec{u} + 3\vec{v}$  (ii)  $4\vec{u} - 2\vec{v}$

$$(i) \quad 2\vec{u} + 3\vec{v} = 2(5\hat{i} + \hat{j}) + 3(2\hat{i} - 9\hat{j}) = 10\hat{i} + 2\hat{j} + 6\hat{i} - 27\hat{j} = 16\hat{i} - 25\hat{j}$$

$$(ii) \quad 4\vec{u} - 2\vec{v} = 4(5\hat{i} + \hat{j}) - 2(2\hat{i} - 9\hat{j}) = 20\hat{i} + 4\hat{j} - 4\hat{i} + 18\hat{j} = 16\hat{i} + 22\hat{j}$$

**Example:** Check whether the vectors  $\vec{u} = 3\hat{i} + 2\hat{j}$  and  $\vec{v} = 6\hat{i} - 4\hat{j}$ , are parallel or not.

**Solution:**

Given that  $\vec{u} = 3\hat{i} + 2\hat{j}$  and  $\vec{v} = 6\hat{i} - 4\hat{j}$

$$\vec{v} = 6\hat{i} - 4\hat{j} = 2(3\hat{i} - 2\hat{j}) \neq 2\vec{u}$$


Therefore  $\vec{u}$  and  $\vec{v}$  are not parallel.

#### Check Point:

If  $\vec{a} = x\hat{i} + 2\hat{j}$  and  $\vec{b} = 6\hat{i} - (x + y)\hat{j}$  are equal vectors, find the values of  $x$  and  $y$ .

### Exercise 7.1


- Draw the following vectors.
  - 10 N force along x-axis
  - 50 m/s velocity at  $150^\circ$  with x-axis
  - 220 m displacement towards north
  - 24 m/s<sup>2</sup> acceleration at  $45^\circ$  with x-axis
- Draw the vectors on the graph paper when  $\vec{p}$  is 4cm long making an angle of  $45^\circ$  with x-axis.
  - $2\vec{p}$
  - $-\vec{p}$
  - $0.5\vec{p}$
  - $-1.5\vec{p}$
  - $-0.5\vec{p}$
- Draw vectors  $\vec{a} = 3$  units west and  $\vec{b} = 3$  units north on graph paper. Draw the following vectors on the graph paper.
  - $2\vec{a} + \vec{b}$
  - $\vec{a} - 2\vec{b}$
  - $3\vec{a} + 1.5\vec{b}$
  - $(2\vec{a} + \vec{b}) + (\vec{a} - 2\vec{b})$
  - $0.5(\vec{a} + \vec{b})$
  - $3\vec{a} - 2\vec{b}$
  - $2\vec{a} - 2.5\vec{b}$
  - $(2\vec{a} + \vec{b}) - (\vec{a} - 2\vec{b})$
- In the figure, direction of  $\vec{a}$  is 4 square units to the right and 3 square units up. Find the relation of other vectors with  $\vec{a}$ .



- A point with coordinates  $(5, -7)$  have been translated by the vector  $[0, -3]$ . Find new position of the point.

6. A vector  $[-5, 4]$  have been translated by another vector  $[4, -3]$ . Find new location of the given vector.
7. Triangle ABC is a pre-image with its vertices at  $A = (-4, 6)$ ,  $B = (-1, 4)$  and  $C = (-6, 1)$ . Find the coordinates of the image if:
- the triangle is translated by  $[5, 0]$
  - the triangle is translated by  $[-3, -4]$
8. What translation by vector would be needed to bring the point  $E(-6, 5)$  to the origin?

9. ABCD is a rectangle as shown in the adjoining figure.



Represent the following vectors in terms of  $\vec{a}$  and  $\vec{b}$ .

- $\overrightarrow{DC}$
- $\overrightarrow{DA}$
- $\overrightarrow{AC}$
- $\overrightarrow{BD}$
- $\overrightarrow{AO}$
- $\overrightarrow{BO}$
- $\overrightarrow{AB} + \overrightarrow{AC}$

10. Given that  $\overrightarrow{OP} = \vec{p} + \vec{q}$ ,  $\overrightarrow{OQ} = \vec{p} - \vec{q}$  and  $M$  is mid-point of  $PQ$ . Find in terms of  $\vec{p}$  and  $\vec{q}$ .

- $\overrightarrow{PQ}$
- $\overrightarrow{PM}$
- $\overrightarrow{QM}$
- $\overrightarrow{OM}$

11. Express  $\overrightarrow{AB}$  in the form of  $[x, y]$  and  $x\hat{i} + y\hat{j}$ , when:

- $A(1, 3)$ ,  $B(4, 6)$
- $A(-2, 5)$ ,  $B(2, -5)$
- $A(4, -7)$ ,  $B(-1, -9)$

12. Find the unit vectors of the following vectors.

- $\overrightarrow{PQ}$  when  $P(1, 4)$ ,  $Q(4, 8)$
- $\vec{u} = 5\hat{i} - 12\hat{j}$
- $\vec{b} = -6\hat{i} - 8\hat{j}$

13. If  $\vec{a} = 3\hat{i} - 2\hat{j}$  and  $\vec{b} = \hat{i} + 4\hat{j}$ , then find  $x$  and  $y$  such that  $x\vec{a} + y\vec{b} = 4\hat{i} - 12\hat{j}$ .

14. Given that  $D = (3, -2)$ ,  $E = (-3, -5)$ ,  $F = (0, 5)$  and  $G = (3, 6)$ . Find the following vectors.

- $\overrightarrow{DE}$
- $\overrightarrow{DE} + \overrightarrow{FG}$
- $3\overrightarrow{EF} - 2\overrightarrow{EG}$
- $\frac{3}{2}\overrightarrow{DF} + \frac{1}{2}\overrightarrow{EG}$
- $\overrightarrow{DF} - \overrightarrow{EF} + 2\overrightarrow{FG}$

15. If  $\vec{a} = 3\hat{i} + 5\hat{j}$  and  $\vec{b} = 2\hat{i} - 4\hat{j}$ , then find:

- $\vec{a} + 2\vec{b}$
- $2\vec{a} - 3\vec{b}$
- $5(2\vec{a} + \vec{b})$
- $0.5(\vec{a} - \vec{b})$
- $|\vec{a} + \vec{b}|$
- $|\vec{a} - \vec{b}|$
- $|\vec{a}| + 2|\vec{b}|$
- $|\vec{a}| - |\vec{b}|$

16. Find the value of  $p$  for which the vector  $\vec{u} = 3\hat{i} + 6\hat{j}$  is parallel to  $\vec{v} = \hat{i} + p\hat{j}$ .

17. If  $\vec{a} = 2\hat{i} - 4\hat{j}$ ,  $\vec{b} = \hat{i} - 2\hat{j}$  and  $\vec{c} = [1, 3]$ , then find a unit vector in the direction of:

- $\vec{a} + \vec{b}$
- $2\vec{a} - \vec{c}$
- $2\vec{a} - \vec{b}$
- $3(\vec{a} + \vec{c})$
- $\vec{a} + \vec{b} + 2\vec{c}$
- $4\vec{a} - 2\vec{b} + \vec{c}$

18. If  $\vec{a} = \hat{i} - 3\hat{j}$  and  $\vec{b} = 2\hat{i} + 4\hat{j}$  then find:

- A vector of magnitude 4 in the direction  $2\vec{a} - \vec{b}$ .
- A vector of magnitude 6 in the opposite direction of  $\vec{a} + 2\vec{b}$ .

19. An airplane leaves the airport on the bearing of  $45^\circ$  traveling at 380 mph. The wind is blowing at a bearing of  $135^\circ$  at a speed of 50 mph. What is the actual velocity of the airplane?

20. A truck leaves dry port on a bearing of  $30^\circ$  and travels 15 km. It then turns due west and travels 8 km. How far is the truck from the dry port and what is its bearing?

## Ratio Formula for Finding Position Vector of a Point on a Vector

Let  $\overline{AB}$  be any line segment and  $P$  be the point which divides this line segment in the given ratio  $m : n$  internally. The position vectors of the given points  $A$  and  $B$  are  $\vec{a}$  and  $\vec{b}$  respectively. Let  $\vec{r}$  be the position vector of point  $P$ . Given that:

$$|\overrightarrow{AP}| : |\overrightarrow{PB}| = m : n$$

$$\Rightarrow \frac{|\overrightarrow{AP}|}{|\overrightarrow{PB}|} = \frac{m}{n} \Rightarrow n|\overrightarrow{AP}| = m|\overrightarrow{PB}|$$

Because  $\overrightarrow{AP}$  and  $\overrightarrow{PB}$  have the same direction, so

$$n\overrightarrow{AP} = m\overrightarrow{PB} \quad (i)$$

$$\text{From figure, } \overrightarrow{OA} + \overrightarrow{AP} = \overrightarrow{OP}$$

$$\Rightarrow \vec{a} + \overrightarrow{AP} = \vec{r} \Rightarrow \overrightarrow{AP} = \vec{r} - \vec{a}$$

$$\text{Similarly, } \overrightarrow{OP} + \overrightarrow{PB} = \overrightarrow{OB}$$

$$\Rightarrow \vec{r} + \overrightarrow{PB} = \vec{b} \Rightarrow \overrightarrow{PB} = \vec{b} - \vec{r}$$

Substituting the values in equation (i), we get:

$$\begin{aligned} n(\vec{r} - \vec{a}) &= m(\vec{b} - \vec{r}) \Rightarrow n\vec{r} - n\vec{a} = m\vec{b} - m\vec{r} \\ \Rightarrow n\vec{r} + m\vec{r} - n\vec{b} + n\vec{a} &= (n+m)\vec{r} = m\vec{b} + n\vec{a} \Rightarrow \vec{r} = \frac{m\vec{b} + n\vec{a}}{m+n} \end{aligned}$$

### Particular Case:

If  $m = n$ , then  $P$  will be the midpoint of  $\overline{AB}$  and position vector of  $P$  in this case is:

$$\vec{r} = \frac{n\vec{b} + n\vec{a}}{n+n} = \frac{n(\vec{b} + \vec{a})}{2n} = \frac{\vec{a} + \vec{b}}{2}$$

### Example:

The position vectors of points  $A$  and  $B$  are  $2\hat{i} - \hat{j}$  and  $3\hat{i} + 2\hat{j}$  respectively. Find the position vector of point  $P$  dividing the line segment joining  $A$  and  $B$  in the ratio  $3 : 4$  internally.

### Solution:

Let  $\vec{a} = 2\hat{i} - \hat{j}$ ,  $\vec{b} = 3\hat{i} + 2\hat{j}$  and  $m : n = 3 : 4$

If  $\vec{r}$  be the position vector of  $P$ , then by using ratio formula, we have:

$$\vec{r} = \frac{m\vec{b} + n\vec{a}}{m+n} = \frac{3(3\hat{i} + 2\hat{j}) + 4(2\hat{i} - \hat{j})}{3+4} = \frac{9\hat{i} + 6\hat{j} + 8\hat{i} - 4\hat{j}}{7} = \frac{17\hat{i} + 2\hat{j}}{7} = \frac{17}{7}\hat{i} + \frac{2}{7}\hat{j}$$


## Application to Geometry

There are many theorems and geometrical problems that can be solved by using vector methods.

We solve some example here..

### Example:

What type of a quadrilateral  $ABCD$  is if  $2\overrightarrow{AB} = \overrightarrow{DC}$ ?



### Key Fact:

When the point  $P$  divides the line segment  $\overline{AB}$  in the ratio  $m : n$  externally then:


$$\vec{r} = \frac{n\vec{a} - m\vec{b}}{n-m}$$

**Solution:**

$2\vec{AB} = \vec{DC}$  implies that:

$$2|\vec{AB}| = |\vec{DC}| \text{ and } \vec{AB} \parallel \vec{DC}$$

These two conditions show that ABCD is a trapezium.

**Example:**

A( $x, y$ ), B(-2, 3), C(-3, -4) and D(4, -5) are vertices of a parallelogram taken in order. Use vector method to find vertex A.

**Solution:**


Position vectors of vertices of parallelogram are:

$$\vec{OA} = x\hat{i} + y\hat{j}, \quad \vec{OB} = -2\hat{i} + 3\hat{j}$$

$$\vec{OC} = -3\hat{i} - 4\hat{j}, \quad \vec{OD} = 4\hat{i} - 5\hat{j}$$

$$\vec{AB} = \vec{OB} - \vec{OA} = (-2\hat{i} + 3\hat{j}) - (x\hat{i} + y\hat{j}) = (-2 - x)\hat{i} + (3 - y)\hat{j}$$

$$\vec{DC} = \vec{OC} - \vec{OD} = (-3\hat{i} - 4\hat{j}) - (4\hat{i} - 5\hat{j}) = (-3 - 4)\hat{i} + (-4 + 5)\hat{j} = -7\hat{i} + \hat{j}$$



As, ABCD is a parallelogram, therefore:

$$\begin{aligned} \vec{AB} = \vec{DC} &\Rightarrow (-2 - x)\hat{i} + (3 - y)\hat{j} = -7\hat{i} + \hat{j} \\ &\Rightarrow -2\hat{i} - x\hat{i} + 3\hat{j} - y\hat{j} = -7\hat{i} + \hat{j} \Rightarrow -2\hat{i} + 7\hat{i} + 3\hat{j} - \hat{j} = x\hat{i} + y\hat{j} \\ &\Rightarrow 5\hat{i} + 2\hat{j} = x\hat{i} + y\hat{j} \Rightarrow x = 5, y = 2 \end{aligned}$$

∴ Fourth vertex of parallelogram is A(5, 2).

**Example:**

Use vectors to prove that the diagonals of a rhombus bisect each other.

**Solution:**

Consider any rhombus PQRS. Let  $\vec{p}, \vec{q}, \vec{r}$  and  $\vec{s}$  be the position vectors of the vertices P, Q, R and S respectively.

Let M and N be the mid points of diagonal vectors  $\vec{PR}$  and  $\vec{QS}$  respectively.


$$\text{p.v. of } M = \frac{\vec{p} + \vec{r}}{2} \quad (\text{i}) \quad \text{p.v. of } N = \frac{\vec{q} + \vec{s}}{2} \quad (\text{ii})$$

Since PQRS is a rhombus, therefore:

$$\begin{aligned} \vec{PQ} &= \vec{SR} \\ \Rightarrow \vec{q} - \vec{p} &= \vec{r} - \vec{s} \\ \Rightarrow \vec{q} + \vec{s} &= \vec{p} + \vec{r} \quad (\text{iii}) \end{aligned}$$

Using equation (iii) in (i) and (ii), we get:

$$\text{p.v. of } M = \text{p.v. of } N$$

**Check Point:**

Prove that the diagonals of parallelogram, square and rectangle bisect each other.

Since the position vectors of the mid-points of both the diagonals are same.

Therefore, the diagonals of a rhombus bisect each other.


- What type of a quadrilateral EFGH is if  $\overrightarrow{EF} = \overrightarrow{HG}$  and  $\overrightarrow{EH} = \overrightarrow{FG}$ ?
- Use vectors to find the length of diagonals of a parallelogram having adjacent sides  $2\hat{i} + \hat{j}$  and  $\hat{i} - 3\hat{j}$ .
- (i) Show that  $A(0, 2), B(\sqrt{3}, -1), C(0, -2)$  are the vertices of a right-angled triangle.  
(ii) Show that  $A(3, 1), B(-2, -3), C(2, 2)$  are the vertices of an isosceles triangle.
- The position vectors of points C and D are  $3\hat{i} + 2\hat{j}$  and  $\hat{i} - 4\hat{j}$  respectively. Find the position vector of point O dividing the line segment joining C and D in the ratio 2 : 3 internally.
- If  $M(2, 4)$  is mid-point of  $\overline{XY}$  such that  $X(-1, 3)$  and  $Y(x, y)$ . Find the point Y.
- If  $\overrightarrow{OA} = -\hat{i} + 3\hat{j}$ ,  $\overrightarrow{OB} = 6\hat{i} + 2\hat{j}$ , find the point trisecting the join of  $\overline{AB}$ .
- Find x, such that the points  $P(-1, x), Q(3, 2)$  and  $R(7, 3)$  are collinear.
- Find the coordinates of vertex D of a parallelogram ABCD if  $A(-3, 0), B(1, -2)$  and  $C(5, 0)$  are its three vertices.
- If  $\overrightarrow{AB} = \overrightarrow{CD}$  and  $A(0, 2), C(-2, 4), D(-1, 5)$ , then find vertex B.
- Vertices of a quadrilateral are  $U(9, 4), V(-7, 7), W(-4, -7)$  and  $X(5, -5)$ . Find mid points of its sides and prove that the figure obtained by joining mid points consecutively is a parallelogram.
- Prove that the line segments joining the midpoints of two sides of a triangle are half in length of third side.
- Prove that the joining the midpoints of the two non-parallel sides of a trapezium is parallel to its parallel sides.
- Prove that the line segment joining mid points of adjacent sides of a square or rectangle or parallelogram divides the corresponding diagonal in the ratio 1 : 3.

**I have Learnt**

- Introduction of rectangular coordinate system in a plane.
- Representing vector as a directed line segment.
- Expressing a vector in terms of two non-zero and non-parallel coplanar vector.
- Expressing a vector in terms of position vector.
- Expressing translation by a vector and finding the magnitude of a vector.
- Adding and subtracting vectors, and multiplying a vector by a scalar.
- Solving geometrical problems involving the use of a vector.
- Applying concepts of vectors in geometrical problems such as parallel and perpendicular lines in geometrical shapes, vector projectile motion, crosswinds aviation, military usage, designing roller coasters.

1. Choose the correct option.
- Which of the following is a scalar quantity?  
a. velocity      b. speed      c. torque      d. force
  - Which of the following is a vector quantity?  
a. velocity      b. speed      c. distance      d. work
  - If  $\vec{a}$  and  $\vec{b}$  are position vectors of points A and B respectively, then  $\overrightarrow{AB}$  is:  
a.  $\vec{b} + \vec{a}$       b.  $\vec{a} - \vec{b}$       c.  $\vec{b} - \vec{a}$       d.  $-\vec{b} - \vec{a}$
  - Which of the following is not a symbol of vector  $a$ ?  
a.  $\vec{a}$       b.  $\underline{a}$       c.  $a$       d.  $|\vec{a}|$
  - If  $\overrightarrow{OP} = [-6, 7]$ , then  $-\overrightarrow{OP}$  is equal to:  
a.  $[-6, 7]$       b.  $[6, 7]$       c.  $[6, -7]$       d.  $[-6, -7]$
  - If  $\vec{u} = -5\hat{i} + 12\hat{j}$ , then  $|\vec{u}|$  is equal to:  
a. 17      b. 7      c. 169      d. 13
  - Given that  $\vec{u}$  is any vector. Which of the following is true?  
a.  $|- \vec{u}| = |\vec{u}|$       b.  $-|\vec{u}| = |\vec{u}|$       c.  $|\vec{u}| + |- \vec{u}| = 0$       d.  $|\vec{u}| = 0$
  - The unit vector of the vector  $\vec{u} = 6\hat{i} + 10\hat{j} - 2\hat{j}$  is:  
a.  $\frac{3\hat{i}}{5} - \frac{4\hat{j}}{5}$       b.  $\frac{3\hat{i}}{5} + \frac{4\hat{j}}{5}$       c.  $-\frac{3\hat{i}}{5} - \frac{4\hat{j}}{5}$       d.  $-\frac{3\hat{i}}{5} + \frac{4\hat{j}}{5}$
  - If  $\vec{a} = \lambda\vec{b}$ ,  $\vec{a} = 12\hat{i} - 18\hat{j}$  and  $\vec{b} = -2\hat{i} + 3\hat{j}$ , then  $\lambda$  is equal to:  
a. 3      b. -3      c. 6      d. -6
  - If  $\vec{u} = [-5x, 8]$  and  $\vec{v} = [10, 4y]$  are equal vectors, then:  
a.  $x = 2, y = 2$       b.  $x = -2, y = -2$       c.  $x = -2, y = 2$       d.  $x = 2, y = -2$
  - If  $\vec{p} = [5, -6]$  and  $\vec{q} = [2, 6]$ , then  $\vec{p} - 2\vec{q}$  is:  
a.  $[1, -18]$       b.  $[9, 18]$       c.  $[1, 18]$       d.  $[-9, -18]$
  - If  $\vec{u} = 5\hat{i} + 10\hat{j}$  and  $\vec{v} = 4\hat{j}$ , then  $|\vec{u} - \vec{v}|$  is:  
a.  $\sqrt{123}$       b.  $\sqrt{61}$       c.  $\sqrt{11}$       d.  $-\sqrt{61}$
  - Which of the following vectors represents a position vector?  
a.  $\overrightarrow{OP}$       b.  $-\overrightarrow{OP}$       c.  $\overrightarrow{PO}$       d.  $\overrightarrow{PQ}$
  - What type of a quadrilateral ABCD is, if  $\overrightarrow{AB} = \frac{2}{3} \overrightarrow{DC}$ ?  
a. kite      b. rectangle      c. trapezium      d. rhombus
2. Given that  $\vec{p} = 3\hat{i} - 4\hat{j}$  and  $\vec{q} = -3\hat{i} - 4\hat{j}$ . Prove that  $|\vec{p}| = |\vec{q}|$ . Is  $\vec{p} = \vec{q}$ ?
3. If  $\vec{a} = 2\hat{i} - 4\hat{j}$  and  $\vec{b} = -2\hat{i} + x\hat{j}$ , then find the value of  $x$  if  $|\vec{a} + 2\vec{b}| = 6$ .

4. In  $\Delta OAB$ ,  $\overrightarrow{OA} = \vec{a}$ ,  $\overrightarrow{OB} = \vec{b}$  and M is mid-point of  $\overrightarrow{OA}$ .  
 Write the following vectors in terms of  $\vec{a}$  and  $\vec{b}$ .  
 (i)  $\overrightarrow{OM}$     (ii)  $\overrightarrow{AM}$     (iii)  $\overrightarrow{BM}$     (iv)  $\overrightarrow{OB} + \overrightarrow{BA}$



5. Two tractors are pulling a truck stuck in the mud. First tractor is pulling with a force of 250 N at an angle of  $50^\circ$  with the horizontal while the second tractor is pulling with a force of 300 N at an angle of  $40^\circ$  with the horizontal. What is the magnitude and direction of the resultant force?
6. Ammar and Javed are playing with a ball. Ammar throws two balls for Javed to catch. The path of the two balls is described by the vectors  $\mathbf{V}_1 = 120\hat{i} + 12\hat{j}$  and  $\mathbf{V}_2 = 90\hat{i} - 30\hat{j}$  where the distances are expressed in meters.  
 (i) How much farther did the first ball travel as compared with the second ball?  
 (ii) What is the distance between the two balls thrown?
7. Ahmad can swim in still water with a speed of 6 m/s. He goes swimming in a river Chenab which has a current flowing towards west with a speed of 1.5 m/s. Assuming his speed is the same, what is his resultant velocity if:  
 (i) He tries to swim due west along the current?  
 (ii) He tries to swim due east against the current?  
 (iii) He tries to swim north across the river?
8. A plane is travelling north with a speed of 150 km/h. A steady wind is blowing due east with a speed of 50 km/h.  
 (i) What is the resultant speed of the plane?  
 (ii) How far is the plane from its starting point after 10 hours?

# 08

## Application of Trigonometry

**After studying this unit students will be able to:**

- Extend sine and cosine functions to angles between  $90^\circ$  and  $180^\circ$ .
- Solve problems using laws of sines, cosines and the area formula for any triangle.
- Solve simple trigonometric problems in three dimensions.
- Apply concepts of trigonometry to real life world problems such as video games, flight engineering, navigation and sound waves.

Burj Khalifa at Dubai is the tallest building in the world with tallest free-standing structure and highest number of stories. It is 828 metres (2,716.5 feet) high and consists of more than 160 stories. How do scientists know its true height? A common way to measure the height involves determining the angle of elevation, which is formed by the top of building and the ground at a point some distance away from the base of the building. This method is much more practical than climbing the building and dropping a very long tape measure or rope.



# Trigonometric Ratios

## Quadrantal Angles

In a coordinate plane the two axes divide the plane in four equivalent parts called quadrants. If terminal ray of an angle in standard position coincides with any axis, then it is called quadrantal angle. The measure of a quadrantal angle is multiple of  $90^\circ$ . For example;  $0^\circ, 90^\circ, 180^\circ, 270^\circ, 360^\circ, 450^\circ, \dots$  are quadrantal angles.

In radian measure quadrantal angle is a multiple of  $\frac{\pi}{2}$ .

For example;  $0, \frac{\pi}{2}, \pi, \frac{3\pi}{2}, 2\pi, \frac{5\pi}{2}, 3\pi, \dots$  are quadrantal angles.

## Trigonometric Ratios

The ratio between any two sides of a right-angled triangle is called trigonometric ratio.

In triangle ABC,  $\angle C = 90^\circ$  and  $\angle A = \theta$ .

With respect to acute angle  $\theta$ , BC =  $a$  is length of perpendicular, AC =  $b$  is length of base and AB =  $c$  is length of hypotenuse.

Various trigonometric ratios are defined as:

- The ratio of perpendicular and hypotenuse is sine of  $\theta$  denoted by  $\sin \theta$ .

$$\therefore \sin \theta = \frac{\text{Perpendicular}}{\text{Hypotenuse}} = \frac{a}{c}$$


- The ratio of base and hypotenuse is cosine of  $\theta$  denoted by  $\cos \theta$ .

$$\therefore \cos \theta = \frac{\text{Base}}{\text{Hypotenuse}} = \frac{b}{c}$$

- The ratio of perpendicular and base is tangent of  $\theta$  denoted by  $\tan \theta$ .

$$\therefore \tan \theta = \frac{\text{Perpendicular}}{\text{Base}} = \frac{a}{b}$$

- Furthermore,  $\operatorname{cosec} \theta = \frac{\text{Hypotenuse}}{\text{Perpendicular}} = \frac{c}{a} = \frac{1}{\sin \theta}$ ,  $\sec \theta = \frac{\text{Hypotenuse}}{\text{Base}} = \frac{c}{b} = \frac{1}{\cos \theta}$   
and  $\cot \theta = \frac{\text{Base}}{\text{Perpendicular}} = \frac{b}{a} = \frac{1}{\tan \theta}$



## Trigonometric Ratios with the help of Unit Circle

A circle whose radius is 1 unit is called a unit circle.

Consider a unit circle with centre O.

Take a point P( $x, y$ ) on the circle. Join P with O.

Draw PC perpendicular to x-axis.

Let OC =  $x$ , PC =  $y$  and OP = 1 (unit circle).


If  $\angle COP = \theta$  where  $0 < \theta < 180^\circ$ , then:

$$\cos \theta = \frac{x}{1} \text{ and } \sin \theta = \frac{y}{1} \dots \text{ (i)}$$

Other trigonometric ratios can be defined as:

$$\tan \theta = \frac{y}{x}, \cot \theta = \frac{x}{y}, \sec \theta = \frac{1}{x} \text{ and } \operatorname{cosec} \theta = \frac{1}{y}$$

Now from (i), we have:




$$x = \cos \theta \text{ and } y = \sin \theta$$

So,  $P(x, y) = P(\cos \theta, \sin \theta)$ .

$\therefore$  If  $P$  is a point on a unit circle with centre  $O$  and  $\angle COP = \theta$ , then coordinates of  $P$  are:  $(\cos \theta, \sin \theta)$ .

### Signs of Trigonometric Ratios in Different Quadrants

As  $\cos \theta = x$  and  $\sin \theta = y$  in the unit circle. Therefore, depending upon the signs of  $x$  and  $y$  in different quadrants, the signs of trigonometric ratios change. The adjoining figure shows signs of trigonometric ratios in different quadrants.



### Trigonometric Ratios between $90^\circ$ and $180^\circ$

We have discussed finding the sine and cosine for angles in the first quadrant, but what if our angle is in second quadrant ( $90^\circ < \theta < 180^\circ$ )?

In the diagram,  $R(1, 0)$  and  $P(a, b)$  are points on the unit circle. Under a reflection in the  $y$ -axis, the image of  $P(a, b)$  is  $P'(-a, b)$  and the image of  $R(1, 0)$  is  $R'(-1, 0)$ .

Since angle measure is preserved under a line reflection,  
 $\therefore \angle ROP = \angle R'OP'$

Also  $\angle ROP'$  and  $\angle R'OP'$  are supplementary angles.

$$\therefore \angle ROP' + \angle R'OP' = 180^\circ \quad (\text{i})$$

$$\text{But, } \angle R'OP' = \angle ROP$$

Therefore, (i) becomes:

$$\angle ROP' + \angle ROP = 180^\circ$$

$$\angle ROP = 180^\circ - \angle ROP' = 180^\circ - \theta$$

Case 1: If  $\angle ROP' = \theta$ , then:

$$\cos \theta = -a \text{ and } \sin \theta = b \dots \text{(ii)} \text{ (using the properties of unit circle)}$$

Case 2: If  $\angle ROP = 180^\circ - \theta$ , then:


$$\cos(180^\circ - \theta) = a \text{ and } \sin(180^\circ - \theta) = b \dots \text{(iii)}$$

Comparing (ii) and (iii), we get:

$$\cos(180^\circ - \theta) = -\cos \theta \quad \text{and} \quad \sin(180^\circ - \theta) = \sin \theta$$

$$\text{As, } \frac{\sin(180^\circ - \theta)}{\cos(180^\circ - \theta)} = \tan(180^\circ - \theta) \quad \text{and} \quad \frac{\sin \theta}{-\cos \theta} = -\tan \theta$$

$$\therefore \tan(180^\circ - \theta) = -\tan \theta$$



To find sine and cosine of angle lying between  $90^\circ$  and  $180^\circ$ , proceed as follows:

- Measure the angle between the terminal side of the given angle and the horizontal axis. This is the reference angle.
- Determine the values of the cosine and sine of the reference angle.
- Give the cosine the same sign as the  $x$ -values in the quadrant of the original angle.
- Give the sine the same sign as the  $y$ -values in the quadrant of the original angle.

**Example:**

Find the exact values of  $\sin 150^\circ$ ,  $\cos 150^\circ$ , and  $\tan 150^\circ$ .


**Solution:**

Using above relations, we have:

$$\sin 150^\circ = \sin(180^\circ - 150^\circ) = \sin 30^\circ = \frac{1}{2}$$

$$\cos 150^\circ = -\cos(180^\circ - 150^\circ) = -\cos 30^\circ = -\frac{\sqrt{3}}{2}$$

$$\tan 150^\circ = -\tan(180^\circ - 150^\circ) = -\tan 30^\circ = -\frac{1}{\sqrt{3}}$$



**Key Fact:**

- Angles have cosines and sines with the same absolute value as their reference angles.
- The sign (positive or negative) can be determined from the quadrant of the angle.


### Exercise 8.1

- Find the reference angles of the following angles.
  - $125^\circ$
  - $138^\circ$
  - $111^\circ$
  - $142^\circ$
- If  $\theta = 36^\circ$ . Find equivalent angle  $\alpha$  in the second quadrant for which  $\sin \alpha = \sin \theta$ .
- If  $\theta = 87^\circ$ . Find equivalent angle  $\beta$  in the second quadrant for which  $\cos \beta = -\cos \theta$ .
- Using a reference angle, find the exact values of  $\sin 120^\circ$ ,  $\cos 120^\circ$  and  $\tan 120^\circ$ .
- Using the reference angle, find  $\cos \frac{3\pi}{4}$ ,  $\sin \frac{3\pi}{4}$  and  $\tan \frac{3\pi}{4}$ .
- If  $\cos \theta = 0.559$ , find the values of:
  - $\cos(180^\circ - \theta)$
  - $\sin(180^\circ - \theta)$
  - $\tan(180^\circ - \theta)$
  - $\cot(180^\circ - \theta)$
  - $\sec(180^\circ - \theta)$
  - $\cosec(180^\circ - \theta)$

### Solution of Right Angled Triangles

We have studied about the solution of right triangles in the previous grade. We know that any triangle (in particular a right triangle) has six quantities, three sides and three angles. In the adjoining figure, a right triangle ABC contains three sides  $a$ ,  $b$ ,  $c$  and three angles  $\alpha$ ,  $\beta$ ,  $\gamma$ . We can solve a right-angled triangle when its:

- Two sides are given,
- One acute angle and one side are given.



Let's solve some examples to discuss the two cases.

### Case 1: When Measures of two Sides are Given

**Example:** Solve the right triangle ABC in which  $\gamma = 90^\circ$ ,  $a = 5\text{cm}$ ,  $c = 13\text{cm}$ .

**Solution:** Given: A right triangle ABC in which  $\gamma = 90^\circ$ ,  $a = 5\text{cm}$ ,  $c = 13\text{cm}$ .

To Find:  $b$ ,  $\alpha$  and  $\beta$ .

Using Pythagoras theorem, we have:

$$a^2 + b^2 = c^2$$

Substituting values of  $a$  and  $c$ , we get:

$$5^2 + b^2 = 13^2 \Rightarrow 25 + b^2 = 169$$

$$\Rightarrow b^2 = 169 - 25 = 144$$

$$\Rightarrow b = 12\text{cm}$$

$$\text{Now } \cos \beta = \frac{5}{13} = 0.385$$

$$\Rightarrow \beta = \cos^{-1}(0.385) = 67.4^\circ$$

$$\text{As } \alpha + \beta = 90^\circ$$

$$\Rightarrow \alpha = 90^\circ - \beta = 90^\circ - 67.4^\circ = 22.6^\circ$$

### Case 2: When Measures of one Side and one Angle are Given

**Example:** Solve the right triangle ABC in which  $\gamma = 90^\circ$ ,  $\beta = 45^\circ$ ,  $a = 10\text{cm}$ .

**Solution:** Given: A right triangle ABC in which  $\gamma = 90^\circ$ ,  $\beta = 45^\circ$ ,  $a = 10\text{cm}$ .

To Find:  $b$ ,  $c$  and  $\alpha$ .

$$\text{As } \alpha + \beta = 90^\circ$$

$$\text{As } \alpha + \beta = 90^\circ$$

$$\Rightarrow \alpha = 90^\circ - \beta = 90^\circ - 45^\circ = 45^\circ$$

$$\text{Now } \frac{b}{a} = \tan 45^\circ$$

$$\Rightarrow b = a \times \tan 45^\circ = 10 \times 1 = 10\text{cm}$$

To find  $c$ , we use Pythagoras theorem as follows:

$$c^2 = a^2 + b^2$$

Substituting values of  $a$  and  $b$ , we get:



$$c^2 = 10^2 + 10^2 = 100 + 100 = 200$$

$$\Rightarrow c = 10\sqrt{2}\text{cm}$$

**Example:** From a small plane Z the angle of depression of a boat is  $25^\circ$ . The angle of depression of a boat on the other side of the plane is  $50^\circ$ . The plane is flying at an altitude of 1500 m. How far apart are the boats?

**Solution:** In the figure,

distance between the boats = XY = XU + UY



Using the properties of angles,

$$\angle X = 25^\circ \text{ and } \angle Y = 50^\circ$$

In right  $\triangle XUZ$ ,

$$\tan 25^\circ = \frac{ZU}{XU}$$

$$XU = \frac{ZU}{\tan 25^\circ} = \frac{1500}{0.466} = 3216.76m$$

In right  $\triangle YUZ$ ,

$$\tan 50^\circ = \frac{ZU}{YU}$$

$$YU = \frac{ZU}{\tan 50^\circ} = \frac{1500}{1.192} = 1258.65m$$

$$\therefore \text{Distance between the boats} = XY = XU + UY \\ = 3216.76 + 1258.65 = 4475.41 \text{ m}$$

### Example:

Find the measure of  $\angle F$  in the adjoining figure.

**Solution:**

To find  $\angle F$ , we need measure of  $\overline{GE}$ .

In right  $\triangle AEG$ ,



$$\sin 34^\circ = \frac{GE}{GD}$$

$$GE = GD \times \sin 34^\circ = 14.4 \times 0.559 = 8.05 \text{ cm}$$

Now in right  $\triangle EGF$ ,

$$\sin(\angle F) = \frac{GE}{GF} = \frac{8.05}{9.6} = 0.839$$

$$\angle F = \sin^{-1}(0.839) = 57^\circ 2'$$




### Check Point:


Abbas is flying a kite to which the angle of elevation is  $70^\circ$ . The string of the kite is 65 meters long. How far is the kite above the ground?

### Exercise 8.2


1. Solve the following right-angled triangles.



(i)




(ii)




(iii)

2. Solve right-angled triangles ABC in which  $\gamma = 90^\circ$  and
- $a = 12\text{cm}$ ,  $\beta = 35^\circ$
  - $b = 30\text{cm}$ ,  $\alpha = 25^\circ 35'$
  - $a = 50\text{cm}$ ,  $b = 25\text{cm}$
  - $a = 30\text{cm}$ ,  $c = 40\text{cm}$
  - $a = 24\text{cm}$ ,  $\alpha = 36^\circ 15'$
  - $b = 10\text{cm}$ ,  $\alpha = 70.5^\circ$
3. A tower casts a shadow that is 20 m long when the angle of elevation of the sun is  $65^\circ$ . How tall is the tower?
4. Arif is standing on top of a cliff 105 m above a lake. The measurement of the angle of depression to a boat on the lake is  $42^\circ$ . How far is the boat from Arif?
5. A ladder that is 20 ft. long is leaning against the side of a building. If the angle formed between the ladder and the ground is  $75^\circ$ , how far is the bottom of the ladder from the base of the building?
6. Uzma is standing 50 meters from a hot air balloon that is preparing to take off. The angle of elevation to the top of the balloon is  $28^\circ$ . Find the height of the balloon.
7. A man is in a boat that is floating 175 feet from the base of a 200-foot cliff. What is the angle of depression between the cliff and the boat?
8. A flagpole casts a shadow 40 feet long when the measurement of the angle of elevation to the sun is  $31^\circ$ . How tall is the flagpole?
9. A straight waterslide is 175 feet above ground and is 200 feet long. What is the angle of depression to the bottom of the slide?
10. Zain wants to measure the height of a tree. He walks exactly 50 m from the base of the tree and looks up that the angle from the ground to the top of the tree is  $33^\circ$ . How tall is the tree?
11. From a 100m observation tower on the beach, a man sights a whale in difficulty. The angle of depression of the whale is  $7^\circ$ . How far is the whale from the shoreline?
12. Urooba is sitting on the ground midway between two trees, 100 m apart. The angles of elevation of the tops of the trees are  $13^\circ$  and  $18^\circ$ . How much taller is one tree than the other?
13. The angle of elevation of the top of a tree, T is  $27^\circ$ . From the same point on the ground, the angle of elevation of a hawk H, flying directly above the tree is  $43^\circ$ . The tree is 12.7m tall. How high is the hawk above the ground?
14. This diagram shows a falcon F, on a tree, with a squirrel S, and a chipmunk C, on the ground. From the falcon, the angles of depression of the animals are  $36^\circ$  and  $47^\circ$ . How far apart are the animals on the ground?
15. Two guy wires support a flagpole FH. The first wire is 11.2 m long and has an angle of elevation of  $39^\circ$ . The second wire has an angle of elevation of  $47^\circ$ . How tall is the flagpole?




## Oblique Triangles

An oblique triangle is any triangle that is not a right triangle. It could be an acute triangle or it could be an obtuse triangle.



Acute Triangle



Obtuse Triangle

The Pythagoras Theorem along with trigonometric ratios allow us to easily handle any given right triangle problem, but what if the triangle is not a right triangle i.e. the triangle is oblique?

To solve oblique triangles, we use certain laws known as "law of cosines", "law of sines" and "law of tangents".

### Law of Cosines

When we know the measures of two sides and the included angle of a triangle (SAS), the size and shape of the triangle are determined. Therefore, we should be able to find the measure of the third side of the triangle. Sometimes three sides of oblique triangles are given (SSS) and we have to find the three angles of triangle. Such cases are easily handled by using law of cosines.

#### Statement:

If  $a, b, c$  are three sides and  $\alpha, \beta, \gamma$  are three angles of an oblique triangle with usual notation, then law of cosines is stated as follows:

$$a^2 = b^2 + c^2 - 2bc \cos \alpha \dots\dots (1)$$

$$b^2 = a^2 + c^2 - 2ac \cos \beta \dots\dots (2)$$

$$c^2 = a^2 + b^2 - 2ab \cos \gamma \dots\dots (3)$$

**Proof:** Consider an acute angled triangle ABC with usual notations as shown in the figure.

Draw  $\overline{AD} \perp \overline{BC}$ .

In right  $\triangle ABD$ ,

$$\frac{BD}{AB} = \cos \beta$$


$$\Rightarrow BD = AB \cos \beta = c \cos \beta \dots\dots (i)$$

$$\text{Again, } \frac{AD}{AB} = \sin \beta$$

$$\Rightarrow AD = AB \sin \beta = c \sin \beta \dots\dots (ii)$$

$$\text{Now, } CD = BC - BD = a - c \cos \beta$$

In right  $\triangle ACD$ ,



$$(AC)^2 = (AD)^2 + (CD)^2 \quad \dots \text{(Pythagoras Theorem)}$$

$$b^2 = (c \sin \beta)^2 + (a - c \cos \beta)^2$$

$$b^2 = c^2 \sin^2 \beta + a^2 + c^2 \cos^2 \beta - 2ac \cos \beta$$

$$b^2 = a^2 + c^2 (\sin^2 \beta + \cos^2 \beta) - 2ac \cos \beta$$

$$b^2 = a^2 + c^2 - 2ac \cos \beta$$

Similarly, we can prove that:

$$a^2 = b^2 + c^2 - 2bc \cos \alpha \quad \text{and} \quad c^2 = a^2 + b^2 - 2ab \cos \gamma$$

From laws of cosines,

$$\cos \alpha = \frac{b^2 + c^2 - a^2}{2bc}, \quad \cos \beta = \frac{a^2 + c^2 - b^2}{2ac}, \quad \cos \gamma = \frac{a^2 + b^2 - c^2}{2ab}$$

These laws are useful when three sides of oblique triangles are given and we have to find their angles.

#### Check Point:

Prove law of cosines by taking an obtuse triangle.

**Example:** Use the law of cosines to solve the triangle ABC when:

$$a = 12\text{cm}, b = 7\text{cm}, \gamma = 59^\circ 30'$$

**Solution:** Using law of cosines, we have:

$$c^2 = a^2 + b^2 - 2ab \cos \gamma$$

$$c^2 = 12^2 + 7^2 - 2 \times 12 \times 7 \times \cos 59^\circ 30'$$

$$c^2 = 144 + 49 - 168 \times 0.506 = 108$$

$$c = \sqrt{108} = 10.4\text{cm}$$

Now to find angle  $\alpha$ , we use following formula.

$$\begin{aligned} \cos \alpha &= \frac{b^2 + c^2 - a^2}{2bc} = \frac{7^2 + 10.4^2 - 12^2}{2 \times 7 \times 10.4} \\ &= \frac{13.16}{145.6} = 0.0904 \end{aligned}$$

$$\alpha = \cos^{-1}(0.0904) = 84^\circ 49'$$

$$\text{Finally, } \alpha + \beta + \gamma = 180^\circ$$

$$\beta = 180^\circ - \alpha - \gamma = 180^\circ - 84^\circ 49' - 59^\circ 30'$$


$$\beta = 35^\circ 41'$$



#### Key Fact:

Law of cosines is applicable when:

- ❖ Two sides and included angle are given.
- ❖ Three sides are given.



#### Key Fact:

If in the law of cosines,

$$a^2 = b^2 + c^2 - 2bc \cos \alpha$$

angle  $\alpha$  is  $90^\circ$ , then

$$a^2 = b^2 + c^2 - 2bc \cos 90^\circ$$

$$a^2 = b^2 + c^2 - 2bc \times 0 \Rightarrow a^2 = b^2 + c^2$$

which becomes Pythagoras Theorem.

Thus Pythagoras Theorem is a special case of "law of cosines".

#### Check Point:

Two forces of magnitude 20N and 30N are inclined at angle of  $105^\circ$  with each other. Find the magnitude of resultant force. Can you find the angle between force of magnitude 20N and resultant force?

## Law of Sines

Law of cosines is not applicable when two angles and one side (AAS or ASA) or one angle and two sides (angle not included (SSA)) of an oblique triangle are given. In order to handle such situation, we use another law known as “law of sines”.

### Statement:

If  $a, b, c$  are three sides and  $\alpha, \beta, \gamma$  are three angles of an oblique triangle with usual notation, then law of sines is stated as follows:

$$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma}$$

**Proof:** Consider an obtuse triangle ABC as shown in the figure below.

Let  $h$  be the height of triangle ABC with respect to base  $\overline{AC}$ . In right  $\Delta ABD$ ,

$$\frac{BD}{AB} = \sin \alpha$$

$$\Rightarrow BD = AB \sin \alpha = c \sin \alpha$$


$$\Rightarrow h = c \sin \alpha \dots\dots(i)$$

In right  $\Delta CBD$ ,

$$\frac{BD}{CB} = \sin(\pi - \gamma)$$

$$\Rightarrow BD = CB \sin(\pi - \gamma) \quad [\text{as } \sin(\pi - \gamma) = \sin \gamma]$$

$$\Rightarrow h = a \sin \gamma \dots\dots(ii)$$



From (i) and (ii), we get:

$$a \sin \gamma = c \sin \alpha$$

$$\text{Or } \frac{a}{\sin \alpha} = \frac{c}{\sin \gamma}$$

Similarly, by drawing perpendiculars on other two sides of  $\Delta ABC$ , we get:

$$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} \quad \text{and} \quad \frac{c}{\sin \gamma} = \frac{b}{\sin \beta}$$

Combining above results, we obtain the following law known as “law of sines”.

$$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma} \quad \text{or} \quad \frac{\sin \alpha}{a} = \frac{\sin \beta}{b} = \frac{\sin \gamma}{c}$$

### To Determine the Number of Solutions in $\Delta ABC$

**Task:** Given measure of sides  $a, b$  and angle  $\alpha$ . Use the law of sines to solve the  $\Delta ABC$  for  $\sin \beta$ .

#### Case-1:

- (i) When  $\sin \beta > 1$ , there is no triangle, so no solution.

#### Check Point:

Prove law of sines by taking an acute triangle.

- (ii) When  $\sin \beta = 1$ , there is one right triangle (one solution) if  $\alpha$  is acute and no triangle (no solution) if  $\alpha$  is obtuse.

**Case-2:**

When  $\alpha$  is acute and  $\sin \beta < 1$ , there are two possible values of  $\beta$ , acute or obtuse.

$$0 < \beta < 90^\circ \text{ and } \beta' = 180^\circ - \beta$$

- (i) If  $\alpha + \beta' < 180^\circ$ , there are two possible triangles  $\Delta ABC$  and  $\Delta AB'C$ .  
(ii) If  $\alpha + \beta' \geq 180^\circ$ , there is only one  $\Delta ABC$ .

**Case-3:**

When  $\alpha$  is obtuse and  $\sin \beta < 1$ , then  $\beta$  must be acute.

- (i) If  $\alpha + \beta < 180^\circ$ , there is only one  $\Delta ABC$ .  
(ii) If  $\alpha + \beta \geq 180^\circ$ , there is no triangle, so no solution.

Alternatively, if we let  $h = b \sin \alpha$ , the height of triangle, we can summarize the number of possible triangles given  $a$ ,  $b$  and  $\alpha$  in  $\Delta ABC$ .

Angle is $\alpha$	Acute $a < h$	Acute $a = h$	Acute $h < a < b$	Acute $a > b$	Obtuse $a < h$	Obtuse $a > b$
Possible Triangles	none	one right triangle	two triangles	one triangle	none	one triangle

### Example:

Use the law of sines to solve the triangle ABC when:

$$a = 25\text{cm}, \alpha = 66^\circ 51', \gamma = 44^\circ 12'$$

### Solution:

$$\text{Given: } a = 25\text{cm}, \alpha = 66^\circ 51', \gamma = 44^\circ 12'$$

To Find:  $b$ ,  $c$  and  $\beta$

$$\alpha + \beta + \gamma = 180^\circ$$

$$66^\circ 51' + \beta + 44^\circ 12' = 180^\circ$$

$$\beta = 180^\circ - 66^\circ 51' - 44^\circ 12' = 68^\circ 57'$$

Using law of sines, we have:

$$\frac{a}{\sin \alpha} = \frac{c}{\sin \gamma} \Rightarrow \frac{25}{\sin 66^\circ 51'} = \frac{c}{\sin 44^\circ 12'}$$

$$c = \frac{25 \times \sin 44^\circ 12'}{\sin 66^\circ 51'} = 18.96\text{cm}$$

Again, by law of sines,

### Check Points

Solve the triangle ABC by using both law of sines and law of cosines when:

$$a = 25\text{cm}, \alpha = 66^\circ 51', \gamma = 44^\circ 12'$$



$$\frac{b}{\sin \beta} = \frac{a}{\sin \alpha} \Rightarrow \frac{b}{\sin 68^\circ 57'} = \frac{25}{\sin 66^\circ 51'}$$

$$\Rightarrow b = \frac{25}{\sin 66^\circ 51'} \times \sin 68^\circ 57' = 25.37 \text{ cm}$$

**Example:**

Find the values of unknowns in the adjoining figure.

Figure is not drawn according to scale.

**Solution:**

In  $\Delta ABC$ ,

$$\angle ACB = 180^\circ - 70^\circ = 110^\circ$$

$$\text{and } \angle ABC = 180^\circ - 32^\circ - 110^\circ = 38^\circ$$

Using law of sines for triangle ABC,

$$\frac{a}{\sin 32^\circ} = \frac{900}{\sin 38^\circ}$$

$$\Rightarrow a = \frac{900}{\sin 38^\circ} \times \sin 32^\circ = 774.66 \text{ m}$$

Again for the  $\Delta ABC$ ,

$$\frac{c}{\sin 110^\circ} = \frac{774.66}{\sin 32^\circ}$$


$$c = \frac{774.66}{\sin 32^\circ} \times \sin 110^\circ = 1373.69 \text{ m}$$

Now in  $\Delta BCD$ ,

$$\frac{a}{\sin \angle D} = \frac{1200}{\sin 70^\circ} \Rightarrow \frac{774.66}{\sin \angle D} = \frac{1200}{\sin 70^\circ}$$

$$\sin \angle D = \frac{774.66 \times \sin 70^\circ}{1200} = 0.607$$

$$\angle D = \sin^{-1}(0.607) = 37^\circ 21'$$



## Law of Tangents

Law of tangents is also used to solve oblique triangles. This law handles the same situations for which the law of cosines is used.

**Statement:**

If  $a, b, c$  are three sides and  $\alpha, \beta, \gamma$  are three angles of an oblique triangle with usual notation, then law of tangents is stated as follows:

$$\frac{a+b}{a-b} \cdot \frac{\tan\left(\frac{\alpha+\beta}{2}\right)}{\tan\left(\frac{\alpha-\beta}{2}\right)}, \quad \frac{b+c}{b-c} = \frac{\tan\left(\frac{\beta+\gamma}{2}\right)}{\tan\left(\frac{\beta-\gamma}{2}\right)}, \quad \frac{c+a}{c-a} = \frac{\tan\left(\frac{\gamma+\alpha}{2}\right)}{\tan\left(\frac{\gamma-\alpha}{2}\right)}$$

**Proof:**

For any oblique triangle ABC, the law of sines is given as follows:

$$\frac{a}{\sin \alpha} = \frac{b}{\sin \beta}$$

$$\frac{a}{b} = \frac{\sin \alpha}{\sin \beta}$$

Using componendo and dividendo property, we get:

$$\frac{a+b}{a-b} = \frac{\sin \alpha + \sin \beta}{\sin \alpha - \sin \beta}$$

$$\frac{a+b}{a-b} = \frac{2 \sin \frac{\alpha+\beta}{2} \cos \frac{\alpha-\beta}{2}}{2 \cos \frac{\alpha+\beta}{2} \sin \frac{\alpha-\beta}{2}}$$

$$\frac{a+b}{a-b} = \frac{\tan \frac{\alpha+\beta}{2}}{\tan \frac{\alpha-\beta}{2}}$$

Similarly, we can prove that:

$$\frac{b+c}{b-c} = \frac{\tan \left( \frac{\beta+\gamma}{2} \right)}{\tan \left( \frac{\beta-\gamma}{2} \right)}, \quad \frac{c+a}{c-a} = \frac{\tan \left( \frac{\gamma+\alpha}{2} \right)}{\tan \left( \frac{\gamma-\alpha}{2} \right)}$$

**Example:**

Use the law of tangents to solve the triangle ABC when  $a = 500\text{mm}$ ,  $b = 600\text{mm}$ ,  $\gamma = 50^\circ 20'$

**Solution:**

Given:  $a = 500\text{cm}$ ,  $b = 600\text{cm}$ ,  $\gamma = 50^\circ 20'$

To Find:  $c, \alpha$  and  $\beta$

As in any triangle,


$$\alpha + \beta + \gamma = 180^\circ \Rightarrow \beta + \alpha = 180^\circ - \gamma$$

$$\Rightarrow \beta + \alpha = 180^\circ - 50^\circ 20' = 129^\circ 40' \dots\dots (i)$$

Now using law of tangents for  $b > a$ ,

$$\frac{b+a}{b-a} = \frac{\tan \frac{\beta+\alpha}{2}}{\tan \frac{\beta-\alpha}{2}}$$

Substituting the values of  $b$ ,  $c$  and  $\beta + \alpha$ , we get:


**Check Point:**

Prove that

$$\frac{a+c}{a-c} = \frac{\tan \left( \frac{\alpha+\gamma}{2} \right)}{\tan \left( \frac{\alpha-\gamma}{2} \right)}$$

**Key Fact:**

It is better to use formula

$$\frac{a+c}{a-c} = \frac{\tan \left( \frac{\alpha+\gamma}{2} \right)}{\tan \left( \frac{\alpha-\gamma}{2} \right)}$$

rather than the formula

$$\frac{c+a}{c-a} = \frac{\tan \left( \frac{\gamma+\alpha}{2} \right)}{\tan \left( \frac{\gamma-\alpha}{2} \right)}$$

when  $a > c$  in the given situation.

$$\frac{600+500}{600-500} = \frac{\tan \frac{129^\circ 40'}{2}}{\tan \frac{\beta-\alpha}{2}} \Rightarrow \frac{1100}{100} = \frac{\tan 64^\circ 50'}{\tan \frac{\beta-\alpha}{2}}$$

$$\tan \frac{\beta-\alpha}{2} = \frac{\tan 64^\circ 50'}{11} \Rightarrow \tan \frac{\beta-\alpha}{2} = 0.193$$

$$\frac{\beta-\alpha}{2} = \tan^{-1}(0.193) = 10^\circ 57'$$

$$\beta-\alpha = 21^\circ 54' \dots\dots (ii)$$

Solving (i) and (ii), we get:

$$\beta = 75^\circ 47', \alpha = 53^\circ 53'$$

To find  $c$ , we use law of sines (we can also use law of cosines).


$$\frac{c}{\sin \gamma} = \frac{b}{\sin \beta} \Rightarrow c = \frac{b \times \sin \gamma}{\sin \beta}$$

$$c = \frac{600 \times \sin 50^\circ 20'}{\sin 75^\circ 47'} = 476.45 \text{ mm}$$

### When Law of Sines and Law of Cosines are Used


The Law of Cosines and the Law of Sines can be used to find the remaining three measures of any triangle when we know the measure of a side and the measures of any two other components (two sides, two angles, one side one angle) of the triangle.

**1. Given:** Two sides and the included angle.



- Use the Law of Cosines to find the measure of the third side.
- Use the Law of Sines or the Law of Cosines to find the measure of another angle.
- Use the sum of the angles of a triangle to find the measure of the third angle.

**2. Given:** Three sides.



- Use the Law of Cosines to find the measure of an angle.
- Use the Law of Sines or the Law of Cosines to find the measure of another angle.
- Use the sum of the angles of a triangle to find the measure of the third angle.

### Check Point;

Solve the triangle ABC by using both law of tangents and law of sines when:


$$b = 50 \text{ cm}, c = 40, \alpha = 80^\circ 30'$$

3. Given: Two angles and a side.



- Use the sum of the angles of a triangle to find the measure of the third angle.
- Use the Law of Sines to find the remaining sides.

4. Given: Two sides and an angle opposite to one of them



- Use the Law of Sines to find the possible measure(s) of another angle.
- Determine if there are two, one, or no possible triangles.
- If there is a triangle, use the sum of the angles of a triangle to find the measure(s) of the third angle.
- Use the Law of Sines or the Law of Cosines to find the measure(s) of the third side.

## Half Angle Formulae

Sometimes it is convenient to use half angle formulae for the solution of oblique triangles when measures of three sides of a triangle are given. In this section, we will study half angles formulae for sines, cosines and tangents. Let us derive them one by one.

### (a) Cosines of Half Angle

**Statement:** In any triangle ABC with usual notations,

$$\cos \frac{\alpha}{2} = \sqrt{\frac{S(S-a)}{bc}}, \quad \cos \frac{\beta}{2} = \sqrt{\frac{S(S-b)}{ac}}, \quad \cos \frac{\gamma}{2} = \sqrt{\frac{S(S-c)}{ab}}$$

$$\text{where } S = \frac{a+b+c}{2}.$$

**Proof:** We know that:

$$\cos \alpha = \frac{b^2 + c^2 - a^2}{2bc}$$

$$2\cos^2 \frac{\alpha}{2} - 1 = \frac{b^2 + c^2 - a^2}{2bc} \Rightarrow 2\cos^2 \frac{\alpha}{2} = \frac{b^2 + c^2 - a^2}{2bc} + 1$$

$$\Rightarrow 2\cos^2 \frac{\alpha}{2} = \frac{b^2 + c^2 + 2bc - a^2}{2bc} = \frac{(b+c)^2 - (a)^2}{2bc}$$

$$\Rightarrow 2\cos^2 \frac{\alpha}{2} = \frac{(b+c+a)(b+c-a)}{2bc}$$

$$\text{As } a+b+c=2S \Rightarrow b+c-a=2S-2a=2(S-a)$$


$$\therefore 2\cos^2 \frac{\alpha}{2} = \frac{2S \times 2(S-a)}{2bc}$$

$$\cos^2 \frac{\alpha}{2} = \frac{S(S-a)}{bc}$$

$$\cos \frac{\alpha}{2} = \sqrt{\frac{S(S-a)}{bc}}$$

### Check Point:

Find the diameter of the circle shown in the figure below.



Similarly, it can be proved that:

$$\cos \frac{\beta}{2} = \sqrt{\frac{S(S-b)}{ac}} \text{ and } \cos \frac{\gamma}{2} = \sqrt{\frac{S(S-c)}{ab}}$$

### (b) Sines of Half Angle

**Statement:** In any triangle ABC with usual notations,

$$\sin \frac{\alpha}{2} = \sqrt{\frac{(S-b)(S-c)}{bc}}, \quad \sin \frac{\beta}{2} = \sqrt{\frac{(S-a)(S-c)}{ac}}, \quad \sin \frac{\gamma}{2} = \sqrt{\frac{(S-a)(S-b)}{ab}}$$

$$\text{where } S = \frac{a+b+c}{2}.$$

**Proof:** We know that

$$\cos \alpha = \frac{b^2 + c^2 - a^2}{2bc}$$

$$1 - 2\sin^2 \frac{\alpha}{2} = \frac{b^2 + c^2 - a^2}{2bc} \Rightarrow 2\sin^2 \frac{\alpha}{2} = 1 - \frac{b^2 + c^2 - a^2}{2bc}$$

$$\Rightarrow 2\sin^2 \frac{\alpha}{2} = \frac{(a)^2 - (b-c)^2}{2bc} = \frac{(a-b+c)(a+b-c)}{2bc}$$

$$\text{As } a+b+c=2S \Rightarrow a+c-b=2S-2b=2(S-b) \text{ and } a+b-c=2S-2c=2(S-c)$$

$$\therefore 2\sin^2 \frac{\alpha}{2} = \frac{2(S-b) \times 2(S-c)}{2bc} = \frac{2(S-b)(S-c)}{bc}$$

$$\sin^2 \frac{\alpha}{2} = \frac{(S-b)(S-c)}{bc}$$

$$\sin \frac{\alpha}{2} = \sqrt{\frac{(S-b)(S-c)}{bc}}$$

### Check Point:

Prove that

$$\cos \frac{\gamma}{2} = \sqrt{\frac{S(S-c)}{ab}} \text{ and } \sin \frac{\gamma}{2} = \sqrt{\frac{(S-a)(S-b)}{ab}}$$

Similarly, it can be proved that:

$$\sin \frac{\beta}{2} = \sqrt{\frac{(S-a)(S-c)}{ac}} \text{ and } \sin \frac{\gamma}{2} = \sqrt{\frac{(S-a)(S-b)}{ab}}$$

### (c) Tangents of Half Angle

**Statement:** In any triangle ABC with usual notations,

$$\tan \frac{\alpha}{2} = \sqrt{\frac{(S-b)(S-c)}{S(S-a)}}, \quad \tan \frac{\beta}{2} = \sqrt{\frac{(S-a)(S-c)}{S(S-b)}}, \quad \tan \frac{\gamma}{2} = \sqrt{\frac{(S-a)(S-b)}{S(S-c)}}$$

where  $S = \frac{a+b+c}{2}$ .

$$\text{Proof: } \tan \frac{\alpha}{2} = \frac{\sin \frac{\alpha}{2}}{\cos \frac{\alpha}{2}} = \frac{\sqrt{\frac{(S-b)(S-c)}{bc}}}{\sqrt{\frac{S(S-a)}{bc}}} = \sqrt{\frac{(S-b)(S-c)}{S(S-a)}}$$

$$\therefore \tan \frac{\alpha}{2} = \sqrt{\frac{(S-b)(S-c)}{S(S-a)}}$$

Similarly, it can be proved that:

$$\tan \frac{\beta}{2} = \sqrt{\frac{(S-a)(S-c)}{S(S-b)}}, \quad \tan \frac{\gamma}{2} = \sqrt{\frac{(S-a)(S-b)}{S(S-c)}}$$

### Example:

Use half angle formulae to solve the triangle ABC when  $a = 5\text{cm}$ ,  $b = 6\text{cm}$ ,  $c = 7\text{cm}$ .

### Solution:

Given:  $a = 5\text{cm}$ ,  $b = 6\text{cm}$ ,  $c = 7\text{cm}$ .

To Find:  $\alpha$ ,  $\beta$  and  $\gamma$

We use half angle formula for cosines here.

$$\text{Now } S = \frac{a+b+c}{2} = \frac{5+6+7}{2} = 9$$

$$\therefore \cos \frac{\alpha}{2} = \sqrt{\frac{S(S-a)}{bc}} = \sqrt{\frac{9(9-5)}{6 \times 7}} = \sqrt{\frac{36}{42}} = 0.926$$

$$\frac{\alpha}{2} = \cos^{-1}(0.926) = 22.18^\circ \Rightarrow \alpha = 44.36^\circ$$

$$\begin{aligned} \text{Now } \cos \frac{\beta}{2} &= \sqrt{\frac{S(S-b)}{ac}} = \sqrt{\frac{9(9-6)}{5 \times 7}} \\ &= \sqrt{\frac{27}{35}} = 0.878 \end{aligned}$$

$$\frac{\beta}{2} = \cos^{-1}(0.878) = 28.58^\circ \Rightarrow \beta = 57.20^\circ$$

To find third angle, we have:

$$\alpha + \beta + \gamma = 180^\circ \Rightarrow \gamma = 180^\circ - \alpha - \beta$$

$$\Rightarrow \gamma = 180^\circ - 44.36^\circ - 57.20^\circ = 78.44^\circ$$

### Check Point:

Solve the triangle ABC when  $a = 5\text{cm}$ ,  $b = 6\text{cm}$ ,  $c = 7\text{cm}$  using half angle formulae for sines and tangents. Which one method do you think easy?


### Check Point:

The base of an isosceles triangle measures  $14.5\text{ cm}$  and the vertex angle measures  $110^\circ$ .



- Find the measure of one of the congruent sides of the triangle to the nearest hundredth.
- Find the perimeter of the triangle.

### Exercise 8.3



1. Use the Law of Cosines to find the remaining side(s) and angle(s) in triangle ABC.
  - (i)  $a = 7, b = 12, \gamma = 58.5^\circ$
  - (ii)  $b = 20, c = 15, \alpha = 36^\circ$
  - (iii)  $a = 150, c = 150, \beta = 15^\circ$
  - (iv)  $\beta = 45^\circ, a = 30, c = 42$
  - (v)  $a = 7, b = 10, c = 12$
  - (vi)  $a = 3, b = 3, c = 5$
  - (vii)  $a = 30, b = 45, c = 50$
  - (viii)  $a = 6, b = 6, c = 6$
  - (ix)  $a = 5, b = 12, c = 13$
2. Use the half angles formulae to solve parts (v) to (ix) in Q.1 where possible.
3. Use the Law of Sines to solve triangle ABC in the following (where possible).
  - (i)  $a = 10, \alpha = 40^\circ, \beta = 60^\circ$
  - (ii)  $b = 20, \alpha = 50^\circ, \beta = 70^\circ$
  - (iii)  $c = 12, \alpha = 45^\circ, \beta = 75^\circ$
  - (iv)  $b = 18, \beta = 40^\circ 35', \gamma = 120^\circ$
  - (v)  $a = 14.6, \alpha = 25^\circ 10', \beta = 85.5^\circ$
  - (vi)  $c = 52, \alpha = 42.3^\circ, \gamma = 85^\circ 14'$
4. Solve parts (i) to (iv) of Q.1 by using the Law of Tangents.
5. Solve by using appropriate law.
  - (i)  $a = 10, b = 8, \beta = 80^\circ$
  - (ii)  $b = 20, c = 14, \beta = 70^\circ$
  - (iii)  $c = 12, b = 10, \gamma = 64^\circ$
  - (iv)  $b = 18, \alpha = 55^\circ 5', a = 37$
  - (v)  $a = 14.6, b = 10.6, c = 17.2$
  - (vi)  $c = 88, \beta = 23.2^\circ, \gamma = 73^\circ 14'$
6. A pilot is flying from city A to city C, 500 km apart. He starts his flight  $20^\circ$  off course and flies on this course for 150 km and is above city B. How far is he from city C?




7. Two sides of a triangular plot have lengths 400m and 600m. The measurement of angle between the sides is  $45^\circ$ . Find the perimeter and area of the plot.
8. The sides of a triangle are 6.5cm, 8.2cm and 5.8cm. Find the measurement of smallest and largest angles.
9. The sides of a parallelogram are 50cm and 70cm. Find the length of each diagonal if the larger angle measures  $110^\circ$ .
10. For parallelogram ABCD, find :
  - (i) BC
  - (ii)  $\angle BDC$




11. For the figure below find:




12. Find the height of the building in the figure below.



13. Fire towers A and B are located 10 miles apart at the same level of ground. Rangers at fire tower A spots a fire at  $42^\circ$ , and rangers at fire tower B spot the same fire at  $64^\circ$ . How far from tower A is the fire to the nearest tenth of a mile?



14. Circle O has a radius of 15cm. The angle between radii  $\overline{OA}$  and  $\overline{OB}$  is  $120^\circ$ . Find the length of chord  $\overline{AB}$ .



15. Two lighthouses are 12 miles apart along a straight shore. A ship is 15 miles from one lighthouse and 20 miles from the other. Find, to the nearest degree, the measure of the angle between the lines of sight from the ship to each lighthouse.


## Area of Triangular Region

### (a) Area of Triangular Region when Two Sides and Included Angle are Given

When the measures of two sides and the included angle of a triangle are known, the size and shape of the triangle is determined. Therefore, it is possible to use these known values to find the area of the triangle. Let us find the area of an acute angled triangle ABC given below.

We know that:

$$\text{Area } (\Delta) \text{ of triangle ABC} = \frac{1}{2} \times \text{base} \times \text{height}$$



$$\text{Area } (\Delta) \text{ of triangle ABC} = \frac{1}{2} \times AB \times CD = \frac{1}{2} \times c \times h \dots\dots \text{ (i)}$$

Now in right angled triangle ACD,  $AB = c$

$$\text{and } \frac{h}{b} = \sin \alpha \Rightarrow h = b \sin \alpha$$

Therefore from (i),

$$\Delta = \frac{1}{2} \times c \times b \sin \alpha = \frac{1}{2} b c \sin \alpha$$

Again, in right angled triangle BCD

$$\frac{h}{a} = \sin \beta \Rightarrow h = a \sin \beta$$

Therefore from (i),

$$\Delta = \frac{1}{2} \times c \times a \sin \beta = \frac{1}{2} a c \sin \beta$$

Similarly, we can prove that:

$$\Delta = \frac{1}{2}ab\sin\gamma$$

$$\text{Hence } \Delta = \frac{1}{2}bc\sin \alpha = \frac{1}{2}ac\sin \beta = \frac{1}{2}ab\sin \gamma$$

### **Example:**

Find the area of triangle DEF if  $DE = 10$ ,  $EF = 8$  and  $\angle E = 30^\circ$ .

**Solution:**

$$\begin{aligned}\text{Area of triangle DEF} &= \frac{1}{2} \times DE \times EF \times \sin(\angle E) \\ &= \frac{1}{2} \times 10 \times 8 \times \sin 30^\circ \\ &= 40 \times 0.5 = 20 \text{ sq. units}\end{aligned}$$

**(b) Area of Triangular Region when Two Angles and One Side are Given**

If  $a$  is measure of side of a triangle ABC and any two angles let's say  $\beta$  and  $\gamma$  are given then:

$$\text{Area of triangle } ABC = \Delta = \frac{1}{2} a^2 \frac{\sin \beta \sin \gamma}{\sin \alpha}$$

Let us prove this formula.

We have proved in the previous section that:

#### **Check Point**

Urwa said that the area of rhombus  $PQRS$  is  $(PQ)^2(\sin P)$ . Do you agree with Urwa? Explain why or why not.

$$\text{Area of triangle ABC} = \Delta = \frac{1}{2} ab \sin \gamma \dots\dots(i)$$

But in the current situation,  $b$  is not given. We can replace  $b$  by using law of sines as follows.

$$\frac{b}{\sin \beta} = \frac{a}{\sin \alpha} \Rightarrow b = \frac{a \sin \beta}{\sin \alpha}$$

Substituting the value of  $b$  in (i), we get:

$$\Delta = \frac{1}{2} a \left( \frac{a \sin \beta}{\sin \alpha} \right) \sin \gamma = \frac{1}{2} a^2 \frac{\sin \beta \sin \gamma}{\sin \alpha}$$

In the same way, we can prove that:

$$\Delta = \frac{1}{2} b^2 \frac{\sin \alpha \sin \gamma}{\sin \beta} \quad \text{and} \quad \Delta = \frac{1}{2} c^2 \frac{\sin \alpha \sin \beta}{\sin \gamma}$$

$$\text{Hence, } \Delta = \frac{1}{2} a^2 \frac{\sin \beta \sin \gamma}{\sin \alpha}, \quad \Delta = \frac{1}{2} b^2 \frac{\sin \alpha \sin \gamma}{\sin \beta}, \quad \Delta = \frac{1}{2} c^2 \frac{\sin \alpha \sin \beta}{\sin \gamma}$$

### Example:

Calculate the cost of grass cutting @ Rs.10 per sq. unit of a triangular plot ABC if  $a=16.5$ ,  $\alpha=40.5^\circ$ ,  $\beta=65^\circ$ .

**Solution:** To find the cost of grass cutting, we need area of plot and to find area, we need the third angle  $\gamma$ .

$$\begin{aligned}\gamma &= 180^\circ - \alpha - \beta \\ &\Rightarrow \gamma = 180^\circ - 40.5^\circ - 65^\circ = 74.5^\circ\end{aligned}$$

$$\begin{aligned}\text{Now, area of triangle ABC} &= \frac{1}{2} a^2 \frac{\sin \beta \sin \gamma}{\sin \alpha} \\ &= \frac{1}{2} \times (16.5)^2 \times \frac{\sin 65^\circ \times \sin 74.5^\circ}{\sin 40.5^\circ} \\ &= 183.05 \text{ sq. units}\end{aligned}$$

Therefore, the cost of grass cutting @ Rs.10 per sq. unit = Rs.  $10 \times 183.05$  = Rs.1830.50

### (c) Area of Triangular Region when Three Sides are Given

If  $a$ ,  $b$ ,  $c$  are measure of sides of a triangle ABC then:

$$\text{Area of triangle ABC} = \Delta = \sqrt{S(S-a)(S-b)(S-c)} \text{ where } S = \frac{a+b+c}{2}$$

Let us prove this formula.

We have proved in the previous section that:

$$\text{Area of triangle ABC} = \Delta = \frac{1}{2} bc \sin \alpha$$

Using half angle formula, we have:

$$\Delta = \frac{1}{2} bc \times 2 \sin \frac{\alpha}{2} \cos \frac{\alpha}{2} = bc \times \sqrt{\frac{(S-b)(S-c)}{bc}} \times \sqrt{\frac{S(S-a)}{bc}}$$

### Check Point:

Prove that

$$(i) \Delta = \frac{1}{2} b^2 \frac{\sin \alpha \sin \gamma}{\sin \beta}$$

$$(ii) \Delta = \frac{1}{2} c^2 \frac{\sin \alpha \sin \beta}{\sin \gamma}$$

### Check Point:

Two sides of a triangle are 20cm long each. Find the angle between both sides if area of triangle is 200 sq. unit.

$$\Delta = bc \times \sqrt{\frac{(S-b)(S-c)}{bc}} \times \sqrt{\frac{S(S-a)}{bc}} = \sqrt{S(S-a)(S-b)(S-c)}$$

This formula is known as hero's formula or Heron's formula.

**Example:**

Find area of a triangle with dimensions 16, 18, 20.

**Solution:** Let  $a=16, b=18, c=20$ .

$$\text{Area of triangle} = \sqrt{S(S-a)(S-b)(S-c)}$$

$$\text{Where } S = \frac{16+18+20}{2} = \frac{54}{2} = 27$$

$$\text{Now, area of triangle} = \sqrt{27(27-16)(27-18)(27-20)} = \sqrt{18711}$$

$$= 136.79 \text{ sq. unit}$$


### Exercise 8.4

1. Find the area of triangular region ABC in the following.


- |  |  |  |
|--|--|--|
| (i) $a=8, b=14, \gamma=68.7^\circ$                   | (ii) $b=30, c=25, \alpha=46^\circ$                   |  |
| (iii) $a=20, c=15, \beta=25^\circ$                   | (iv) $\beta=46^\circ 50', a=43, c=52$                |  |
| (v) $b=4.5, c=2.5, \alpha=65.2^\circ$                | (vi) $a=2, b=2, c=5$                                 |  |
| (vii) $a=12, \alpha=44^\circ, \beta=60^\circ$        | (viii) $b=30, \alpha=40^\circ, \beta=80^\circ$       |  |
| (ix) $c=10, \alpha=75^\circ, \beta=45^\circ$         | (x) $b=21, \beta=30^\circ 15', \gamma=110^\circ$     |  |
| (xi) $a=18.4, \alpha=65^\circ 10', \beta=95.5^\circ$ | (xii) $c=25, \alpha=52.7^\circ, \gamma=79^\circ 24'$ |  |
| (xiii) $18, 21, 32$                                  | (xiv) $20, 26, 37$                                   | (xv) $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}$ |
| (xvi) $12.4, 13.7, 20.2$                             | (xvii) $1.6, 2.6, 4.1$                               |  |

2. The adjacent sides of parallelogram ABCD measure 12 and 15. The measure of one angle of the parallelogram is  $135^\circ$ . Find the area of the parallelogram.
3. Three streets intersect in pairs enclosing a small triangular park. The measures of the distances between the intersections are 30 m, 34 m, and 27 m. Find the area of the park.
4. A field is bordered by two pairs of parallel roads so that the shape of the field is a parallelogram. The lengths of two adjacent sides of the field are 2 km and 3 km, and the length of the shorter diagonal of the field is 3 km.
  - a. Find the cosine of the acute angle of the parallelogram.
  - b. Find the exact value of the sine of the acute angle of the parallelogram.
  - c. Find the exact value of the area of the field.
  - d. Find the area of the field to the nearest integer.
5. The roof of a shed consists of four congruent isosceles triangles. The length of each equal side of one triangular section is 22.0 feet and the measure of the vertex angle of each triangle is  $75^\circ$ . Find the area of one triangular section of the roof.

6. A garden is in the shape of an isosceles trapezoid. The lengths of the parallel sides of the garden are 30 feet and 20 feet, and the length of each of the other two sides is 10 feet. If a base angle of the trapezoid measures  $60^\circ$ , find the exact area of the garden.
7. In triangle ABC,  $\angle B = 30^\circ$  and in triangle DEF,  $\angle E = 150^\circ$ . Show that if AB = DE and BC = EF, the areas of the two triangles are equal.
8. Area of a triangular garden is  $150\text{m}^2$ . If two corner angles of a side measure  $40^\circ$  and  $65^\circ$ , find the length of that side. Also find the third angle of garden.
9. An equilateral triangle is inscribed in a circle of radius 6cm.  
Find the area of triangle.



10. Laiba wants to draw triangle ABC with  $AB = 15\text{ cm}$ ,  $BC = 8\text{ cm}$ , and an area of  $40\text{ cm}^2$ .
- What must be the sine of  $\angle B$ ?
  - Find the measure of  $\angle B$ .
11. Let ABCD be a parallelogram with  $AB = c$ ,  $BC = a$ , and  $\angle B = \theta$ .
- Write a formula for the area of parallelogram in terms of  $c$ ,  $a$ , and  $\theta$ .
  - For what value of  $\theta$  does parallelogram have the greatest area?



## Circles Connected with Triangles

### (a) Circumcircle

A circle passing through the vertices of any triangle (polygon) is called circumcircle.

The centre of this circle is called the circumcentre and its radius is called the circumradius.

The radius of circumcircle is denoted by R.

### (b) Incircle

A circle drawn inside and touching the sides of a triangle (polygon) is called incircle or inscribed circle. It is the largest circle contained in the triangle. The centre of this circle is called the incentre usually denoted by I and its radius is called the inradius denoted by r.


### (c) Escribed Circle

A circle touching one of the sides of a triangle externally and the extensions of its two other sides internally, is called an escribed circle (e-circle or ex-circle). The centre of this circle is called the e-centre and its radius is called the e-radius.

If a circle touches extended arms of angle A (circle is drawn opposite to vertex A) of a triangle ABC, then its centre and radius are respectively denoted by  $I_1$  and  $r_1$ . Similarly,

(i) Centre and radius of e-circle opposite to vertex B are denoted by  $I_2$  and  $r_2$ .


(ii) Centre and radius of e-circle opposite to vertex C are denoted by  $I_3$  and  $r_3$ .



Circumcircle



Incircle




Escribed circle

#### Key Fact:


- (i) Circumcentre is the point of intersection of right bisectors of sides of a triangle.
- (ii) Incentre is the point of intersection of angle bisectors of a triangle.
- (iii) E-centre is the point of intersection of one internal bisector of angle and external bisectors of remaining angles of a triangle.

### Circum-Radius of Circle In-terms of Side and Opposite Angle of Triangle


Let O be the circumcentre of triangle ABC. Draw a diameter BD. Also join C to D.



(i)



(ii)



(iii)

#### Case-1: When $\alpha$ is Acute

In figure (i),  $\alpha$  and  $\angle BDC$  are angles in the same segment associated with chord BC.  
 $\therefore \angle BDC = \alpha$

Now in right angled triangle BCD,

$$\frac{BC}{BD} = \sin(\angle BDC) = \sin \alpha$$

$$\frac{a}{2R} = \sin \alpha \Rightarrow R = \frac{a}{2 \sin \alpha} \text{ where } BD = 2R \text{ and } BC = a.$$

#### Case-2: When $\alpha$ is Obtuse

In figure (ii),  $\alpha + \angle BDC = 180^\circ$  because ABDC is a cyclic quadrilateral.  
 $\therefore \angle BDC = \pi - \alpha$

Now in right angled triangle BCD,

$$\frac{BC}{BD} = \sin(\angle BDC) = \sin(\pi - \alpha) \text{ where } \sin(\pi - \alpha) = \sin \alpha. \quad \frac{a}{2R} = \sin \alpha \Rightarrow R = \frac{a}{2\sin \alpha}$$

### Case-3: When $\alpha$ is Right

As, angle in a semicircle is right angle. Therefore, ABC is a right angled triangle and  $\overline{BC}$  is a diameter. Hence  $a = BC = 2R$

$$\therefore \frac{a}{2R} = 1 = \sin \frac{\pi}{2} = \sin \alpha$$

$$\frac{a}{2R} = \sin \alpha \Rightarrow R = \frac{a}{2\sin \alpha}$$

Similarly, we can prove that:

$$R = \frac{b}{2\sin \beta} \text{ and } R = \frac{c}{2\sin \gamma}$$

Combining all results, we get:

$$R = \frac{a}{2\sin \alpha} = \frac{b}{2\sin \beta} = \frac{c}{2\sin \gamma}$$

### Circum-Radius of Circle In-terms of Sides of Triangle

We have proved in the previous section, that:

$$R = \frac{a}{2\sin \alpha} = \frac{a}{2 \times 2\sin \frac{\alpha}{2} \cos \frac{\alpha}{2}} = \frac{a}{4 \times \frac{\sqrt{(S-b)(S-c)}}{bc} \frac{\sqrt{S(S-a)}}{bc}}$$

$$R = \frac{a}{4 \times \sqrt{\frac{(S-b)(S-c)}{bc}} \sqrt{\frac{S(S-a)}{bc}}} = \frac{abc}{4\sqrt{S(S-a)(S-b)(S-c)}}$$

$$R = \frac{abc}{4\Delta}$$

Where  $\Delta = \sqrt{S(S-a)(S-b)(S-c)}$  and  $\sin \alpha = 2 \sin \frac{\alpha}{2} \cos \frac{\alpha}{2}$ .

### In-Radius of a Circle in a Triangle


Consider a triangle ABC. Draw the three angle bisectors of triangle which intersect at point I which is in-centre of circle.

By drawing angle bisectors, the given triangle ABC has been divided into three triangles IAB, IBC and IAC.

Draw altitude ID of triangle IBC, altitude IE of triangle IAC and altitude IF of triangle IAB. Note that

$ID = IE = IF = \text{radius of in-circle} = r$

Now if  $AB = c$ ,  $BC = a$ ,  $AC = b$  and  $\Delta$  is the area of



triangle ABC, then from the figure:

area of triangle ABC = area of triangle IBC + area of triangle IAC + area of triangle IAB

$$\Delta = \frac{1}{2} \times BC \times ID + \frac{1}{2} \times AC \times IE + \frac{1}{2} \times AB \times IF$$

$$\Delta = \frac{1}{2} \times a \times r + \frac{1}{2} \times b \times r + \frac{1}{2} \times c \times r = \frac{1}{2} \times r \times (a+b+c)$$

$$\Delta = r \times \frac{a+b+c}{2} = rS$$

$$r = \frac{\Delta}{S}$$

#### Key Fact:

The two geometrical figures, circle and triangle are made up of distinct conception. The triangle is made with straight lines, whereas circle is made by curve lines (arcs). The beauty of circum-circle and in-circle of a triangle is a combination of a triangle and a circle. Engineers use this combination in making parts of machines.

#### E-Radius of a Circle of a Triangle

Consider a triangle ABC. Draw internal bisector of  $\angle A$  and external bisectors of  $\angle B$  and  $\angle C$  of triangle intersecting at  $I_1$  which is the centre of ex-circle.

By drawing angle bisectors, the given triangle ABC has been divided into three triangles  $I_1AB$ ,  $I_1BC$  and  $I_1AC$ .

Draw altitude  $I_1D$  of triangle  $I_1AB$ , altitude  $I_1E$  of triangle  $I_1BC$  and altitude  $I_1F$  of triangle  $I_1AC$ . Note that

$$I_1D = I_1E = I_1F = \text{radius of e-circle} = r_1$$

Now if  $AB = c$ ,  $BC = a$ ,  $AC = b$  and  $\Delta$  is the area of triangle ABC, then from the figure:


area of triangle ABC = area of triangle  $I_1AB$  + area of triangle  $I_1AC$  + area of triangle  $I_1BC$

$$\Delta = \frac{1}{2} \times AB \times I_1D + \frac{1}{2} \times AC \times I_1F - \frac{1}{2} \times BC \times I_1E$$

$$\Delta = \frac{1}{2} \times c \times r_1 + \frac{1}{2} \times b \times r_1 - \frac{1}{2} \times a \times r_1 = \frac{1}{2} \times r_1 \times (c+b-a)$$

$$\Delta = r_1 \times \frac{2S - a - b}{2} = r_1 \times \frac{2S - 2a}{2} = r_1 \times \frac{2(S - a)}{2}$$

$$r_1 = \frac{\Delta}{S - a}$$



Similarly, we can prove that:

$$r_2 = \frac{\Delta}{S - b} \quad \text{and} \quad r_3 = \frac{\Delta}{S - c}$$

**Example:**

Find  $r$ ,  $R$  and  $r_2$  when  $a = 8$ ,  $b = 9$ ,  $c = 11$ .

**Solution:** Given:  $a = 8$ ,  $b = 9$ ,  $c = 11$

To find:  $r$ ,  $R$  and  $r_2$

First of all, we find  $S$  and  $\Delta$ .

$$S = \frac{8+9+11}{2} = \frac{28}{2} = 14$$

$$\text{and } \Delta = \sqrt{S(S-a)(S-b)(S-c)} \\ = \sqrt{14(14-8)(14-9)(14-11)} = \sqrt{1260} = 35.5$$

$$\text{Now, } r = \frac{\Delta}{S} = \frac{35.5}{14} = 2.54$$

$$R = \frac{abc}{4\Delta} = \frac{792}{142} = 5.58$$

$$r_2 = \frac{\Delta}{S-b} = \frac{35.5}{14-9} = \frac{35.5}{5} = 7.1$$

**Example:**

Prove that  $r_2 = 4R \cos \frac{\alpha}{2} \cos \frac{\gamma}{2} \sin \frac{\beta}{2}$

**Solution:**

$$\begin{aligned} \text{R.H.S} &= 4R \cos \frac{\alpha}{2} \cos \frac{\gamma}{2} \sin \frac{\beta}{2} \\ &= 4 \times \frac{abc}{4\Delta} \times \sqrt{\frac{S(S-a)}{bc}} \times \sqrt{\frac{S(S-c)}{ab}} \times \sqrt{\frac{(S-a)(S-c)}{ac}} \\ &= \frac{abc}{\Delta} \times \sqrt{\frac{S^2(S-a)^2(S-c)^2}{a^2b^2c^2}} = \frac{abc}{\Delta} \times \frac{S(S-a)(S-c)}{abc} \\ &= \frac{1}{\Delta} \times \frac{S(S-a)(S-b)(S-c)}{S-b} = \frac{\Delta^2}{\Delta(S-b)} \\ &= \frac{\Delta}{S-b} = r_2 = \text{L.H.S} \end{aligned}$$

We know that

$$R = \frac{a}{2 \sin \alpha} = \frac{b}{2 \sin \beta} = \frac{c}{2 \sin \gamma}$$

$$\Rightarrow \frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma} = 2R$$

$$\Rightarrow \frac{a}{\sin \alpha} = \frac{b}{\sin \beta} = \frac{c}{\sin \gamma}$$

Which is law of sines.

Find  $r_1$ ,  $r_2$  and  $r_3$

when  $a = 5.4$ ,  $b = 7.6$  and  $c = 10.2$ .

Show that

$$4R \cos \frac{\alpha}{2} \cos \frac{\beta}{2} \sin \frac{\gamma}{2} = r_3$$

**Exercise 8.5**

1. Find  $r$ ,  $R$ ,  $r_1$ ,  $r_2$  and  $r_3$  when

- (i)  $a = 10$ ,  $b = 13$ ,  $c = 17$   
 (iii)  $a = 3$ ,  $b = 4$ ,  $c = 5$

- (ii)  $a = 22$ ,  $b = 24$ ,  $c = 30$   
 (iv)  $a = 50$ ,  $b = 60$ ,  $c = 70$

2. If ABC is an equilateral triangle, prove that:

- (i)  $r : R : r_1 = 1 : 2 : 3$   
 (ii)  $r_1 : r_2 : r_3 = 1 : 1 : 1$

3. Show that:

- (i)  $\frac{1}{ab} + \frac{1}{bc} + \frac{1}{ca} = \frac{2S}{abc} = \frac{1}{2rR}$  (ii)  $\frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3} = \frac{S}{\Delta}$   
(iii)  $r_1 r_2 + r_2 r_3 + r_1 r_3 = S^2$  (iv)  $r_1 + r_2 + r_3 - r = 4R$   
(v)  $\sqrt{r r_1 r_2 r_3} = \Delta$  (vi)  $r_1 r_2 r_3 = r S^2$

4. Show that:  $(\sin \alpha + \sin \beta + \sin \gamma) = \frac{S}{R}$

5. Prove that:  $(r_1 + r_2) \tan \frac{\gamma}{2} = (r_3 - r) \cot \frac{\gamma}{2} = c$

6. Show that:

- (i)  $4rR \cos \frac{\alpha}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2} = r^2 \cot \frac{\alpha}{2} \cot \frac{\beta}{2} \cot \frac{\gamma}{2} = \Delta$   
(ii)  $r = 4R \sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sin \frac{\gamma}{2} = S \tan \frac{\alpha}{2} \tan \frac{\beta}{2} \tan \frac{\gamma}{2}$   
(iii)  $r = a \sin \frac{\beta}{2} \sin \frac{\gamma}{2} \sec \frac{\alpha}{2} = b \sin \frac{\alpha}{2} \sin \frac{\gamma}{2} \sec \frac{\beta}{2} = c \sin \frac{\alpha}{2} \sin \frac{\beta}{2} \sec \frac{\gamma}{2}$

7. Prove that:

(i)  $4R \sin \frac{\alpha}{2} \cos \frac{\beta}{2} \cos \frac{\gamma}{2} = r_1$  (ii)  $4R \cos \frac{\alpha}{2} \sin \frac{\beta}{2} \cos \frac{\gamma}{2} = r_2$

8. Prove that:

(i)  $r_1 = S \tan \frac{\alpha}{2}$  (ii)  $r_2 = S \tan \frac{\beta}{2}$  (iii)  $r_3 = S \tan \frac{\gamma}{2}$

9. Prove that:

(i)  $r = (S-a) \tan \frac{\alpha}{2} = (S-b) \tan \frac{\beta}{2} = (S-c) \tan \frac{\gamma}{2}$

10. Find the radius of in-circle and circum-circle of a triangle having sides: 7cm, 12cm and 15cm.

11. The measures of sides of a triangle are 5cm, 9cm, 10cm. Find the circumference of ex-circle:

(i) opposite to smaller side (ii) opposite to larger side

12. Find the area and circumference of in-circle of a triangle having sides: 2.5cm, 7.3 cm and 6.2cm.


### Trigonometric Problems in Three Dimensions

#### 3-D Trigonometry

3-D trigonometry is an application of the trigonometric skills developed for 3-dimensional shapes in order to find unknown sides and angles.

In this topic, we will learn about 3-D trigonometry. We will use combined knowledge of Pythagoras' theorem, trigonometric ratios, the law of sines (sine rule) and the law of cosines (cosine rule) to find missing angles and sides of triangles in 3-dimensional shapes.

The flowchart below can help to determine which path we need to use:



In order to find a missing angle or side within a 3-dimensional shape:


**Step 1:** Find the missing side or angle of a triangle.

**Step 2:** Sketch and label the second triangle using information from Step 1.

**Step 3:** Calculate the missing side or angle of the final triangle.

Let us solve some examples to understand the method.

**Example:** Find the length of  $\overline{EC}$  in the given cuboid.




**Step 1:** Finding the length of missing side  $\overline{AC}$  of right triangle ABC.


$\overline{AC}$  is hypotenuse of right triangle ABC. We can find its length by using simple trigonometric ratios.

In right triangle ABC:

$$\begin{aligned}\sin \theta &= \frac{16}{H} \\ \Rightarrow H &= \frac{16}{\sin \theta} = \frac{16}{\sin 45^\circ} \\ &= \frac{16}{0.707} = 22.63\end{aligned}$$



**Step 2:** Sketching and labeling the second right triangle EAC using information from step 1.



**Step 3:** EAC is a right-angled triangle, so we can use Pythagoras' Theorem to find the length of  $\overline{EC}$ .

Let  $EC = x \text{ cm}$

$$x^2 = (12)^2 + (22.63)^2$$

$$x^2 = 144 + 512.12 = 656.12$$

Taking square root on both sides, we get:

$$x = \sqrt{656.12} = 25.61$$

Therefore,  $EC = 25.61 \text{ cm}$


**Example:** ABCDEF is an isosceles triangular prism.

BC is 8cm,  $\angle DFE = 70^\circ$  and angle FCD =  $45^\circ$ . Find CD.

**Solution:**

To find CD, we are required DF.

Therefore, first consider  $\triangle DEF$  to find DF.




In right  $\triangle DGF$ :

$$\cos 70^\circ = \frac{4}{DF}$$

$$\Rightarrow DF = \frac{4}{\cos 70^\circ}$$

$$= 11.695 \text{ cm}$$




Now in right  $\triangle CFD$ :


$$\sin 45^\circ = \frac{11.695}{CD}$$

$$\Rightarrow CD = \frac{11.695}{\sin 45^\circ}$$

$$= 16.54 \text{ cm}$$



**Example:** Find  $\angle x$  in the following figure where P is mid-point of CD.



**Solution:** To find the value of  $\angle x$ , we need  $EF = a$  which is length of a side of  $\triangle AEF$ .

In right  $\triangle AEF$ :

$$\frac{EF}{AF} = \sin 30^\circ$$

$$\Rightarrow \frac{a}{40} = \frac{1}{2}$$

$$\Rightarrow a = 20$$

$$\therefore EF = 20 \text{ cm}$$

As P is a midpoint of  $\overline{DC}$ , therefore  $\triangle PEF$  is an isosceles triangle.

Therefore, using law of cosines, we have:



$$\cos x = \frac{50^2 + 50^2 - 20^2}{2(50)(50)}$$

$$\cos x = \frac{2500 + 2500 - 400}{5000}$$

$$\cos x = \frac{4600}{5000} = 0.92$$


$$\Rightarrow x = \cos^{-1}(0.92) = 23.1^\circ$$

$$\therefore \angle EPF = 23.1^\circ$$




### Exercise 8.6


1. In the figure, ABCDEFGH is a cuboid. Calculate CF.




2. Find the angle ACF in the given cuboid in the following figure.




3. In the figure, perimeter of square based pyramid is 36 cm. Find the length of diagonal of the base and height of the pyramid.



4. The diagram shows a cylinder. Find the value of  $\gamma$  in triangle ABC.




5. In the figure, three workers are standing at positions B, D and H of a container of cuboid shape.



- (i) Find the distance between workers at B and D.
- (ii) Find the distance between workers at B and H.
- (iii) Find  $\angle DBH$ .

6. In the figure, O is the centre of the square based pyramid ABCDE. Calculate the angle between side AB and plane BCDE if  $OA = 6 \text{ cm}$ .




7. Floor of a room is 9 m long and 6 m wide. The angle of elevation from the bottom left corner to the top right corner of the room is  $49^\circ$ .
- Find the distance from one corner of the floor to the opposite corner of the floor.
  - Find the height of the room.
  - Find the angle of elevation from the bottom corner of the 9 m long wall to the opposite top corner of the wall.
  - Find the angle of depression from the top corner of the 6 m long wall to the opposite bottom corner of the wall.
8. Three satellites A, B, and C, used for GPS navigation are orbiting the Earth in the same plane. The distance between satellites A and B is 15 km. If D is a house on Earth, such that  $\angle BAD = 70^\circ$  and ACBD is a rhombus then:
- Determine the distance between satellite A and satellite C.
  - Determine the distance between satellites B and C.
  - Find the distance from satellite C to the house D.

#### MISCELLANEOUS EXERCISE-8

1. Select the correct option in the following.

- In right  $\triangle ABC$ ,  $a = 2\text{cm}$ ,  $c = 4\text{cm}$ , what is  $\alpha$ ?
   
 (a)  $30^\circ$       (b)  $45^\circ$       (c)  $60^\circ$       (d)  $120^\circ$
- If in a triangle,  $a = 10$ ,  $b = 15$ ,  $\alpha = 32^\circ$  then  $\beta = \dots$ 
  
 (a)  $42.5^\circ$       (b)  $46.5^\circ$       (c)  $52.7^\circ$       (d)  $62.8^\circ$
- Area of an equilateral triangle having side  $a$  is:
   
 (a)  $\frac{\sqrt{3}}{8}a^2$       (b)  $\frac{\sqrt{3}}{4}a^2$       (c)  $\frac{\sqrt{3}}{16}a^2$       (d)  $\frac{\sqrt{3}}{16}a^2$
- If  $a$ ,  $a$  and  $b$  are length sides of an isosceles triangle, then  $S = \dots$ 
  
 (a)  $\frac{a}{2} + b$       (b)  $a + \frac{a+b}{2}$       (c)  $a - \frac{a+b}{2}$       (d)  $a + \frac{b}{2}$

- v. Area of a triangle ABC with  $a = 20, b = 30, \gamma = 90^\circ$  is:  
 (a) 0 (b) 30 (c) 300 (d) 600
- vi. For an equilateral triangle,  $r : r_1 : R = \dots$   
 (a)  $1 : 2 : 3$  (b)  $3 : 2 : 1$  (c)  $1 : 1 : 2$  (d)  $1 : 3 : 2$
- vii. Radius of circum-circle for sides 6, 8, 10 is:  
 (a) 6 (b) 5 (c) 4 (d) 2
- viii. If  $a = 5, b = 10, c = 20$  are sides of a triangle ABC, then angle  $\alpha$  is:  
 (a) not possible (b) acute (c) obtuse (d)  $0^\circ$
- ix. Radius of circum-circle  $R = \dots$   
 (a)  $\frac{a}{2} \sec \frac{\alpha}{2}$  (b)  $\frac{b}{2} \sec \frac{\beta}{2}$  (c)  $\frac{c}{2} \csc \frac{\gamma}{2}$  (d)  $\frac{c}{2} \cos \frac{\gamma}{2}$
- x. If in a triangle ABC,  $a = b = c$ , then  $\tan \frac{\alpha}{2} = \dots$   
 (a)  $\sqrt{\frac{S-a}{a}}$  (b)  $\sqrt{\frac{S-b}{b}}$  (c)  $\sqrt{\frac{S-c}{c}}$  (d) all (a), (b) & (c)
- xi. Shadow of a man 5.6ft tall is making an angle of elevation of  $45^\circ$  with the Sun. What is the length of shadow?  
 (a) 2.8ft (b) 5.6ft (c) 8.4ft (d) 11.2ft
- xii.  $R(\sin \alpha + \sin \beta + \sin \gamma) = \dots$   
 (a)  $S$  (b)  $S^2$  (c)  $\frac{1}{S}$  (d)  $\frac{1}{S^2}$
2. Solve the following oblique triangles.
- (i)  $a = 8, \alpha = 15^\circ, \beta = 20^\circ$  (ii)  $a = 34, b = 41, \alpha = 115^\circ$   
 (iii)  $b = 17.3, c = 14.5, \beta = 98.2^\circ$  (iv)  $c = 55.4, \alpha = 115^\circ, \beta = 45^\circ$   
 (v)  $a = 34, c = 48, \beta = 108^\circ$  (vi)  $a = 44, b = 33, c = 55$
3. The diagonals of a parallelogram measure 12 centimeters and 22 centimeters and intersect at an angle of 143 degrees. Find the length of the longer sides of the parallelogram.




4. Usman and Abubakar follow a triangular path when they take a walk. They walk from home for  $1.5\text{ km}$  along a straight road, turn at an angle of  $100^\circ$ , walk for another  $0.95\text{ km}$  and then return home.
- Find the length of the last portion of their walk.
  - Find the total distance that they walk.
5. A kite is in the shape of a quadrilateral with two pair of congruent adjacent sides. The lengths of two sides are  $20.0$  inches and the lengths of the other two sides are  $35.0$  inches. The two shorter sides meet at an angle of  $115^\circ$ .
- Find the length of the diagonal between the points at which the unequal sides meet.
  - Using the answer to part a, find the measure of the angle at which the two longer sides meet.
6. Three streets intersect in pairs enclosing a small triangular park. The measures of the distances between the intersections are  $55\text{ m}$ ,  $63\text{ m}$ , and  $77\text{ m}$ . Find the area of the park.
7. The base of an isosceles triangle measures  $14.5$  centimeters and the vertex angle measures  $110^\circ$ .
- Find the measure of one of the congruent sides of the triangle.
  - Find the perimeter of the triangle.
8. Three streets intersect in pairs enclosing a small park. Two of the angles at which the streets intersect measure  $80^\circ$  and  $60^\circ$ . The length of the longest side of the park is  $90\text{ m}$ . Find the lengths of the other two sides of the park.
9. Aamir wants to draw a parallelogram with the measure of one side  $12\text{ cm}$ , the measure of one diagonal  $10\text{ cm}$  and the measure of one angle  $120^\circ$ . Is this possible? Explain why or why not. (Hint: Use law of sines.)
10. The angle of depression from an observer in an apartment complex to a pipe on the building next door is  $55^\circ$ . From a point five stories below the original observer, the angle of inclination to the pipe is  $20^\circ$ . Find the distance from each observer to the pipe and the distance from the pipe to the apartment complex. It is given that one story of a building is 9 feet high.
11. A geologist wants to measure the diameter of a crater. From her camp, it is 4 miles to the northern-most point of the crater and 2 miles to the southern-most point. If the angle between the two lines of sight is  $117^\circ$ , what is the diameter of the crater?
12. Side of an equilateral triangle is  $6\text{cm}$  long. Find the circumference of circum-circle and ex-circle.
13. Use the Law of Cosines to prove that if the angle between two congruent sides of a triangle measures  $60^\circ$ , the triangle is equilateral.
14. Prove that:  $\frac{1}{r_1^2} + \frac{1}{r_2^2} + \frac{1}{r_3^2} + \frac{1}{r_s^2} = \frac{a^2 + b^2 + c^2}{\Delta^2}$

15. Prove that: (i)  $r_1 = a \cos \frac{\beta}{2} \cos \frac{\gamma}{2} \sec \frac{\alpha}{2}$     (ii)  $r_2 = b \cos \frac{\alpha}{2} \cos \frac{\gamma}{2} \sec \frac{\beta}{2}$


(iii)  $r_3 = c \cos \frac{\alpha}{2} \cos \frac{\beta}{2} \sec \frac{\gamma}{2}$

16. A mountain climber is on top of a mountain that is 680 m high. The angles of depression of two points on opposite sides of the mountain are  $48^\circ$  and  $32^\circ$ . How long would a tunnel be that runs between the two points?

17. Find the measure of  $\angle C$  in the adjoining figure.



18. Sides of a square prism are 12cm, 5cm and 5cm long as shown below. Find the measure of angle  $a$ .



# Chord and Arcs of a Circle


**After studying this unit students will be able to:**

- One and only one circle can pass through three non collinear points.
- A straight line drawn from the centre of a circle to bisect the chord is perpendicular to the chord.
- Perpendicular drawn from the centre of a circle on a chord, bisects it.
- Two congruent chords of a circle are equidistant from the centre.
- Two chords of a circle equidistant from the centre are congruent.
- If two arcs of a circle (or of congruent circles) are congruent, then the corresponding chords are equal.
- If two chords of a circle (or of congruent circles) are equal, then their corresponding arcs (minor, major or semicircle) are congruent.
- Equal chords of a circle (or of congruent circles) subtend equal angles at the centre (at the corresponding centres).
- If the angles subtended by two chords of a circle (or congruent circles) at the centre (corresponding circles) are equal, the chords are equal.
- Apply concepts of chords and arcs to real life word problems such as decorative features, rainbow, bridges and roller coaster track.

Circle theorems are statements in geometry that state important results related to circles that are used to solve various questions in geometry. They have very useful applications within design and engineering.

These theorems show relationships between angles within the geometry of a circle. We can use these theorems along with prior knowledge of other angle properties to calculate missing angles, without the use of a protractor.

Can you find the value of angle in the figure below.



# Chords of a Circle

## Definition


A line segment joining two points of a circle is called chord of that circle.

## Theorem 9.1

**Statement:** One and only one circle can pass through three non collinear points.

### Class Activity:

- Take three non-collinear points A, B and C on the paper and join them to obtain the triangle.
- Draw right bisectors of the sides. They will meet at only one point P (say).
- With centre P, draw a circle of radius PA.  
What do you notice? Does the circle pass through B and C also?



**Corollaries:** (i) One and only one circle can pass through two points lying on the same line.  
(ii) Two circles cannot intersect each other at more than two points.

### Example:

Construct a square with sides 6 cm long. Show that one and only one circle can be drawn through vertices of the square. Measure the radius of the circle.

### Solution:

- Construct a square ABCD having side equal to 6 cm long.
- Draw diagonals  $\overline{AC}$  and  $\overline{BD}$ .
- The two diagonals bisect each other at point O.
- With centre O, construct a circle with radius OA.
- You will see that the circle will also pass through vertices B, C and D.

As diagonals  $\overline{AC}$  and  $\overline{BD}$  bisect each other. Therefore:

$$OA = OB = OC = OD$$

Furthermore, there is only one centre O of the circle, therefore we can say that only one circle can be drawn through vertices of the square.

Now in right triangle ABC:


$$(AC)^2 = (AB)^2 + (BC)^2$$

$$(AC)^2 = (6)^2 + (6)^2 = 36 + 36 = 72$$

Taking square root on both sides, we have:

$$AC = \sqrt{72} = 6\sqrt{2} \text{ cm}$$

$$\text{Radius} = OA = \text{half of } AC = 3\sqrt{2} \text{ cm}$$



### Key Fact:

If two chords of a circle do not pass through the centre, they cannot bisect each other.

## Theorem 9.2


**Statement:** A straight line drawn from the centre of a circle to bisect the chord is perpendicular to the chord.

### Class Activity:

- Draw a chord AB of a circle with centre O.
- Take a point C on the chord such that  $AC = BC$ .
- Join O to C and measure  $\angle OCA$  and  $\angle OCB$ .

Are the angles equal? Yes! they are.

$$\therefore \angle OCA = \angle OCB \text{ i.e., } \overline{OC} \perp \overline{AB}$$



### Corollaries:

- If a diameter or radial segment of a circle bisects the chord, it is perpendicular to the chord.
- The diameter or radial segment of a circle passes through the mid points of two parallel chords of a circle.

## Theorem 9.3


**Statement:** Perpendicular drawn from the centre of a circle on a chord, bisects it.

### Class Activity:

- Draw a chord AB of a circle with centre O.
- Take a point C on the chord such that  $\angle OCA = 90^\circ$ .
- Measure AC and BC.

Are they equal?

Yes! they are. i.e.,  $AC = BC$



### Corollaries:

- If a diameter of a circle is perpendicular to the chord, it bisects the chord.
- Perpendicular (right) bisector of a chord of a circle passes through the centre of a circle.

### Example:

Diameter and chord of a circle are 20 cm and 16 cm long respectively and are perpendicular to each other. Calculate the distance of chord from the centre.

### Solution:

$$\text{Diameter} = CD = 20 \text{ cm}$$

$$\text{Radius} = OA = 10 \text{ cm}$$

$$\text{Chord} = AB = 16 \text{ cm}$$


As the diameter and chord are perpendicular to each other.

Therefore, diameter bisects the chord.

$$\text{Hence, } AE = BE = 8 \text{ cm}$$

Now in right triangle AEO:

$$\begin{aligned}
 (AE)^2 + (OE)^2 &= (OA)^2 \Rightarrow (8)^2 + (OE)^2 = (10)^2 \\
 \Rightarrow (OE)^2 &= (10)^2 - (8)^2 = 100 - 64 = 36 \\
 \Rightarrow OE &= \sqrt{36} = 6 \text{ cm} \\
 \therefore \text{The distance of chord from the centre} &= 6 \text{ cm}
 \end{aligned}$$




### Key Fact:

Diameters of a circle bisect each other.


### Exercise 9.1

- Construct an equilateral triangle with side 5 cm long. Show that one and only one circle can be drawn through the vertices of triangle. What is the radius of circle drawn?
- Given that P is the centre of each of the circles. Find the values of unknown if line segment drawn from the centre of each circle is perpendicular to the chord.


(i)




(ii)




- Find the length of diameter CD of the circle when  $AB = 10\text{ cm}$ ,  $PE = 12\text{ cm}$ . Also find  $CE$ .




- In the following diagram,  $OB = 15\text{ cm}$ ,  $OF = 9\text{ cm}$ . Find the length of chord CD given that  $\overline{OF} \perp \overline{CD}$ .




- Calculate the length of chord AB in the following diagram if  $\overline{OC} \perp \overline{AB}$ .



6. Construct a rectangle EFGH such that  $EF = 6\text{ cm}$  and  $FG = 4\text{ cm}$ . Draw a circle passing through its vertices and prove that it is the only circle that can be drawn through vertices.
7. Take four non collinear points A, B, C and D such that  $AB = BC = DB = 4\text{ cm}$ . Draw all possible circles that can pass through A, C and D.
8. In the diagram,  $PU = 16\text{ cm}$  and  $RS = 10\text{ cm}$ .
- Express OU in terms of radius r.
  - Form an equation in r and solve it to find radius r.
- Hint:  $16 - OU = r$




9. In the following diagram,  $AB = 16\text{ cm}$ ,  $DE = 4\text{ cm}$ . Find the radius of the circle.



10. Given that diameter and chord of a circle are  $10\text{ cm}$  and  $8\text{ cm}$  long respectively. The diameter bisects the chord and the distance between chord and centre of the circle is  $3\text{ cm}$ . Show that the diameter bisects the chord perpendicularly.

11. In the figure O is centre of the circle. AB is the chord and  $CD \perp AB$  passes through centre. Show that  $AC = BC$ .




### Theorem 9.4

**Statement:** Two congruent chords of a circle are equidistant from the centre.

**Class Activity:**

- Draw two equal chords AB and CD of a circle as shown in the adjoining figure.
- Draw perpendiculars OE and OF on the chords AB and CD respectively.



(iii) Measure OE and OF.

Are they equal? Yes! they are equal.

### Theorem 9.5

#### Statement:


Two chords of a circle equidistant from the centre, are congruent.

#### Class Activity:

(i) Draw two chords AB and CD of a circle such that they are equidistant from the centre as shown in the adjoining figure.

(ii) Measure chords AB and CD.

You will see that they are equal in length.



#### Corollary:

(i) If two chords of a circle are unequal, then longer chord is nearer to the centre and vice versa.

(ii) If two chords of two congruent circles are congruent, then they are equidistant from their respective centres and vice versa.

#### Example:

In the adjoining figure,  $\overline{AB} \parallel \overline{CD}$  such that  $AB = 24\text{ cm}$  and  $CD = 10\text{ cm}$ . Find the radius of the circle if distance between chords is  $17\text{ cm}$ .

#### Solution:

Let the radius of circle be  $r\text{ cm}$  with centre P.

Draw  $\overline{PE} \perp \overline{AB}$  and  $\overline{PF} \perp \overline{CD}$

$$\Rightarrow BE = \frac{1}{2}AB \quad \text{and} \quad CF = \frac{1}{2}CD$$

As the chords AB and CD are parallel, therefore points E, P and F are collinear and  $EF = 17\text{ cm}$

Let  $PE = x$ , then  $PF = 17 - x$

Join P to B and C, then  $PB = PC = r$

Now, PBE and PCF are right angled triangles. Therefore:

$$(PB)^2 = (PE)^2 + (EB)^2 \quad \text{and} \quad (PC)^2 = (PF)^2 + (FC)^2$$

$$r^2 = x^2 + 12^2 \quad \dots\dots(i) \quad \text{and} \quad r^2 = (17 - x)^2 + 5^2 \quad \dots\dots(ii)$$


$$\Rightarrow x^2 + 12^2 = (17 - x)^2 + 5^2 \quad \dots\dots [\text{comparing (i) and (ii)}]$$

$$\Rightarrow x^2 + 144 = 289 + x^2 - 34x + 25$$

$$\Rightarrow 34x = 170$$

$$\Rightarrow x = 5$$

Now from (i), we have:



#### Key Fact:

If AB is chord and diameter CD is right bisector of AB, then:

$$AC = BC \text{ and } AD = BD$$

$$r^2 = 5^2 + 12^2 = 25 + 144 = 169$$

Taking square roots on both sides, we get  $r = 13$ .

$\therefore$  Radius of circle =  $r = 13$  cm

### Example:

Two parallel chords AB and CD of lengths 16 cm and 12 cm respectively are drawn on the opposite sides of centre O of the circle. Find the distance between the chords if the radius of the circle is 10 cm.

### Solution:

Radius of circle =  $OA = OC = 10$  cm

$AB = 16$  cm,  $CD = 12$  cm

Draw  $\overline{OE} \perp \overline{AB}$  and  $\overline{OF} \perp \overline{CD}$

$$\Rightarrow AE = \frac{1}{2} AB = 8 \text{ cm} \quad \text{and} \quad CF = \frac{1}{2} CD = 6 \text{ cm}$$

As  $\overline{AB} \parallel \overline{CD}$ , therefore points E, O and F are collinear.

Now, OAE and OCF are right angled triangles. Therefore:

$$\begin{aligned} (OE)^2 + (AE)^2 &= (OA)^2 & \text{and} & \quad (OF)^2 + (CF)^2 = (OC)^2 \\ \Rightarrow (OE)^2 + (8)^2 &= (10)^2 & \text{and} & \quad (OF)^2 + (6)^2 = (10)^2 \\ \Rightarrow (OE)^2 + 64 &= 100 & \text{and} & \quad (OF)^2 + 36 = 100 \\ \Rightarrow (OE)^2 &= 100 - 64 = 36 & \text{and} & \quad (OF)^2 = 100 - 36 = 64 \end{aligned}$$

Taking square roots, we get:


$$OE = 6 \text{ cm} \quad \text{and} \quad OF = 8 \text{ cm}$$

$$\therefore \text{Distance between chords} = OE + OF = 6 + 8 = 14 \text{ cm}$$


### Exercise 9.2

1. In the figure, O is centre of the circle and  $\angle COF = 150^\circ$ .


Find the values of  
 $\angle OCF$  and  $\angle CFD$ .



2. Two circles of radius 3 cm each are drawn with centres P and Q. AB and CD are their chords such that  $AB = CD = 4$  cm. Find shortest distances of the chords from the respective centres. Are they equal?




3. Find the distance between two chords PQ and RS of circle shown in the adjoining figure if:  
 $PQ = 6 \text{ cm}$ ,  $RS = 8 \text{ cm}$  and radius = 5 cm




4. Given that O is the centre of each of the circles. Find the values of unknown. All dimensions are in cm.

(i)




(ii)



5. Two parallel chords of lengths 24 cm and 12 cm are drawn on the opposite sides of a circle of radius 13 cm. Find the distance between the chords.

6. AB is chord of a bigger circle of radius 10 cm centered at O. Find the radius of the smaller circle passing through midpoint of AB.

Also find the difference of radii of both circles.



7. Two parallel chords of lengths 18 cm and 80 cm are drawn on the same side of a circle of radius 41 cm. Find the distance between the chords.

8. Two parallel chords AB and CD are 4 cm apart and lie on the opposite sides of the centre of a circle. If  $AB = 2 \text{ cm}$  and  $CD = 6 \text{ cm}$ , find the radius of the circle.


9. PQ and RS are two parallel chords lying on the same side of the centre of a circle and are 2 cm apart. If  $PQ = 8 \text{ cm}$  and  $RS = 10 \text{ cm}$ , find the radius of the circle.

## Arcs of Circles

### Arc of a Circle

An arc is a part of a circle. In the figure, AB is an arc of the circle P.

Arc AB is denoted by  $\widehat{AB}$ .




### Key Fact:

There is a difference between a circle and a circular region. A circular region is union of circle and its interior.

## Semicircle

A diameter divides a circle into two equal parts. Each part is called semicircle or half circle. We use three letters to name a semicircle. The first and third letter are end points of diameter.


In the figure,  $\widehat{XAY}$  is semicircle. Can you name the other semicircle?



## Minor and Major arcs

Minor arc is the arc included in a semicircle while the arc which includes a semicircle is called major arc.

In the Fig.  $\widehat{EF}$  is minor arc while  $\widehat{ECF}$  is major arc.




### Key Fact:

- (i) Major arc of a circle and semicircle arc denoted by three letters.
- (ii) Minor arc is less than and major arc is greater than the semicircle.

## Central Angle

Angle subtended by an arc at the centre of a circle is called central angle of the arc. An angle whose vertex is the centre of circle and whose arms pass through end points of arc is known as central angle of arc.



### Key Fact:

If an arc is included in a semi-circle, then its central angle is less than  $180^\circ$ . Central angle of semi-circle is straight angle ( $180^\circ$ ). Central angle of major arc of the circle is greater than  $180^\circ$ .

In the figure  $\angle AOB$  is central angle.

## Congruent Arcs

Two arcs of a circle (or of two congruent circles) are congruent if their central angles are congruent.


## Theorem 9.6

### Statement:

If two arcs of a circle (or of congruent circles) are congruent, then the corresponding chords are equal.

### **Class Activity:**

- Take two congruent arcs AB and CD of a circle with centre O.
  - Join C to D and A to B.
  - Measure chords AB and CD.
- You will see that  $AB = CD$



### **Key Fact:**

If two arcs of two congruent circles are congruent, then the corresponding chords of both circles are equal.


### **Theorem 9.7**

#### **Statement:**

If two chords of a circle (or of congruent circles) are equal, then their corresponding arcs (minor, major or semicircle) are congruent.

#### **Class Activity:**

- Draw two equal chords AB and CD of a circle with centre O.
- Measure corresponding arcs AB and CD.  
They will be congruent.



#### **Corollary:**

- If two chords of a circle (or of congruent circles) are equal, then their corresponding minor or major arcs are congruent.
- If two chords of a circle (or of congruent circles) are equal, then their corresponding semicircles are congruent.

### **Theorem 9.8**


#### **Statement:**

Equal chords of a circle (or of congruent circles) subtend equal angles at the centre (at the corresponding centres).

#### **Class Activity:**

- Draw two equal chords AB and CD of a circle with centre O.
- Join O to A, B, C and D.
- Measure corresponding angles AOB and COD.

You will observe that the corresponding angles are equal in measure.



## Theorem 9.9

### Statement:


If the angles subtended by two chords of a circle (or congruent circles) at the centre (corresponding circles) are equal, the chords are equal.

### Class Activity:

- (i) Draw two chords AB and CD of a circle such that  
 $\angle AOB = \angle COD$

- (ii) Measure corresponding chords AB and CD.  
What do you notice?

You will observe that the corresponding chords are equal in measure.



### Example:

In the figure,  $\widehat{AB} = 4\widehat{BC}$  and  $\angle AOB = 120^\circ$ , find:

- (a)  $\angle BOC$    (b)  $\angle OBC$    (c)  $\angle OBA$   
(d)  $\angle OAB + \angle ABC + \angle BCO + \angle COA$


### Solution:

(a) Let  $\angle BOC = x$ , then  $\angle AOB = 4x$  ... ( $\widehat{AB} = 4\widehat{BC}$ )

But  $\angle AOB = 120^\circ$  .... (Given)

$$\therefore 4x = 120^\circ \Rightarrow x = 30^\circ$$

Hence,  $\angle BOC = 30^\circ$



(b)  $\triangle OBC$  is an isosceles triangle. ( $OB = OC = \text{radii}$ )

$\Rightarrow \angle OBC = \angle OCB$  (opposite angles of congruent sides of triangle)

$$\therefore \angle OBC = \frac{1}{2}(180^\circ - 30^\circ) = 75^\circ$$

(c)  $\triangle OAB$  is an isosceles triangle. ( $OA = OB = \text{radii}$ )

$\Rightarrow \angle OAB = \angle OBA$  (opposite angles of congruent sides of triangle)

$$\therefore \angle OBA = \frac{1}{2}(180^\circ - 120^\circ) = 30^\circ$$

(d)  $\angle OAB + \angle ABC + \angle BCO + \angle COA = 30^\circ + (30^\circ + 75^\circ) + 75^\circ + (30^\circ + 120^\circ)$   
 $= 30^\circ + 105^\circ + 75^\circ + 150^\circ = 360^\circ$

### Example:

Diagram shows a pentagon inscribed in a circle with centre P.

If  $\widehat{AB} = \widehat{BC} = \widehat{CD}$  and  $\angle ABC = 140^\circ$ , then find:

- (a)  $\angle AEB$    (b)  $\angle AED$    (c)  $\angle BPD$

### Solution:


(a) Given that,  $\widehat{AB} = \widehat{BC} = \widehat{CD}$ , then:

Correspong chords are equal in length.

i.e.,  $AB = BC = CD$

ABCE is a cyclic quadrilateral.

$$\therefore \angle ABC + \angle AEC = 180^\circ$$



$$\begin{aligned}\Rightarrow \angle AEC &= 180^\circ - \angle ABC = 180^\circ - 140^\circ = 40^\circ \\ \Rightarrow \angle AEB + \angle BEC &= 40^\circ \text{ (angle sum postulate)} \\ \Rightarrow 2 \angle AEB &= 40^\circ \quad (\angle AEB = \angle BEC) \\ \Rightarrow \angle AEB &= 20^\circ\end{aligned}$$

(b) As,  $AB = BC = CD$  (given)

$$\angle AEB = \angle BEC = \angle CED = 20^\circ$$

(Equal chords subtend equal angle at a point on the circumference.)

$$\text{Now, } \angle AED = \angle AEB + \angle BEC + \angle CED = 20^\circ + 20^\circ + 20^\circ = 60^\circ$$

$$\begin{aligned}(c) \angle BPD &= 2 \angle BED \text{ (Angle at the centre is double the angle at a point on circumference.)} \\ &= 2 \times 40^\circ = 80^\circ\end{aligned}$$

**Example:**

In the figure, chords  $\overline{EF} \parallel \overline{GH}$ . Prove that:

$$\widehat{EG} = \widehat{FH}$$

**Solution:**

Draw  $\overline{EF}$  and  $\overline{FG}$  passing through centre of circle P.

$\angle EPG = 2 \angle EFG$  (Angle at the centre is double the angle at a point on the circumference.)


$$\therefore \angle EFG = \frac{1}{2} \angle EPG \quad (i)$$

$$\text{Similarly, } \angle FGH = \frac{1}{2} \angle FPH \quad (ii)$$

But,  $\angle EFG = \angle FGH$  (alternate angles)

$$\therefore \angle EPG = \angle FPH \quad [\text{from (i) and (ii)}]$$

Hence,  $\widehat{EG} = \widehat{FH}$  (Arcs subtending equal angles at the centre of circle are equal.)



**Key Fact:**



If two arcs of a circle are unequal, then chord and angle corresponding to greater arc are greater.

**Check Point:**

Prove that non-parallel sides of a cyclic trapezium are equal.

### Exercise 9.3


1. In the figure, chord  $AB =$  chord  $BC$ .
  - Is  $\widehat{AB} = \widehat{BC}$ ?
  - Is  $\angle APB = \angle BPC$ ?
  - If  $\widehat{AB} < \widehat{AD}$ , then what is relation between corresponding chords?
2. In the figure,  
 $\widehat{AC} = \widehat{BD}$   
 Prove that:  $\overline{AB} \parallel \overline{CD}$



3. In the given figure, chords  $\overline{AB}$  and  $\overline{CD}$  intersect at P.

If  $AB = CD$ , then prove that:

$$\widehat{AB} = \widehat{CD}$$



4. Consider two circles of radii 2 cm and 4 cm centered at P and Q respectively.  $\widehat{AB}$  and  $\widehat{CD}$  are their arcs such that  $\angle APB = \angle CQD$ .

- Are  $\widehat{AB}$  and  $\widehat{CD}$  congruent?
- What is relation between the lengths of  $\widehat{AB}$  and  $\widehat{CD}$ ?
- Show that  $\triangle APB \sim \triangle CQD$

5. O is centre of two concentric circles of radii 3 cm and 6 cm.



OA and OB are radii of larger circle and they meet smaller circle at C and D respectively.

- Are  $\overline{AB}$  and  $\overline{CD}$  parallel?
- If  $AB = 8$  cm, then find  $CD$ .
- Find distance between  $\overline{AB}$  and  $\overline{CD}$ .

6. The length of an arc of a circle is 4 cm and its central angle is of measure  $60^\circ$ . Find central angle of the arc whose length is 8 cm.

7. Given that M is a point on a circle centered at P. If M is equidistant from the two radii PE and PF, show that  $\widehat{ME} = \widehat{MF}$ .

8. Consider two circles centered at A and B respectively.  $\overline{CD}$  and  $\overline{EF}$  are their chords such that  $CD = 12$  cm,  $EF = 8$  cm and  $\angle CAD = \angle EBF$ . Find radius of circle A if radius of circle B is 4 cm.




9. In the figure,  $ABDC$  is cyclic quadrilateral such that:

$$AB = CD.$$


Prove that:

$$AC = BD$$



10. In the given figure, ABC is an isosceles triangle inscribed in a circle centered at O.

Prove that  $\overleftrightarrow{AO}$  bisects  $\angle BDC$ .




11. ABCDE is a pentagon inscribed in a circle.

If  $AB = BC = CD$ ,  $\angle BCD = 130^\circ$  and  $\angle BAE = 120^\circ$


Find:

- (a)  $\angle ABC$
- (b)  $\angle CDE$
- (c)  $\angle AED$
- (d)  $\angle EAD$



12. Given that O is the centre of a circle. Diameters  $\overline{AB}$  and  $\overline{CD}$  are perpendicular to each other.

Prove that the line segments joining in order the end points of diameters form a square.




13. In the figure, O is centre of the circle.

$AB = CD$  and  $\angle OAB = 40^\circ$ .

Find  $\angle AOB$  and  $\angle COD$ .

Are the angles equal? If yes, then why?



14. In the figure, diameters AB and CD of circle are perpendicular.


(i) Prove that all the four chords make equal central angles.

(ii) Find  $\angle PCB$  and  $\angle ACB$ .

(iii) Find other angles that are equal to  $\angle ACB$ .

(iv) What is the name of inscribed polygon?


(v) How many circles can be drawn through the vertices of polygon formed?



15. In the adjoining figure,  $RS = 3PQ$  and  $\angle PTQ + \angle RTS = 180^\circ$ .

(i) Find  $\angle PTQ$  and  $\angle RTS$ .

(ii) What is the ratio between measures of  $\angle PTQ$  and  $\angle RTS$ ?




## Applications of Chords and Arcs

### Arc Length

The length of an arc of a circle of radius 'r' that subtends an angle  $\theta$  at the center is calculated by the following formula:

$$\begin{aligned}\text{Arc length } l &= \frac{\theta}{360^\circ} \times \text{circumference of circle} \\ &= \frac{\theta}{360^\circ} \times 2\pi r\end{aligned}$$

Where  $360^\circ$  is the interior angle subtended by a circle.



### Chord Length


Length of a chord of a circle of radius 'r'

lying at a distance 'd' from the centre is:

In the adjoining figure, length of chord AB is:

$$AB = 2\sqrt{r^2 - d^2}$$

Note that AC is half of AB.



### Perimeter of a Semicircle

The perimeter of a semicircle is given by the following formula.


$$P = 2 \times \text{radius} + \text{arc length} = 2r + \pi r$$

### Perimeter of a Segment of a Circle

Perimeter of segment of a circle is defined as:

$$P = \text{arc length} + \text{chord length}$$

$$= \frac{\theta}{360^\circ} \times 2\pi r + 2\sqrt{r^2 - d^2}$$



### Area of Sector of Circle

Area of the sector of a circle having radius  $r$  and central angle  $\theta$  is:

$$A = \frac{\theta}{360^\circ} \times \pi r^2$$

Where the angle  $\theta$  is measured in degrees.

#### Example:

Find the length of arc and area of sector of a circle with central angle of  $60^\circ$  and radius 14cm.

**Solution:** Given that:

$$\theta = 60^\circ \text{ and } r = 14\text{cm}$$

$$\begin{aligned}\text{Arc length} &= \frac{\theta}{360^\circ} \times 2\pi r \\ &= \frac{60^\circ}{360^\circ} \times 2 \times \frac{22}{7} \times 14 = 14.67 \text{ cm}\end{aligned}$$

$$\text{Area} = \frac{\theta}{360^\circ} \times \pi r^2 = \frac{60^\circ}{360^\circ} \times \frac{22}{7} \times (14)^2 = 102.67 \text{ cm}^2$$

**Example:**

A circular decoration piece consists of two equal parts as shown in the figure. Find the length of one part of the piece. What is the cost of decoration piece @ Rs.8 per centimetre?

**Solution:**


Given that  $r = 56\text{cm}$ ,  $\theta = 90^\circ$ , then:

$$l = \frac{\theta}{360^\circ} \times 2\pi r = \frac{90^\circ}{360^\circ} \times 2 \times \frac{22}{7} \times 56$$

$$l = 88\text{ cm}$$

Length of one part of the decoration piece is  $= 2 \times 88 = 176\text{ cm}$

Cost of decoration piece @ Rs.8 per cm = Rs.  $2 \times 176 \times 8 = \text{Rs. } 2,816$




### Exercise 9.4

1. In the figure, radius of circle is 7cm. Find:

- (i) the length of minor arc  $\widehat{PQ}$  and major arc  $\widehat{PAQ}$ .
- (ii) the circumference of circle.


Is the sum of lengths of minor arc and major arc equal to circumference of circle?



2. Given the radius of circle is 8cm. Find:

- (i) the area of minor sector and major sector.
- (ii) the area of circle..


Is the sum of areas of minor and major sectors equal to area of circle?



3. Find the distance covered by the tip of hour hand of a clock in 5 hours if the length of hour hand is 10cm.


4. Given is the sector of circle. Find:

- (i) the length of outer minor arc.
- (ii) the length of inner minor arc.
- (iii) the difference of lengths of both arcs.
- (iv) the area of minor sector of bigger circle.
- (v) the area of minor sector of smaller circle.
- (vi) the area of shaded part of sector.



5. Radius of the given circle is 14 cm. Find:


- (i) length of  $\overline{LM}$ .
- (ii) length of  $\widehat{LM}$ .
- (iii) perimeter of segment along chord  $\overline{LM}$ .
- (iv) area of triangle OLM.
- (v) area of sector OLM.
- (vi) area of segment along chord  $\overline{LM}$ .




6. In a circle whose diameter is 12 cm, there is a central angle whose measure is  $120^\circ$ . A chord joins the endpoints of the arc cut off by the angle. Find the length of arc along the chord.

7. The diameter of a circle is 10 cm long and a chord parallel to it is 6 cm long. Find the distance between the chord and the diameter of the circle

8. Find the area and perimeter of semicircular window if its radius is 2.8 feet.



9. The building shown resembles a flying saucer that has landed on its four legs. Find the length of supporting beam AB if it makes an angle of  $98^\circ$  with center and the radius of the beam is 21 m.



10. Figure shows a bridge of semi-circular shape with arc length of 44 m.

Find the length of road constructed below.




11. Find the perimeter of lower portion (below the line PQ) of circular window if:

radius of window = 2.1 feet,


$\angle POQ = 105^\circ$ ,

and shortest distance between centre O and  $\overline{PQ} = 1.4$  feet.


Also find the area of upper portion (above line PQ) of window.



12. A circular roller coaster is shown in the figure. Find the distance covered by driver of train between two points making an angle of  $80^\circ$  with the centre of roller coaster if its radius is 21 ft.



13. Building shown in the figure is located in Guangzhou province, China. The building's height is 138 m with 33 stories. It has an empty circular core almost 59 m in diameter.
- What is the covered area of the building?
  - Find the circular length of the part of building that touches ground if it makes an angle of  $54^\circ$  with the centre.



### I have Learnt


- Solving the problems by using the properties of a circle that:
  - One and only one circle can pass through three non collinear points.
  - A straight line drawn from the centre of a circle to bisect the chord is perpendicular to the chord and vice versa.
  - Two congruent chords of a circle are equidistant from the centre and vice versa.
  - If two arcs of a circle (or of congruent circles) are congruent, then the corresponding chords are equal and vice versa.
  - Equal chords of a circle (or of congruent circles) subtend equal angles at the centre (at the corresponding centres) and vice versa.
  - Applying concepts of chords and arcs to real life word problems such as decorative features, rainbow, bridges and roller coaster track.

### MISCELLANEOUS EXERCISE-9



- Encircle the correct option in the following.
  - One and only one circle can pass through ..... non collinear points.
    - 2
    - 3
    - 4
    - 5
  - ..... number of circles can pass through a point.
    - 1
    - 2
    - 3
    - infinite
  - Diameter of circle which bisects the chord is ..... to the chord.
    - collinear
    - parallel
    - perpendicular
    - equal
  - In the figure,  $OC = 3 \text{ cm}$ ,  $AB = 8 \text{ cm}$ .

The radius of circle is:

- 4 cm
- 4.5 cm
- 5 cm
- 10 cm



- v. Diameter of circle perpendicular to the chord ..... the chord.  
 (a) intersects (b) bisects (c) trisects (d) touches
- vi. Two chords which are equidistant from the ..... are congruent.  
 (a) centre (b) diameter (c) circle (d) chord
- vii. Two ..... which are equidistant from the centre are congruent.  
 (a) circles (b) diameters (c) segments (d) chords
- viii. Length of chord of a circle of radius 5 cm is 8 cm. The distance of chord from centre is:  
 (a) 3cm (b) 4cm (c) 5cm (d) 6cm
- ix. What is the radius of circle in the figure?  
 (a) 26cm (b) 10cm (c) 5cm (d) 13cm
- x. In the figure,  $AB = 4\text{cm}$ ,  $OC = 3\text{cm}$ . Chord DE passes through mid of OC and is parallel to AB. What is the length of DE?  
 (a) 4.5 cm (b) 5.6 cm (c) 6.6 cm (d) 3.3 cm



- xi. An angle whose vertex is centre of circle and whose arms pass through end points of an arc is known as ..... angle.  
 (a) inscribed (b) central (c) interior (d) exterior
- xii. Corresponding arcs of two congruent chords of a circle are \_\_\_\_\_.  
 (a) unequal (b) major (c) congruent (d) minor
- xiii. Length of two chords of a circle are in the ratio  $1 : 3$ , and central angle of one arc is  $60^\circ$ , the second arc is \_\_\_\_\_.  
 (a) minor (b) major (c) semicircle (d) circle
- xiv. The central angle of quadrant of a circle is \_\_\_\_\_.  
 (a)  $30^\circ$  (b)  $45^\circ$  (c)  $60^\circ$  (d)  $90^\circ$
- xv. If a circle is divided into ten equal arcs, then central angle of each arc is \_\_\_\_\_.  
 (a)  $10^\circ$  (b)  $36^\circ$  (c)  $60^\circ$  (d)  $90^\circ$
- xvi. How many central angles of an arc can be drawn?  
 (a) one (b) two (c) finite (d) infinite
- xvii. Two congruent chords of two congruent circles have ..... central angles.  
 (a) different (b) same (c) proportional (d) acute
- xviii. Central angle of an arc which includes a semicircle in it is \_\_\_\_\_.  
 (a)  $< 90^\circ$  (b)  $> 90^\circ$  (c)  $< 180^\circ$  (d)  $> 180^\circ$


# Tangents and Angles of a Circle

**After studying this unit students will be able to:**

- If a line is drawn perpendicular to a radial segment of a circle at its outer end point, it is tangent to the circle.
- The tangent to a circle and the radial segment joining the point of contact and the centre are perpendicular to each other.
- The two tangents drawn to a circle from a point outside the circle are equal in length.
- If two circles touch externally (internally), the distance between their centres is equal to the sum (difference) of their radii.
- The measure of a central angle of a minor arc of a circle, is double that of the angle subtended by the corresponding major arc.
- Any two angles in the same segment of a circle are equal.
- The angle in a semi-circle is a right angle.
- The angle in a segment greater than semi-circle is less than a right angle.
- The angle in a segment less than semi-circle is greater than a right angle.
- The opposite angles of any quadrilateral inscribed in a circle are supplementary.
- Apply concepts of tangents and angles to real life world problems such as architecture, monuments and pyramids etc.

Tangents have several important applications across various fields, particularly in mathematics, physics, engineering, and geometry. Some applications of tangents are:

- Tangents play an important role in the study of curves. They are lines that touch a curve at a single point, perpendicular to the curve's radius at that point. Tangents are used to define geometric properties such as the radius, diameter, chord, and arc length of circles.
- Tangents are used to design curves in buildings, such as arches and domes.
- Tangents are used to design curves in roads, bridges, and other structures. By using tangents, engineers can create curves that are safe and efficient for travel.
- Tangents are used to analyse the motion of objects. For example, when a car or bike travels down a road, the road becomes tangent at each location where the wheels roll.



**Tangent Line**

# Tangent to a Circle

## Tangent Line

A line which touches the circle at only one point is called tangent line. A tangent line is perpendicular to the radial segment. Moreover the distance of tangent line from the centre is equal to radius of the circle.


### Theorem 10.1

#### Statement:

If a line is drawn perpendicular to a radial segment of a circle at its outer end point, it is tangent to the circle.

#### Class Activity:

- (i) Draw a radial segment  $\overline{AB}$  ( $\overrightarrow{AB}$ ) of a circle as shown in the adjoining figure.
- (ii) Draw a perpendicular line  $\overrightarrow{PQ}$  ( $\overrightarrow{PQ}$ ) at the outer end of  $\overline{AB}$ .
- (iii) Check whether  $\overrightarrow{PQ}$  touches the circle at only one point or not.  
Is  $\overrightarrow{PQ}$  tangent line?
- (iv) Take any point C on  $\overrightarrow{PQ}$  and join C to A.  
Why  $\overline{AC}$  is not the radial segment?




### Theorem 10.2

#### Statement:

The tangent to a circle and the radial segment joining the point of contact and the centre are perpendicular to each other.

#### Class Activity:

- (i) Draw a radial segment  $\overline{AB}$  ( $\overrightarrow{AB}$ ) of a circle as shown in the adjoining figure.
- (ii) Draw a tangent  $\overrightarrow{PQ}$  at point B.
- (iii) Measure the angle between  $\overline{AB}$  and tangent  $\overrightarrow{PQ}$ , i.e.  $\angle ABC$   
What is the measure of  $\angle ABC$ ?
- (v) Take any point C on  $\overrightarrow{PQ}$  and join C to A. Which is shortest  $\overline{AB}$  or  $\overline{AC}$ ? Why  $\angle ABQ = 90^\circ$  and  $\angle ACB \neq 90^\circ$ ?



#### Corollaries:

- (i) The line perpendicular to the tangent at point of contact passes through the centre of circle.
- (ii) The lines perpendicular to the diameter of a circle at its end points are tangents to the circle.

- (iii) The line perpendicular to the tangent of circle from centre passes through point of contact.
- (iv) The tangents at end points of diameter of a circle are parallel.


### Theorem 10.3

**Statement:**

The two tangents drawn to a circle from a point outside the circle are equal in length.

**Class Activity:**

- (i) Draw a circle with centre O.
- (ii) Take a point A outside the circle and join with O.
- (iii) Draw two tangents  $\overrightarrow{AB}$  and  $\overrightarrow{AC}$  where B and C are points on the circle.
- (iv) Measure  $\overrightarrow{AB}$  and  $\overrightarrow{AC}$ .  
Are they equal?



**Corollary:**

The two tangents drawn to a circle from an outer point make equal angles with the line segment joining the centre and outer point.


### Theorem 10.4

**Statement:**

If two circles touch externally, the distance between their centres is equal to the sum of their radii.

**Class Activity:**

- (i) Draw two circles of radii 2 cm and 1 cm with centres A and B respectively such that the circles touch externally as shown in the figure.
- (ii) Join the centres of both circles by drawing AB.
- (iii) Measure AB. It will be 3 cm.



Is 3 cm equal to the sum of radii of both the circles? Can we write  $AB = AC + BC$  ?

**Corollary:**

If two congruent circles touch externally, the distance between their centres is equal to diameter of each circle.

### Theorem 10.5

**Statement:**

If two circles touch internally, the distance between their centres is equal to the difference of their radii.

### Class Activity:

- (i) Draw two circles of radii 3.5 cm and 1 cm with centres A and B respectively such that the circles touch internally as shown in the figure.


- (ii) Draw AB.

- (iii) Measure AC. It will be 2.5 cm.

Is 2.5 cm equal to the difference of radii of both the circles?

Can we write  $AC = AB - BC$  ?

where AB and BC are radii of both circles.



### Corollary:

If two congruent circles touch internally, the distance between their centres is equal to zero.

### Theorem 10.6 (Alternate Segment Theorem)

#### Statement:


An angle between a tangent and chord of a circle through the point of contact is equal to the angle in the alternate segment.

### Class Activity:

- (i) Draw a tangent  $\overrightarrow{PT}$  and a chord  $\overline{PA}$  at the point of contact T of a circle centered at O.

- (ii) Measure  $\angle APT$  and note it.

- (iii) Draw angle  $\angle ABP$  in the alternate segment and measure it.



You will see that  $\angle APT = \angle ABP$


### Example:

In the figure,  $\overrightarrow{AB}$  is tangent to the circle of centre P.

- (i) Express the length AP in terms of r.

- (ii) What is measure of  $\angle ABP$ ? and why?

- (iii) What is the radius of the circle?



### Solution:

- (i)  $AP = AC + PC = (9 + r) \text{ cm}$

- (ii)  $\angle ABP = 90^\circ$  because tangent line and radial segment are perpendicular to each other.

- (iii)  $\triangle ABP$  is right angled triangle, therefore using Pythagoras theorem, we have:

$$\begin{aligned} (AP)^2 &= (AB)^2 + (BP)^2 \\ \Rightarrow (9+r)^2 &= (15)^2 + r^2 \quad \Rightarrow 81 + r^2 + 18r = 225 + r^2 \\ \Rightarrow 18r &= 225 - 81 = 144 \quad \Rightarrow r = 144 \div 18 = 8 \\ \therefore \text{Radius of the circle} &= 8 \text{ cm} \end{aligned}$$

### Example:

In the figure,  $\overline{OP}$  and  $\overline{OQ}$  are tangents to a circle

centered at S. If  $\angle POQ = 40^\circ$ , then find:

- (i)  $\angle PSQ$  (ii)  $\angle SPQ$  (iii)  $\angle OPQ$

**Solution:**

- (i) Join O and S.

As OPS is right angled triangle, therefore:

$$\angle OPS = 90^\circ$$

Also  $\angle POS = 20^\circ$  (Half of  $40^\circ$ .  $\overline{OS}$  is angle bisector of  $\angle POQ$ .)

$$\therefore \angle PSO = 90^\circ - 20^\circ = 70^\circ$$

$$\text{Hence, } \angle PSQ = \angle PSO + \angle QSO = 70^\circ + 70^\circ = 140^\circ \quad (\angle PSO = \angle QSO)$$

- (ii) In  $\triangle PQS$ ,  $PS = QS$  (Radii of same circle)

$\therefore \angle SPQ = \angle SQP$  (Opposite angles of congruent sides of triangle)

$$\text{Let } \angle SPQ = \angle SQP = x$$

$$\text{Then, } x + x + \angle PSQ = 180^\circ \Rightarrow 2x + 140^\circ = 180^\circ$$

$$\Rightarrow 2x = 180^\circ - 140^\circ = 40^\circ \Rightarrow x = 20^\circ$$

$$\text{Hence, } \angle SPQ = \angle SQP = 20^\circ$$

- (iii)  $\angle OPQ = \angle SPO - \angle SPQ = 90^\circ - 20^\circ = 70^\circ$

**Example:** Find the measures of angles  $a$  and  $b$  in the adjoining figure.

**Solution:**

As  $\overline{OQ}$  and  $\overline{OR}$  are tangents, therefore:

$$\angle ORP = \angle OQP = 90^\circ$$

Now  $OQPR$  is a kite.

$$\therefore 45^\circ + 90^\circ + a + 90^\circ = 360^\circ$$

$$\Rightarrow a = 360^\circ - 90^\circ - 90^\circ - 45^\circ = 135^\circ$$

As  $\overline{RS}$  is a straight line and P lies on it.

$$\therefore a + \angle QPS = 180^\circ \Rightarrow 135^\circ + \angle QPS = 180^\circ$$



$$\Rightarrow \angle QPS = 180^\circ - 135^\circ = 45^\circ$$

- In  $\triangle PQS$ ,  $PQ = PS$  (Radii of same circle)

$\therefore \angle PQS = \angle PSQ$  (Opposite angles of congruent sides of triangle)

$$\text{Then, } b + b + 45^\circ = 180^\circ \Rightarrow 2b + 45^\circ = 180^\circ$$

$$\Rightarrow 2b = 180^\circ - 45^\circ = 135^\circ \Rightarrow b = 67.5^\circ$$



#### Check Point:

The distance of a point P from the centre O of a circle of radius 6 cm, is 10 cm. Find the length of tangents to the circle from P.

**Example:** Find the measures of angles  $x, y$  and  $z$  in the adjoining figure.

**Solution:**

We know that:


$$\angle BDC = \angle BCA \text{ (Theorem 10.6)}$$

$$\therefore y = 60^\circ$$

In triangle BCD:

$$56^\circ + y + z = 180^\circ$$

$$\Rightarrow 56^\circ + 60^\circ + z = 180^\circ$$



$$\Rightarrow z = 180^\circ - 56^\circ - 60^\circ = 64^\circ$$









Now,

$$x = \angle CBD \text{ (Theorem 10.6)}$$

$$\Rightarrow x = 56^\circ$$


### Exercise 10.1

1. Find the values of unknown in the following figures.




2. In the figure AB and AC are two tangents of a circle with centre O at point B and C respectively. If  $\angle BOC = 120^\circ$ , then find:

- (i)  $\angle OBA$       (ii)  $\angle OCB$   
 (iii)  $\angle BAC$       (iv)  $\angle ABC$




3. O is centre of two concentric circles. AB and CD are chords of outer circle tangent to inner circle. Prove that AB = CD.




4. Ali is whirling a stone tied with a string of length 3 feet. All of a sudden string is broken and the stone moves along tangent direction of circular path and hits a point 5 feet away from Ali. Find the distance covered by the stone.


5. Three circles touch externally. The ratio of distances between centers is  $2 : 3 : 4$ . The perimeter of triangle joining the centers is 36 cm. Find radius of each circle.




6. In the figure, AD = 5 cm, CD = 4 cm and BD = 4.5 cm. Find the distance between centres of circles.



7. In the adjoining figure,  $\angle PEF = 20^\circ$ . H is mid-point of  $\overline{FG}$ . Find  $\angle PGH$ .



8. In the figure, triangle ABC, AB = AC.  
Find the values of angles  $x$  and  $y$ .




## Angle in a Segment of a Circle

### Segment of the Circle

A chord of a circle divides the circular region into two parts, called segment of the circle.


The region between chord and minor arc is called minor segment and the region between chord and major arc is called major segment. In the figure, the shaded region is minor segment and remaining region is major segment.



### Angle in a Segment

An angle whose vertex is a point on the arc of a circle and whose arms pass through end points of chord, is known as inscribed angle of the arc.

In the adjoining figure, angles  $a$  and  $b$  are examples of angles in the segment.



#### Key Fact:

Major segment is greater than semi-circular region and always contains the centre of the circle while minor segment is smaller than the semi-circular region.

## Theorem 10.7


### Statement:

The measure of a central angle of a minor arc of a circle, is double that of the angle subtended by the corresponding major arc.

### Class Activity:

- Draw the circle O and take points E, F and G on the same segment of the circle.
- Measure angles E, F, G and central angle COD.
- Check whether:

$$2\angle E = 2\angle F = 2\angle G = \angle COD$$




Prove theorem 10.7 by taking central angle  $\angle COD = 180^\circ$  and  $\angle COD > 180^\circ$  (reflex angle).

### Theorem 10.8

**Statement:** Any two angles in the same segment of a circle are equal.

**Class Activity:**

- Draw two circles of different radii.
- Draw a chord AB in both the circles.
- Take three points C, D and E on the same segment and join them with end points of chord AB.



- Complete the following table.

S. No	$\angle C$	$\angle D$	$\angle E$	Result
1.				
2.				

Are  $\angle C = \angle D = \angle E$  ?

**Result:**

Any number of angles in the same segment of a circle are equal.

### Theorem 10.9

**Statement:**


The angle in a semi-circle is a right angle.

**Class Activity:**

- Draw a semicircle O.
- Take a point P on the semi-circle and draw AP and PB.
- Measure angle APB.

What do you notice?

You will see that angle APB is equal to  $90^\circ$ .



### **Corollary:**

If an arc of a circle subtends a right angle at any point of the remaining part of the circle, then the arc is a semi-circle.

### **Theorem 10.10**


#### **Statement:**

The angle in a segment greater than semi-circle is less than a right angle.

#### **Class Activity:**

- Draw a circle O.
- Draw a chord AB such that the centre lies in the major segment.
- Take three points P, Q and R on the major segment and join them with end points of chord AB.
- Measure  $\angle P = \angle Q = \angle R$

You will see that angles P, Q, R are less than right angles (acute angles).



### **Theorem 10.11**


#### **Statement:**

The angle in a segment less than semi-circle is greater than a right angle.

#### **Class Activity:**

- Draw a circle O.
- Draw a chord AB such that the centre lies in the major segment.
- Take three points P, Q and R on the minor segment and join them with end points of chord AB.
- Measure  $\angle P = \angle Q = \angle R$

You will see that the angles P, Q, R are greater than right angle (obtuse angle)




### **Cyclic Quadrilateral**

A quadrilateral inscribed in a circle is called cyclic quadrilateral.

Or a quadrilateral having vertices on the circle is called a cyclic quadrilateral.

In the adjoining figure, ABCD is cyclic quadrilateral.

The four points on the circle are called concyclic.



### Theorem 10.12


#### Statement:

The opposite angles of any quadrilateral inscribed in a circle are supplementary. Or  
The sum of opposite angles in a cyclic quadrilateral is  $180^\circ$ .

#### Class Activity:

- Draw a circle O and inscribe a quadrilateral ABCD in the circle as shown in the adjoining figure.
- Measure interior angles A, B, C and D of quadrilateral.
- Add  $\angle A$  and  $\angle C$ .  
What is the sum of these angles? Yes! it is  $180^\circ$ .
- Again add  $\angle B$  and  $\angle D$ .  
What is the sum of these angles? Yes! it is  $180^\circ$ .

What is sum of opposite interior angles of cyclic quadrilateral?



#### Example:

In the figure,  $\overline{OP}$  and  $\overline{OQ}$  are tangents to a circle centered at S. If  $\angle POQ = 36^\circ$ , then find:

- $\angle PSQ$
- $\angle SQP$
- $\angle PTQ$

#### Solution:

- Join O and S.

As OPS is right angled triangle, therefore:

$$\angle OPS = 90^\circ$$

Also  $\angle POS = 18^\circ$  (Half of  $36^\circ$ .  $\overline{OS}$  is angle bisector of  $\angle POQ$ .)

$$\therefore \angle PSO = 90^\circ - 18^\circ = 72^\circ$$

$$\text{Hence, } \angle PSQ = \angle PSO + \angle QSO = 72^\circ + 72^\circ = 144^\circ \quad (\angle PSO = \angle QSO)$$

- In  $\Delta PQS$ ,  $PS = QS$  (Radii of same circle)

$\therefore \angle SPQ = \angle SQP$  (Opposite angles of congruent sides of triangles)

$$\text{Let } \angle SPQ = \angle SQP = x$$

$$\text{Then, } x + x + \angle PSQ = 180^\circ \Rightarrow 2x + 144^\circ = 180^\circ$$

$$\Rightarrow 2x = 180^\circ - 144^\circ = 36^\circ \Rightarrow x = 18^\circ$$

$$\text{Hence, } \angle SPQ = \angle SQP = 18^\circ$$

- As, the measure of a central angle of a minor arc of a circle, is double that of the angle subtended by the corresponding major arc.



$$\therefore \angle PTQ = \frac{1}{2}(\angle PSQ) = \frac{1}{2} \times 144^\circ = 72^\circ$$

#### Example:

In the figure, A, B, C and D are points on a circle.

If  $\angle POQ = 100^\circ$ ,  $\angle QRS = 120^\circ$ , then find:

- $\angle PSQ$
- $\angle OPQ$
- $\angle SPO$



**Solution:**

(i)  $\angle POQ = 100^\circ$  (Central angle)

As,  $2\angle PSQ$  is angle subtended by the corresponding major arc.

$$\therefore 2\angle PSQ = \angle POQ = 100^\circ$$

$$\Rightarrow \angle PSQ = 50^\circ$$

(ii)  $\Delta PQS$  is isosceles with  $OP = OQ =$  radii. Therefore:

$$\angle OPQ + \angle OQP + \angle POQ = 180^\circ$$

$$\Rightarrow 2\angle OPQ + 100^\circ = 180^\circ \quad (\angle OPQ = \angle OQP)$$

$$\Rightarrow 2\angle OPQ = 180^\circ - 100^\circ = 80^\circ$$

$$\Rightarrow \angle OPQ = 80^\circ \div 2 = 40^\circ$$

(iii) PQRS is a cyclic quadrilateral. Therefore:

$$\angle SPQ + \angle SRQ = 180^\circ \text{ (Opposite angles in cyclic quadrilateral PQRS.)}$$


$$\Rightarrow \angle SPQ + 120^\circ = 180^\circ \Rightarrow \angle SPQ = 180^\circ - 120^\circ = 60^\circ$$

$$\text{Now, } \angle SPO + \angle OPQ = \angle SPQ \Rightarrow \angle SPO + 40^\circ = 60^\circ$$

$$\Rightarrow \angle SPO = 60^\circ - 40^\circ = 20^\circ$$

**Key Fact:**

A cyclic trapezium is isosceles and its diagonal bisect each other.

**Example:**

In the figure, A, B, C, D and E lie on the circumference of a circle.

If  $\angle ABD = 46^\circ$  and  $\angle ACB = 70^\circ$  and  $\angle EAB = 50^\circ$ , then find:

- (i)  $\angle DCA$  (ii)  $\angle AEB$

**Solution:**

(i)  $\angle ADB = \angle ACB = 70^\circ$  (angles in same segments are equal.)

In  $\Delta ABD$ :

$$\angle ABD + \angle ADB + \angle BAD = 180^\circ$$

$$\Rightarrow 46^\circ + 70^\circ + \angle BAD = 180^\circ \Rightarrow \angle BAD = 180^\circ - 46^\circ - 70^\circ = 64^\circ$$


$\angle BAD + \angle BCD = 180^\circ$  (Opposite angles in cyclic quadrilateral ABCD.)

$$64^\circ + \angle BCD = 180^\circ \Rightarrow 64^\circ + (70^\circ + \angle DCA) = 180^\circ$$


$$\Rightarrow \angle DCA = 180^\circ - 64^\circ - 70^\circ = 46^\circ$$

(ii)  $\angle AEB + \angle ACB = 180^\circ$  (Opposite angles in cyclic quadrilateral AEBC.)


$$\angle AEB + 70^\circ = 180^\circ \Rightarrow \angle AEB = 180^\circ - 70^\circ = 110^\circ$$

**Exercise 10.2**


1. In the adjoining figure  $\angle P = 30^\circ$ .  
Find values of  $\angle Q$  and  $\angle ROS$ .




2. In the adjoining figure  $\angle BAO = 30^\circ$ .  
Find  $\angle ABO$ ,  $\angle AOB$ ,  $\angle ACB$ .




3. In the adjoining figure  $\angle BAC = 40^\circ$  and  $\angle ACD = 60^\circ$ . Find  $\angle ABC$ ,  $\angle BCA$ ,  $\angle CAD$ .




4. In the figure, AOB is equilateral triangle.  
Find the measures of  $\angle O$  and  $\angle C$ .



5. In the figure,  $\overline{AB}$  is parallel to  $\overline{CD}$ .  
Find the values of  $\angle AEB$ ,  $\angle B$ ,  $\angle C$  and  $\angle D$ .




6. In the figure,  $\overline{BE}$  is diameter and  $\overline{CD}$  is tangent touching the circle at B. If  $x = 30^\circ$ , then find:  
 (i) radius of circle  
 (ii)  $\angle ABE$   
 (iii)  $\angle ABC$   
 (iv)  $\angle EBD$




7. In the figure, ABCD is a square having side 10 cm. Triangle BOP is enlarged as triangle BCQ. Find:

- (i) Scale factor of enlargement if  $OP = 2$  cm.
- (ii)  $CQ$
- (iii)  $\angle BQC$  and  $\angle BPO$
- (iv)  $AO$
- (v)  $AR$




8. In the adjoining figure,  $\angle QPR = 60^\circ$ . Find:

- (i)  $\angle QSR$
- (ii)  $\angle PRS$
- (iii)  $\angle PQS$
- (iv)  $\angle QTR$
- (v) Why  $\angle QTR$  is acute angle?




9. In the given figure, P is the centre of the circle and  $\angle BPD = 130^\circ$ . Find:

- (i)  $\angle BAD$
- (ii)  $\angle BCD$
- (iii)  $\angle ABC + \angle ADC$




10. In the given figure,  $\overline{PQ} \parallel \overline{SR}$  and  $\angle QPS = 95^\circ$ . Find:

- (i)  $\angle QRS$
- (ii)  $\angle PQR$
- (iii)  $\angle PSR$




11. In the adjoining figure,  $\angle ADC = 120^\circ$ .

- (i) Find  $\angle BAC$ .
- (ii) Find  $\angle CAD$  if  $AD = CD$ .




12. In the figure,  $\angle ABE = 95^\circ$  and  $\angle BAE = 65^\circ$ .  
 Find (i)  $\angle ECD$  (ii)  $\angle CDE$  (iii)  $\angle CED$




13. A circular wall clock of radius 1 ft is hung by a 4 feet long string with a nail at P.

- (i) Find distance between centre of clock and nail.  
 (ii) How far above the clock is the nail?



14. A wooden wheel moving on the ground is 99 cm away from a point A. Find diameter of the wheel if the distance between centre of wheel and point A is 101 cm. How much distance does the wheel cover in one round?



### I have Learnt

- ❖ If a line is drawn perpendicular to a radial segment of a circle at its outer end point, it is tangent to the circle.
- ❖ The tangent to a circle and the radial segment joining the point of contact and the centre are perpendicular to each other.
- ❖ The two tangents drawn to a circle from a point outside the circle are equal in length.
- ❖ If two circles touch externally (internally), the distance between their centres is equal to the sum (difference) of their radii.

- ❖ The measure of a central angle of a minor arc of a circle, is double that of the angle subtended by the corresponding major arc.
- ❖ Any two angles in the same segment of a circle are equal.
- ❖ The angle in a semi-circle is a right angle.
- ❖ The angle in a segment greater than semi-circle is less than a right angle.
- ❖ The angle in a segment less than semi-circle is greater than a right angle.
- ❖ The opposite angles of any quadrilateral inscribed in a circle are supplementary.

### MISCELLANEOUS EXERCISE-10

1. Encircle the correct option in the following.

- i. A tangent line touches the circle at ..... point(s).
  - (a) 1
  - (b) 2
  - (c) 3
  - (d) 4
- ii. A tangent line is ..... to radial segment.
  - (a) parallel
  - (b) equal
  - (c) perpendicular
  - (d) similar
- iii. How many tangents can be drawn on the circle from a point outside the circle?
  - (a) 1
  - (b) 2
  - (c) 3
  - (d) infinite
- iv. If two tangents are drawn on both ends of diameter of a circle, they are:
  - (a) perpendicular
  - (b) parallel
  - (c) intersecting
  - (d) none
- v. Given that radius of a circle is 4 cm. The distance between two tangents drawn at the outer ends of diameter is:
  - (a) 2 cm
  - (b) 4 cm
  - (c) 6 cm
  - (d) 8 cm
- vi. How many tangents can be drawn on the circle from a point on the circle?
  - (a) 1
  - (b) 2
  - (c) none
  - (d) infinite
- vii. Two tangents drawn from a point outside the circle are:
  - (a) perpendicular
  - (b) parallel
  - (c) not congruent
  - (d) congruent
- viii. If two circles touch externally, the distance between their centres is equal to sum of ..... of both circle.
  - (a) radii
  - (b) diameters
  - (c) circumferences
  - (d) area
- ix. If two congruent circles touch externally, the distance between their centres is equal to ..... of a circle.
  - (a) radius
  - (b) diameter
  - (c) chord
  - (d) sector
- x. If two circles of radii 1.4cm and 2.5 cm touch internally, the distance between their centres is equal to:
  - (a) 1.4 cm
  - (b) 2.5 cm
  - (c) 1.1 cm
  - (d) 3.9 cm
- xi. Angle subtended by an arc at centre of circle is called ..... angle.
  - (a) reflex
  - (b) inscribed
  - (c) straight
  - (d) central
- xii. An angle inscribed in a half circle is:
  - (a)  $0^\circ$
  - (b)  $45^\circ$
  - (c)  $90^\circ$
  - (d)  $180^\circ$

xiii. If central angle of minor arc of a circle is  $100^\circ$ , angle inscribed in corresponding major arc is:

- (a)  $200^\circ$       (b)  $100^\circ$       (c)  $75^\circ$       (d)  $50^\circ$

xiv. Central angle of minor arc of a circle is:

- (a) less than  $360^\circ$       (b) less than  $180^\circ$   
(c) greater than  $360^\circ$       (d) greater than  $180^\circ$

xv. Central angle of major arc of a circle is:

- (a) less than  $90^\circ$       (b) less than  $180^\circ$   
(c) greater than  $90^\circ$       (d) greater than  $180^\circ$

xvi. All angles in a same segment of a circle are:

- (a) equal      (b) acute      (c) obtuse      (d) supplementary

xvii. If ABCD is a cyclic quadrilateral and  $\angle A = 60^\circ$ , then  $\angle C = \dots$

- (a)  $180^\circ$       (b)  $150^\circ$       (c)  $120^\circ$       (d)  $60^\circ$

xviii. An exterior angle of a cyclic quadrilateral is ..... the opposite interior angle.


- (a) greater than      (b) less than      (c) equal to      (d) supplement of

xix. Inscribed angle of a quadrant of a circle is:


- (a)  $45^\circ$       (b)  $90^\circ$       (c)  $145^\circ$       (d)  $130^\circ$

xx. In the figure, angle  $\theta$  is equal to:

- (a)  $45^\circ$       (b)  $94^\circ$   
(c)  $23.5^\circ$       (d)  $47^\circ$



2. In a circle whose diameter is 12 cm, there is a central angle whose measure is  $90^\circ$ . A chord joins the endpoints of the arc cut off by the angle. Find the length of the chord.
3. The diameter of a circle is 20 cm long and a chord parallel to it is 12 cm long. Find the distance between the chord and the center of the circle.
4. In a cyclic quadrilateral ABCD with sides  $a = AB$ ,  $b = BC$ ,  $c = CD$ , and  $d = DA$ , and diagonals  $p = AC$  and  $q = BD$ . We can express diagonals in terms of the sides as:  
$$p \times q = (a \times c) + (b \times d)$$
.  
Find  $q$  if  $a = 4\text{cm}$ ,  $b = 5\text{cm}$ ,  $c = 6\text{ cm}$ ,  $d = 3\text{ cm}$  and  $p = 8\text{ cm}$ .



# 11

## Practical Geometry of Circles

**After studying this unit students will be able to:**

- Locate the centre of a given circle.
- Draw a circle passing through three given non-collinear points.
- Complete the circle (i) by finding its centre, (ii) without finding its centre, when a part of its circumference is given by
- Draw a tangent to a given arc, without using the centre, through a given point P when:  
(i) P is middle point of the arc (ii) P is at the end of arc (iii) P is outside the arc.
- Draw a tangent to a given circle from a point P when:  
(i) P is on the circumference (ii) P is outside the circle.
- Draw two tangents to a circle meeting each other at a given angle.
- Draw (i) direct common tangents or external tangents (ii) transverse common tangents or internal tangents, to two equal/unequal circles.
- Draw a tangent to two unequal (i) touching circles (ii) intersecting circles.
- Apply concepts of practical geometry of circles to real life word problems such as athletic tracks, recreational parks, Ferris wheels and mechanical machines.

A circle is symmetric about its centre and any of its diameters. Architects often use the circle's symmetrical properties when designing athletic tracks, recreational parks, buildings, roundabouts, Ferris-wheels, etc. Artists and painters find the circle almost indispensable in their work. Circular cylinders are used to print newspapers. Engineers exploit the circle's symmetrical properties as seen by the use of the circle in making watches, clocks, bicycles, cars, trains, ships, aero-planes, radios, telephones, trolleys, wheel-barrows, air-conditioners, rockets etc. Civilizations have progressed dramatically because of the invention of the circular wheel.



# Construction of Circles


## To Locate the Centre of a Circle

**Given:** A circle without centre.

**Required:** To locate the center of circle.

### Steps of Construction:

- (i) Take three points A, B and C on a circle.
- (ii) Join these points to obtain a triangle ABC.
- (iii) Draw right bisectors of sides AB, BC and AC which meet each other at point O.  
O is required center of circle.




## To Draw a Circle Passing Through Three Non-Collinear Points

**Given:** Three non-collinear points A, B and C.

**Required:** To draw a circle passing through these points.

### Steps of Construction:

- (i) Take three non-collinear points A, B and C on a paper.
- (ii) Join these points to obtain a triangle ABC.
- (iii) Draw right bisectors of sides BC and AC which meet each other at point O.
- (iv) With center O, construct a circle of radius OA or OB.



## To Complete the Circle when a Part of its Circumference is Given:

**(a) By finding the Centre (b) Without finding the Centre**


### (a) To Complete a Circle by Finding its Centre when Part of Circumference is Given

**Given:** A part of circle i.e. an arc.

**Required:** To complete the circle by finding its center.

### Steps of Construction:

- (i) Take three points A, B and C on the given arc.
- (ii) Join these points to obtain a triangle ABC.
- (iii) Draw right bisectors of sides AB, BC and AC which meet each other at point O.
- (iv) With center O, construct a circle of radius OA or OB of which the given arc is a part.



## (b) To Complete a Circle without Finding its Centre when Part of Circumference is Given


Given: A part of circle i.e. an arc.

Required: To complete the circle without finding its center.

### First Method

#### Steps of Construction:


- (i) Take three points A, B and C on the given arc such that  $AB = BC$ .  
(We can also  $AB \neq BC$ .)
- (ii) Join these points to obtain a triangle ABC.
- (iii) Draw  $BD = AC$  and  $CD = BC$  to obtain a triangle BCD.
- (iv) Again draw  $CE = BD$  and  $DE = CD$  to obtain a triangle CDE.
- (v) Continue in this way and construct triangles DEF, EFG, FGH and GHA.
- (vi) Join points C to D, D to E, E to F, F to G, G to H and H to A in circular path to obtain required circle.




### Second Method

#### Steps of Construction:


- (i) Draw a chord AB of suitable length on an arc PQ.
- (ii) Construct an internal angle of measure  $120^\circ$  at point B and draw BC equal to AB.
- (iii) Again, construct an internal angle of measure  $120^\circ$  at point C and draw CD equal to BC.
- (iv) Continue this procedure to obtain points E and F.
- (v) Draw arcs BC, CD, DE, EF and FA carefully such that length of these arcs is equal to the arc AB.



1. Given a circle, locate its centre.
2. Three points X, Y and Z are located on an arc. Find the centre of that arc.
3. Three non-collinear points A, B and C are such that  $AB = BC = 3\text{cm}$  and  $\angle B = 120^\circ$ . Construct a circle passing through A, B and C.
4. PQ and QR are perpendicular segments in a plane such that  $PQ = 4\text{cm}$  and  $QR = 3\text{cm}$ . Construct a circle passing through P, Q and R. What is the radius of that circle?
5. Draw a circle with the help of any round object, then find its centre.
6. Find the centre of an arc DEF, such that chord DE = 3.5cm, chord EF = 4.2cm and angle between both segments is  $60^\circ$ .
7. Draw an arc of any length. Complete the circle without finding its centre of which the given arc is the part.
8. An engineer is constructing a circular path in a recreational park as shown below. He wants to plant a pine tree in the centre of the circular path.
  - (i) How will he locate the centre of the path?
  - (ii) What is the radius of the circular path?



9. A factory produces arc shaped tiles of different colors each of inner radius 7 inches. Arc length of each tile is 2 inches. Hamza wants to purchase these tiles for making a circular design on the wall. How many tiles are required?
10. In the figure a boat swing is shown. Trace the circular path of swing during a complete round.




## Tangent to the Circle

**To draw a Tangent to a Given Arc without using the Centre, through a Given Point P when P is:**

- (a) The middle point of the arc.
- (b) At the end of the arc.
- (c) Outside the arc.

**(a) Given:** An arc PA without centre.

**Required:** To draw tangent at mid-point of arc.



### **Steps of Construction:**


- Draw an arc PA.
- Joint P to A.
- Draw right bisector EF of PA where E is mid-point of  $\overline{PA}$  and F is a point on given arc.
- Draw a line XY passing through F parallel to  $\overline{PA}$  using set square or by any other method.  
 $\overrightarrow{XY}$  is required tangent.

(b) Given: An arc PQ without centre.

Required: To draw a tangent at end point P of the arc.

### **Steps of Construction:**

- Draw an arc PQ.
- Join P to Q.
- Draw right bisector EF of PQ where E is mid-point of  $\overline{PQ}$  and F is a point on given arc.
- Join point P to F.
- With centre P, draw an arc of radius PE intersecting PF at point H.
- Take point G on this arc such that  $EH = HG$ . Now join P to G.  
 $\overrightarrow{PG}$  is required tangent.




(c) Given: An arc AB without knowing its centre.

Required: To draw a tangent from a point P where P is outside the arc.

### **Steps of Construction:**

- Draw an arc AB.
- Draw PB intersecting the arc AB at point C.
- Bisect PB at point M.
- With centre M, draw a semi-circle of radius MP or MB.
- Draw CD perpendicular to PB intersecting semicircle at point D.
- With centre P, draw another arc of radius PD intersecting given arc AB at point E.
- Joint P to E.  $\overrightarrow{PE}$  is required tangent.



### **To draw a Tangent to a Given Circle from a Point P when:**


- P lies on the Circumference.
- P lies outside the Circle.

**(a) Given:** A circle of radius 3 cm.

**Required:** To draw a tangent from a point P lying on the circle.

**Steps of Construction:**

- (i) Construct a circle of radius 3 cm by taking O as centre.
- (ii) Take a point P on the circle and join point O to point P.
- (iii) Construct an angle of  $90^\circ$  at point P and draw a line PQ.  
The line PQ is required tangent.




**(b) Given:** A circle of radius 3 cm and a point P 8 cm away from the centre of circle.

**Required:** To draw a tangent from a point P at a distance of 8 cm from the center of the circle.

**Steps of Construction:**


- (i) Construct a circle of radius 3 cm by taking O as centre.
- (ii) Take a point P at a distance of 8 cm from the centre.
- (iii) Bisect  $\overline{OP}$  at point Q.
- (iv) Draw a dotted circle of radius OQ or PQ.
- (v) The two circles intersect each other at points A and B.
- (vi) Draw  $\overrightarrow{PA}$  and  $\overrightarrow{PB}$  which are required tangents.



### To Draw two Tangents to a Circle Meeting each other at a Given Angle

**Given:** A circle of radius 3 cm and angle between tangents is  $30^\circ$ .

**Required:** To draw two tangents to the circle.



**Steps of Construction:**


- (i) Construct a circle of radius 3 cm at point P.
- (ii) Draw a diameter AB.
- (iii) Draw a tangent BX to the circle at point B.

- (iv) Construct  $\angle APC = 30^\circ$  and draw another tangent  $\overline{CD}$  at point C intersecting the first tangent at point D.  
 (v) Measure  $\angle BDC$ . It will be  $30^\circ$ .

### To draw Direct Common Tangents (External Tangents) to two Equal Circles

**Given:** Two equal circles each of radius 2.5 cm and distance between centers of circles is 6 cm.

**Required:** To draw direct common tangents or external tangents.




### Steps of Construction:

- Draw  $\overline{AB} = 6 \text{ cm}$ .
  - Construct two circles each of radius 2.5 cm at points A and B.
  - Draw diameters CD at point B and EF at point A, perpendicular to  $\overline{AB}$ .
  - Join E to D and F to C.
- Lines ED and FC are required direct common tangents.

### To draw Transverse Common Tangents (Internal Tangents) to two Equal Circles

**Given:** Two equal circles each of radius 3 cm and distance between centers of circles is 7 cm.

**Required:** To draw transverse common tangents or internal tangents.



### Steps of Construction:


- Draw  $\overline{AB} = 7 \text{ cm}$ .
- Construct two circles each of radius 3 cm at points A and B.
- Bisect  $\overline{AB}$  at point C.

- (iv) Further bisect  $\overline{AC}$  at point D and  $\overline{BC}$  at point E.
- (v) With centre D, construct a dotted circle of radius AD, intersecting the circle with centre A at points F and I.
- (vi) With centre E, construct another dotted circle of radius BE, intersecting the circle with centre B at points H and G.
- (vii) Join H to I and F to G.  
 $\overleftrightarrow{FG}$  and  $\overleftrightarrow{HI}$  are the required transverse common tangents.

### To draw Direct Common Tangents to two Unequal Circles

**Given:** Two circles of radii 2 cm and 1.2 cm respectively and distance between centers of circles is 5 cm.

**Required:** To draw direct common tangents or external tangents.




### Steps of Construction:

- (i) Draw  $AB = 5\text{cm}$ .
- (ii) Construct two circles of radii 2 cm and 1.2 cm respectively at ends A and B.
- (iii) With center as A, construct a circle of radius 0.8 cm (difference of radii of given circles).
- (iv) Bisect  $\overline{AB}$  at M and construct a circle of radius AM which intersects circle of radius 0.8cm at points C and D.
- (v) Produce  $\overline{AC}$  to point E and  $\overline{AD}$  to point F.
- (vi) Draw  $\overline{BG} \parallel \overline{AE}$  and  $\overline{BH} \parallel \overline{AF}$ .
- (vii) Join E to G and F to H.  
 $\overrightarrow{EG}$  and  $\overrightarrow{FH}$  are required direct common tangents.

### To draw Transverse Common Tangents to two Unequal Circles

**Given:** Two circles of radii 2 cm and 1.3 cm respectively and distance between centers of circles is 5 cm.

**Required:** To draw transverse common tangents



**Steps of Construction:**

- (i) Draw  $AB = 5\text{cm}$ .
- (ii) Construct two circles of radii 2 cm and 1.3 cm respectively at ends A and B.
- (iii) With center as A, construct a circle of radius 3.3 cm (sum of radii of given circles).
- (iv) Bisect  $\overline{AB}$  at M and construct a circle of radius AM which intersects circle of radius 3.3 cm at points C and D.
- (v) Draw  $\overline{AC}$  intersecting circle of radius 2 cm at E.
- (vi) Draw  $\overline{AD}$  intersecting circle of radius 2 cm at F.
- (vii) Draw  $\overline{BG} \parallel \overline{AE}$  and  $\overline{BH} \parallel \overline{AF}$ .
- (viii) Join E to G and F to H.

$\overrightarrow{EG}$  and  $\overrightarrow{FH}$  are required transverse common tangents.


**To draw a Tangent to two Unequal Touching Circles**

**Given:** Two touching circles of radius 2.5 cm and 2 cm.

**Required:** To draw a tangent to the circles.

**Steps of Construction:**

- (i) Construct a circle of radius 2.5 cm at point A and a circle of radius 2 cm at point B in such a way that both circles touch each other externally at point C.
- (ii) Join A to B.
- (iii) Bisect  $\overline{AB}$  at point M.
- (iv) With centre M, draw a dotted semi-circle of radius AM or BM.
- (v) Draw a perpendicular  $\overline{CD}$  on  $\overline{AB}$  where D is a point on the semi-circle.
- (vi) With centre D, draw an arc of radius CD intersecting the given circles at points E and F respectively.
- (vii) Join E to F.



Line EF is the required tangent.

**Class Activity:**

Draw a common tangent to two circles that are:

- (i) touching internally,
- (ii) touching externally,

such that tangent drawn is perpendicular to the radial segments of both circles.


**To draw a Tangent to two Unequal Intersecting Circles**

**Given:** Two intersecting circles of radius 2.5 cm and 3 cm.

**Required:** To draw a tangent to the circles.

**Steps of Construction:**


- (i) Construct a circle of radius 2.5 cm at point A.  
Construct another circle of radius 3 cm at point B in such a way that both circles intersect each other at points C and D.
- (ii) Join A to B.
- (iii) Bisect  $\overline{AB}$  at point M. With centre M, draw a dotted semi-circle of radius AM or MB.
- (iv) Join point D to C and produce it to point E lying on semi-circle.  $\overline{DE}$  intersects  $\overline{AB}$  at point F.
- (v) With centre E, draw an arc of radius EF intersecting the given circles at points G and H respectively.
- (vi) Join G to H. Line GH is the required tangent.




**Exercise 11.2**

1. Draw an arc and divide it into two equal parts.
2. Draw an arc of any length. Draw a tangent without using centre, through point A, when point A is: (i) in the middle of arc. (ii) at the right end of the arc.
3. Draw an arc ABC. Take a point X outside the arc and draw a tangent from X to the arc without using centre of the arc.
4. Construct a circle of radius 4cm. Mark a point P on the circumference and draw a tangent passing through P.
5. Construct a circle of radius 3cm. Draw two tangents from a point A, 5.8cm away from the centre of circle.
6. Construct a circle of radius 2.5cm. Draw two tangents making an angle of:  
(i)  $45^\circ$  (ii)  $90^\circ$  with each other.


7. Construct two equal circles each of radius 3cm. Centres of the circles are 6cm apart. Draw their:
- direct common tangents.
  - transverse common tangents.
8. Construct two circles of radii 2cm and 3.5cm. Draw:
- direct common tangents
  - transverse common tangents to the circles, if the distances between centres of circles is 6.4cm.
9. Construct two touching circles of radii 2.6cm and 3.6 cm. Draw:
- a common tangent that passes through the point of contact of both circles.
  - a common tangent that does not pass through the point of contact of both circles.
  - two common tangents.
10. Construct two intersecting circles of radii 2.5cm and 3cm whose centres are 4.8cm apart. Draw two common tangents to the circles.
11. In the figure, two pulleys of radii 7 inches and 14 inches respectively are shown. Centres of pulleys are 28 inches apart. Find the approximate length of rope moving on pulleys.




12. A satellite A is moving around Earth at an altitude of 7600 km from the surface. The radius of Earth is 6400 km. How far is the satellite from the city P?



13. Two cylindrical rods are bound with a strap. Each rod has diameter 12 cm. How long is the strap?



14. A circular mirror with radius 20 cm hangs by a wire from a hook. The wire is 30 cm long and is a tangent to the mirror in two places. How far above the top of the mirror is the hook?



### I have Learnt

- Locating the centre of a given circle.
- Drawing a circle passing through three given non-collinear points.
- Completing the circle (i) by finding its centre, (ii) without finding its centre, when a part of its circumference is given by
- Drawing a tangent to a given arc, without using the centre, through a given point P when:  
(i) P is middle point of the arc (ii) P is at the end of arc (iii) P is outside the arc.
- Drawing a tangent to a given circle from a point P when:  
(i) P is on the circumference (ii) P is outside the arc.
- Drawing two tangents to a circle meeting each other at a given angle.
- Drawing (i) direct common tangents or external tangents (ii) transverse common tangents or internal tangents, to two equal/unequal circles.
- Drawing a tangent to two unequal (i) touching circles (ii) intersecting circles.
- Applying concepts of practical geometry of circles to real life word problems such as athletic tracks, recreational parks, Ferris wheels and mechanical machines.

## MISCELLANEOUS EXERCISE-11

1. Encircle the correct option in the following.

i. Angle between tangent and radial segment of a circle is:

- (a)  $30^\circ$       (b)  $45^\circ$       (c)  $60^\circ$       (d)  $90^\circ$

ii. Direct common tangents of equal circles are:

- (a) parallel      (b) intersecting      (c) converging      (d) not parallel

iii. How many tangents can be drawn to two intersecting circles?

- (a) 1      (b) 2      (c) 3      (d) infinite

iv. Two circles having radii 3cm and 3.2cm respectively, touch externally. The distance between centres of the circles is:

- (a) 6cm      (b) 0.2cm      (c) 6.2cm      (d) 5.3cm

v. Centres and point of contact of two touching circles are: \_\_\_\_\_.

- (a) collinear      (b) non-collinear      (c) converging      (d) coincident

vi. In which type of triangle, incentre and circumcenter are coincident?

- (a) scalene      (b) isosceles      (c) equilateral      (d) right triangle

vii. Two circles with radii 4cm and 4.8cm respectively, touch internally. The distance between centres of the circles is:

- (a) 4cm      (b) 8 cm      (c) 8.8cm      (d) 0.8cm

viii. Two tangents are drawn at the ends of diameter of a circle of radius 3.5cm.

The distance between tangents is:

- (a) 3.5cm      (b) 7cm      (c) 5cm      (d) 10.5cm

ix. Transverse common tangents intersect each other at:

- (a) 1 point      (b) 2 points      (c) 3 points      (d) 4 points

x. A circle is inscribed in a square having length of side 6cm. What is the radius

- (a) 24      (b) 12      (c) 6      (d) 3

xi. A square is inscribed in a circle of radius 5cm. What is the length of diagonal of the square?

- (a) 2.5cm      (b) 5cm      (c) 10cm      (d) 20cm

xii. Perpendicular bisectors of ..... always pass through centre of circle.

- (a) tangents      (b) secants      (c) radial segments      (d) chords

xiii. How many circles can be drawn through three non-collinear points?

- (a) infinite      (b) 1      (c) 2      (d) 3

xiv. A circle have ..... pair(s) of parallel tangents.


- (a) 1      (b) 2      (c) 3      (d) infinite

## **Basic Statistics**

**After studying this unit students will be able to:**

- Construct cumulative frequency table, cumulative frequency polygon and cumulative frequency curve i.e. ogive
  - Interpret the quartiles, deciles, percentiles and inter quartile range(IQR) from cumulative frequency curve
  - Interpret and analyze box and whisker plot correlation
  - Construct and interpret data from scatter diagram and also draw lines of best fit
  - Measure correlation using scatter diagram
  - Calculate range, standard deviation and variance for grouped data
  - Use the mean and standard deviation to compare 2 sets of data
  - Solve real life problems involving variance and SD of grouped data
  - Apply concepts from measures of dispersion to solve real life problems
  - Calculate the probability of combined events using (where appropriate): Sample space diagram, possibility diagrams, tree diagram & venn diagram
  - Apply addition law of Probability to solve problems involving mutually exclusive events
  - Apply Multiplication law of Probability to solve problems involving independent and dependent events

**Haani and Haider participated in a Qiraat competition, with 8 other equally competent contestants. Judges have to decide about top 3 performers. Can you guess the chance of every contestant to be the top first? After decision of First, can you guess the chance of every remaining contestant to be top second? After decision of Second, can you guess the chance of every remaining contestant to be top third?**



## Frequency Distribution

In statistics it is very important to represent a data in a manageable way so that it becomes easy to understand, analyze and to draw useful results. This useful presentation of data can be made by various methods. Grouped frequency distribution is one of the most important methods to represent the quantitative data. Data can be classified in many ways for efficient usage and productivity. In previous grades we have discussed two types of data

- i. raw data (ungrouped data)
- ii. grouped data (frequency distribution)

Here we will study the construction of cumulative frequency distribution by revising frequency polygon.

### Frequency Polygon

Polygon is a many sided closed figure. To draw a frequency polygon mid points (class marks) are marked on the horizontal axis and frequencies are marked on the vertical axis. Points are plotted with class frequencies and their corresponding class marks. These points are joined by straight line segments. To complete a closed polygon we add extra classes at both ends with zero frequencies. In this way two extra points on x-axis are obtained on both ends. The points are joined by their nearer plotted points. Finally a frequency polygon is obtained.

### Cumulative Frequency Distribution

A table in which cumulative frequencies of classes are written against classes is called a cumulative frequency table “The number of values of a data less than an upper-class boundary is called its cumulative frequency”.

The cumulative frequency table is another way to analyze the frequency distribution. The frequency distribution tells us how many values of the data are present within each class interval, while a cumulative frequency tells us how many values of a data are less than or within each class interval.

In the following table cumulative frequencies are found and interval of 0 frequency is added at the top. Column of upper class boundary is also included which are necessary for construction of cumulative frequency polygon or ogive.

Marks	Marks(upper class boundaries)	No. of students	Cumulative frequency	c.f
0	0.5	0	0	0
1 – 10	10.5	2	2	2
11 – 20	20.5	4	4+2=6	2 + 4 = 6
21 – 30	30.5	9	9+4+2=15	6 + 9 = 15
31 – 40	40.5	6	6+9+4+2=21	15 + 6 = 21
41 – 50	50.5	7	7+6+9+4+2=28	21 + 7 = 28
51 – 60	60.5	3	3+7+6+9+4+2=31	28 + 3 = 31
61 – 70	70.5	4	4+3+7+6+9+4+2=35	31 + 4 = 35

# Cumulative Frequency Polygon and Cumulative Frequency Curve (Ogive)


A frequency distribution can also be represented by a cumulative frequency polygon or Ogive. An ogive and a cumulative frequency polygon are related but distinct concepts in statistics and data visualization.

## Cumulative Frequency Polygon

A frequency polygon made by taking cumulative frequency on y axis is called cumulative frequency polygon. It is made to visualize the cumulative distribution of a variable.

To draw a cumulative frequency polygon, the class boundaries are plotted on x-axis and cumulative frequencies on y-axis. The points are marked on the graph using the upper class boundaries and the respective cumulative frequencies. These points are joined by straight line segments and the polygon is completed by drawing a vertical line segment to join the last marked point with x-axis. Following is a cumulative frequency polygon representing the data given above.

Title: A cumulative frequency polygon showing the marks of 35 students in a test.



**Example:** Late arrival fine imposed to a player is shown below against the terms of 4 weeks:

weeks	1- 4	5-8	9-12	13-16
Amount of fine (Rs)	2000	1000	500	1500

- Construct cumulative frequency table including
  - Column of upper class boundaries
  - Interval for 0 frequency
  - Column of cumulative frequencies
- How much fine is to be paid after 3 terms?
- Construct cumulative frequency polygon of the above data.


**Solution: (a) Cumulative frequency table**

weeks	Upper class boundaries	Amount of fine (Rs)	Cumulative frequencies
0	Less than 0.5	0	0
1-4	Less than 4.5	2000	2000
5-8	Less than 8.5	1000	3000
9-12	Less than 12.5	500	3500
13-16	Less than 16.5	1500	5000

b) After 3 terms, The player has to pay a total amount of Rs. 3500.

c) Construction of cumulative frequency polygon is given below:

**Title: Cumulative freq polygon showing fine imposed on a player against terms of 4 weeks**

**Cumulative Frequency Curve (Ogive)**

A graph showing the cumulative frequency (or cumulative relative frequency) of a variable as a curve, with the x-axis showing the values of the variable and the y-axis showing the respective cumulative frequency is called Cumulative Frequency Curve (Ogive). It is also known as a cumulative distribution function (CDF)


**Example:**

The table below shows the cumulative frequency distribution of masses of cakes in Cocobakes made at a festival:

Mass of cakes (x grams)	$x \leq 400$	$x \leq 500$	$x \leq 600$	$x \leq 700$	$x \leq 800$	$x \leq 900$
No. of cakes	0	20	50	70	80	100

- Draw cumulative frequency curve.
- Estimate from the curve, the number of cakes having masses 750 g or less
- Taking class intervals  $400 < x \leq 500$ ,  $500 < x \leq 600$  ...construct a cumulative frequency distribution

**Solution:** (a) Cumulative frequency curve (Ogive)



An ogive is a smooth curve, while a cumulative frequency polygon is a polygon-shaped graph

- From the curve, we observe that the number of cakes having masses 750 g or less is 75.
- Construction of the frequency cumulative distribution.

Mass of cakes (x g)	Cumulative frequency	Mass of cakes (x g)	frequency
$x \leq 500$	20	$400 \leq x \leq 500$	20
$x \leq 600$	50	$500 \leq x \leq 600$	$50 - 20 = 30$
$x \leq 700$	70	$600 \leq x \leq 700$	$70 - 50 = 20$
$x \leq 800$	80	$700 \leq x \leq 800$	$80 - 70 = 10$
$x \leq 900$	100	$800 \leq x \leq 900$	$100 - 80 = 20$

## Measures of Distribution

These measures are used to understand the spread and distribution of the data, and are often used with measures of central tendency to get a better understanding of the dataset. They include:

## i. Quartiles

The three values which divide the whole data into four equal parts are called quartiles. Quartiles are usually denoted by  $Q_1$ ,  $Q_2$  and  $Q_3$ . The lower quartile  $Q_1$  divides the data in 1 : 3. The middle quartile  $Q_2$  divides the data in 1 : 1. It is in fact nothing different from median of data. The upper quartile  $Q_3$  divides the data in 3 : 1.

$$Q_1 = \frac{n}{4}^{\text{th}} \text{ value of data} \quad \& \quad Q_2 = \left( \frac{2n}{4} = \frac{n}{2} \right)^{\text{th}} \text{ value of data} = \text{median}$$

When data is arranged in some order, it can be represented by a single value that lies exactly at the centre of data. "The central or middle most value of an arranged data is called its median".

$$Q_3 = \frac{3n}{4}^{\text{th}} \text{ value of data}$$

The quartiles can also be estimated graphically from a cumulative frequency polygon.

- To estimate  $Q_1$  (lower quartile), mark  $\frac{n}{4}$  on y-axis.
- Draw a line parallel to x-axis from that point to intersect cumulative frequency polygon.
- Drop a perpendicular from this point of intersection to x-axis. The value of data where the perpendicular meets x-axis is lower quartile  $Q_1$ .

Similarly,  $Q_2$  and  $Q_3$  can be estimated from cumulative frequency curve, by marking  $\frac{n}{2}$  and  $\frac{3n}{4}$

on y-axis and following the same procedure used for  $Q_1$ .

### Example:

The scores on a test were compiled into a dataset with the following quartiles:

$$Q_1 = 40, \quad Q_2 \text{ or median} = 70, \quad Q_3 = 90$$

What percentage of students scored above 90 on the test?

**Solution:** From the quartiles it is obvious that one fourth (25 percent) of the students scored above 90.

## ii. Deciles

Deciles divide a dataset or a distribution into 10 equal parts, each representing 10% of the data. By looking at the deciles, we can get a sense of the shape of the distribution, including where the majority of the data points are concentrated and where the outliers are lying.

Ten Deciles of a dataset are described as follows:

D1: lowest 10% of the data

D2: lowest 20% of the data

D3: lowest 30% of the data

.....

D10: highest decile under which 100% of the data lies.

### Key Fact:

Deciles are often used in statistics, economics, and social sciences to understand the distribution of a variable, such as income, wealth or exam scores.

**Example:** A company wants to analyze the salaries of its employees. The related department compiles a dataset and calculates the deciles. The deciles are as follows:

D1 = Rs.40,000 , D5 or median = Rs.60,000 , D10 = Rs.80,000

What percentage of employees earn a salary above Rs.60,000 but below Rs.80,000?

**Solution:** From the above data, percentage of employees earning salaries above 60,000 and below Rs. 80,000 are between D5(median) and D10 i.e  $90\% - 50\% = 40\%$

(Remember: D10 represents highest 10 % i.e Rs. 80,000 and D9 represents below Rs. 80,000)

### III. Percentiles

Percentiles divide a dataset or a distribution into 100 equal parts, each representing 1% of the data. It's a way to understand the distribution of a dataset or a variable by looking at the percentage of data points that are below a specific value.

Some common percentiles include:

- 25th percentile (Q<sub>1</sub>): 25% of data points are below this value.
- 50th percentile (Q<sub>2</sub> or median): 50% of data is below Q<sub>2</sub>.
- 75th percentile (Q<sub>3</sub>): 75% of data points are below this value.
- 90th percentile: 90% of data points are below this value.
- 99th percentile: 99% of data points are below this value

#### Maths Play Ground:

Help students to design quiz wheel or quiz machine for the reinforcement of measures of distribution.

#### Key Fact:

Percentiles are often used to:

understand the distribution of a variable (e.g., income, scores, masses).

identify outliers.

compare individual values to the overall distribution.

set benchmarks (e.g., top 10 %).

evaluate performance (e.g., moving up or down in percentiles).

**Example:** Uzair scored in the 75th percentile on a test. Find how much percentage of the students scored below him and how much percentage of students scored above him.

**Solution:** From the above information it is clear that the student's score is higher than 75% of the total students. It means that

- 75% of the students scored below him
- 25% scored above him

### IV. Inter Quartile Range

$$IQR = Q_3 - Q_1$$

The IQR represents the range of values within which the middle 50% of the data points fall. In other words, it's the spread of the data within the middle half of the distribution.

IQR can describe:

Spread: A small IQR indicates less scattered, while a large IQR indicates more scattered data.

Variability: The IQR is a more efficient measure of variability than the Range.

Outliers: The IQR can help identify outliers. (data points beyond 1.5 times the IQR from the quartiles are considered outliers).

Comparison: IQR is useful for comparing the spread of different datasets.

**Example:**

A teacher wants to analyze the test scores of her students. She compiles a dataset as:

first quartile  $Q_1 = 60$  and third quartile  $Q_3 = 95$

- What is the Interquartile Range (IQR) of the test scores?
- What percentage of students scored between 60 and 95 (inclusive)?

**Solution:**


- The Interquartile Range (IQR) of the test scores is given by the relation:

$$IQR = Q_3 - Q_1 = 95 - 60 = 35$$

- As  $Q_3 - Q_1$  represents spread of the middle half of a data set so 50% of students scored between 60 and 95 (inclusive)

**Example:**

The adjoining figure shows an ogive representing the distribution of time in minutes spent by 60 preachers of SEED to deliver single Seerah lecture. Find  $Q_1$ ,  $Q_2$ ,  $Q_3$  and inter quartile range.

**Solution:**

To estimate the median time spent from ogive, we observe that 50 % (half of) preachers spent less than or equal to the median time to deliver the lecture.

Median corresponds to the 5<sup>th</sup> decile and 50<sup>th</sup> percentile, i.e.

$$\text{Median} = Q_2 = D_5 = P_{50} = \frac{1}{2} \times 60 = 30$$

From 30 on y-axis, draw a horizontal line to meet the curve and then draw a vertical line to meet x-axis as in above figure.

From the figure, we observe that median time to deliver a lecture is 43.5 minutes.

Lower quartile corresponds to 25th percentile i.e.

$$Q_1 \text{ (the lower quartile)} = P_{25} = \text{one quarter (25\% of) total frequency} = \frac{1}{4} \times 60 = 15$$

From the curve, the lower quartile = 37 minutes.

Upper quartile corresponds to 75th percentile i.e.

$$Q_3 \text{ (the upper quartile)} = P_{75} = \text{three quarters (75\% of) total frequency} = \frac{3}{4} \times 60 = 45$$

From the curve, the upper quartile = 50.5 minutes.

$$\text{The inter Quartile Range} = Q_3 - Q_1 = 50.5 - 37 = 13.5$$

## Exercise 12.1

- 1.** Construct cumulative frequency column for following frequency table and give answers.

Class interval	1 – 10	11 – 20	21 – 30	31 – 40	41 – 50	51 – 60
Frequency	3	4	7	9	5	2

- (i) What is the total number of items in the data?
- (ii) In which class interval 8<sup>th</sup> item of the data lies?
- (iii) What is number of items having worth less than 21?
- (iv) Which group contains highest number of items?
- (v) What is the lower boundary of the last class?

- 2.** The mass gained by a newly born baby is given below:

Age in months	1	2	3	4	5	6
No. of kg gained	4	3	2	3	2	1

Draw a cumulative frequency polygon (not ogive) to represent the data. Also find from the graph, how much he/she weighed at the age of 4 and half months?

- 3.** The ages of refugees in Ghaza camp were recorded as below:

Ages in years	20 – 24	25 – 29	30 – 34	35 – 39	40 – 44	45 – 49
No. of refugees	5	16	12	10	8	4

Draw an ogive (not polygon) to represent the data and Find

- a) First quartile.
- b) 7<sup>th</sup> decile.
- c) 60<sup>th</sup> percentile.

- 4.** Scores of a Hifz e Quran test were grouped into the following frequency distribution.

scores	41 – 50	51 – 60	61 – 70	71 – 80	81 – 90	91 – 100
f	3	17	20	30	18	2

Draw a cumulative frequency table and cumulative frequency curve (ogive) and tell:


- a) How many boys appeared in the test?
- b) What is median score?
- c) How many boys scored under median?
- d) How many boys fell in first decile?
- e) How many boys fell in 30<sup>th</sup> percentile?
- f) If excellence award is to be given to boys falling in 100<sup>th</sup> percentile, how many boys got excellence award?

- 5.** The masses measured to the nearest kg of 50 cubs are shown below:


Mass (kg)	60 – 64	65 – 69	70 – 74	75 – 79	80 – 84	85 – 89
f	2	6	12	14	10	6

- (i) Construct a cumulative frequency table
- (ii) Construct a cumulative frequency polygon and estimate number of cubs above 77 kg.
- (iii) Construct a cumulative frequency curve and estimate number of cubs above 77 kg.
- (iv) Compare the results of (ii) & (iii)

6. Study the adjoining curve showing heights (m) of 100 buildings in a megacity carefully and find:



- a) Number of buildings with heights less than 50 m.
  - b) % age of buildings higher than 40 m.
  - c) Ratio of the number of buildings less than 50m and more than 50m
  - d) No. of Buildings falling in first quartile
  - e) Maximum height of buildings falling under 8<sup>th</sup> decile.
  - f) Maximum height of buildings not falling under 30<sup>th</sup> percentile.
7. Mr. Rashid wants to analyze the salaries of his employees. The first quartile (Q1) is of Rs.50,000 a second quartile (Q2) of Rs.70,000, a third quartile (Q3) of Rs.90,000 and maximum and minimum salaries are of Rs.25,000 & Rs.150,000 respectively. Find range and inter quartile range of the data.
8. Khadeeja scored 85 on a test, and the teacher reported that her score was in the 75th percentile. What does this information indicate about status of Khadeeja in her class?
9. Uzair wants to price his house based on the prices of similar houses in the area. The prices have a Q1 of Rs. 20,000,000, Q2 of Rs. 25,000,000, Q3 of Rs. 30,000,000, and a maximum price of Rs. 35,000,000. What is the possible price of the house if it is in the third quartile? Also find IQR of the data.
10. The adjoining cumulative frequency curve shows the number of faulty items in a shipment of 100 items produced by Khani's Collection. Estimate
- i. The median                      ii. Lower quartile
  - iii. Upper quartile                iv. IQR
  - v. 7th decile                       vi. 30th percentile



# The Box and Whisker Plot Correlation

## Box-and-Whisker Plot (or simply box plot)

A box plot typically consists of the following components:

**Box:** A rectangular box that represents the interquartile range (IQR). The box is divided into two parts by a vertical line, which represents the median ( $Q_2$ ) of the data.

**Whiskers:** Two horizontal lines that extend from the box to the minimum and maximum values in the dataset, excluding outliers. The whiskers represent the range of the data.


**Outliers:** Data points that fall beyond 1.5 times the IQR from the quartiles are considered outliers and are plotted individually.

**Median:** The vertical line inside the box represents the median ( $Q_2$ ) of the data.

**Quartiles:** The edges of box represent 25th percentile ( $Q_1$ ) and the 75th percentile ( $Q_3$ ).

Box plots are useful for comparing datasets, identifying outliers, visualizing distribution and summarizing data

**Example:** The given box and whisker plot shows the prices of stationery items, Hashir purchased yesterday.



- a) What is the median price of an item?
- b) what are the min & max prices of the items purchased by Hashir?
- c) What is IQR of the data?      d) what %age of items is above Rs. 800?

## Solution:

- a) From the box plot, it is clearly seen that median price of an item is Rs. 300
- b) From the box plot, it is observed that min price of any item is Rs. 50 & max price of any item is Rs. 1000
- c)  $IQR = Q_3 - Q_1 = 800 - 160 = 640$
- d) As items between Q<sub>3</sub> and Q<sub>4</sub> are 25 % so %age of items above Rs. 800(Q<sub>3</sub>) is 25%.

## Box and Whisker Plot Correlation

The box and whisker plot correlation means the patterns or relationships between the distributions of two or more variables. Box and whisker plot correlations include:

### 1. Positive correlation:

If the boxes and whiskers of both variables tend to move in the same direction (e.g., both increase or decrease together), it may indicate a positive correlation.

### 2. Negative correlation:

If the boxes and whiskers of both variables tend to move in opposite directions (e.g., one increases while the other decreases), it may indicate a negative correlation.


### 3. No correlation:

If the boxes and whiskers of both variables appear unrelated or don't show a clear pattern, it may indicate no correlation.

**Key Fact:**

Box and whisker plots only provide a visual representation and don't quantify the correlation.

Example shows volume of a gas is shown in first box-and-whisker plot and increase in temperature is shown in second box-and-whisker plot in the adjoining figures.



In this example, we observe that the boxes and whiskers of both variables tend to move in the same direction (both increasing together), suggesting a positive correlation.

**Example:**

The following datasets show 2 Variables. Hours of Exercise and Body Mass Index, (BMI) of a gym members.

variables	Min/max	Q1/Q3	median	Q3/Q1	Max/min
Hours	0(min)	2(Q1)	4	6(Q3)	8(max)
BMI	30(max)	28(Q3)	25	22(Q1)	20(min)



Draw box and whisker plots.

- a) Find type of correlation    b) Comment on type of correlation

**Solution:**

- a) As the hours of exercise (Variable A) increase, the BMI (Variable B) tends to decrease. The boxes and whiskers of both variables move in opposite directions, indicating a negative correlation.
- b) The negative correlation suggests that exercise may lead to weight loss, as higher exercise hours are associated with lower BMI values.

**Box-and-whisker plots  
Time (hrs) of exercise & BMI (kg/m<sup>2</sup>)**



# **Construction of Scatter Diagram, Lines of Best Fit and Interpretation of Data**

Construction of a scatter diagram consists of:

- i. Plotting the data points on a coordinate plane.
- ii. Representing each data point by a mark (e.g., dot, circle, or square).


We solve an example to understand the concept.

**Example:** Table below shows the masses and heights of 11 boys.

Heights(cm)	167	165	184	153	180	170	179	174	160	176	157	177
Mass(kg)	60	54	69	48	64	58	68	64	51	67	46	63

- a) Draw a scatter diagram for the above data.
- b) What type of correlation is observed in the scatter diagram?
- c) Draw a line of best fit on the scatter diagram.
- d) Use this line of best fit to estimate mass of a boy whose height is 150cm.

**Solution (a):**



- (b) From the graph, it is observed that there is a strong positive correlation between masses and heights of boys. More height, more the mass of boys.
- (c) Line of best fit is drawn having almost half the data points on either sides of it.
- (d) From scatter diagram, the boy with height 150cm should have a mass of 39 kg app.

## **Interpretation of a Scatter Diagram**

**Pattern:** There are several types of patterns:

- Positive correlation: As x increases, y also increases.
- Negative correlation: As x increases, y decreases.
- No correlation: No apparent relationship between x and y.

**Strength:** Observe the strength of the relationship. A stronger relationship means the points are closer to a straight line.

**Outliers:** Identify data points that don't fit the pattern.

### Drawing Lines of Best Fit

A line of best fit helps to predict the value of a variable  $y$  for a given  $x$ . It is a line that appears to fit the pattern. It may or may not pass through any data points however almost equal number of points should lie on either side of it. The line of best fit can be represented by

$$y = mx + c.$$

Interpretation of a Line of Best Fit includes following attributes:

**Slope ( $m$ ):** The rate of change of  $y$  with respect to  $x$ .

**Intercept ( $c$ ):** The value of  $y$  when  $x$  is 0.

**Prediction:** Use the equation to predict the value of  $y$  for a given  $x$ .

#### Key Fact:

A line of best fit is not always a perfect fit and it's essential to consider the strength of the relation and the presence of outliers when interpreting the results.

### Measuring Correlation using Scatter Diagram

Measuring correlation using a scatter diagram involves analyzing the pattern and strength of the relationship between two variables.

It includes following steps:

- 1 Plot the data
- 2 Identify the pattern (Positive correlation, Negative correlation and No correlation)
- 3 Assess the strength (Strong correlation, Moderate correlation and Weak correlation)
- 4 Calculate the correlation coefficient  $r$  by:
  - Identifying the data points  $(x, y)$  by coordinates of each point on the scatter plot.
  - Finding the mean of the  $x$ -values ( $\bar{x}$ ) and the mean of the  $y$ -values ( $\bar{y}$ ).
  - Calculating the deviation from the mean for  $x$  ( $x_i - \bar{x}$ ) and  $y$  ( $y_i - \bar{y}$ ).
  - Multiplying the  $x$ -deviation by the  $y$ -deviation  $(x_i - \bar{x})(y_i - \bar{y})$ .
  - Adding up the multiplied deviations for all points.
  - Dividing the sum of the products by the square root of the product of the sum of the squared  $x$ -deviations and the sum of the squared  $y$ -deviations. i.e.

$$r = \frac{\sum \{(x_i - \bar{x})(y_i - \bar{y})\}}{\sqrt{(\sum (x_i - \bar{x})^2) \times (\sum (y_i - \bar{y})^2)}}$$

Where  $r$  is the correlation coefficient, ranging from  $-1$  (perfect negative correlation) to  $1$  (perfect positive correlation).

- 5 Interpret the results high  $r$  value i.e. close to  $1$  or  $-1$  means Strong correlation, Low  $r$  value i.e. close to  $0$  means weak correlation & sign of  $r$  indicates the direction of the correlation i.e. positive or negative.

**Example:**


The table below shows the reading time of 10 children aged between 11 and 16 years to win the 50 pages reading activity in 10 minutes.

Age (yrs)	13.2	14.9	13.3	15.3	13.6	15.5	14.0	14.3	12.8	14.8
Time (min)	13.3	12.1	13.9	12.3	13.2	11.6	13.1	12.3	13.9	13.1

- Draw a scatter diagram for the above data.
- What type of correlation is observed in the scatter diagram?
- Draw a line of best fit on the scatter diagram.
- Use your line to estimate the time that a participant aged 14.5 years is expected to complete 50 pages reading activity.
- Measure the correlation coefficient.
- Is your line of best fit reliable to estimate the time taken by a child of age 1 year?

**Solution:**

a)



- From the data it is observed that older the age of child corresponds to greater speed in reading activity. The line is sloping down from left to right showing negative correlation (and neg. slope as well)
- .... drawn on the graph
- From the graph reading for 14.5 years age, expected time to complete 50 pages reading activity is 12.7min.
- For measuring correlation coefficient, let's calculate the means:

$$\text{Age } (\bar{x}): 14.2 \quad \text{Time } (\bar{y}): 12.9$$

Next, we calculate the deviations, their products and squared deviations.

age	time	$(x_i - \bar{x})$	$(y_i - \bar{y})$	$(x_i - \bar{x})(y_i - \bar{y})$	$(x_i - \bar{x})^2$	$(y_i - \bar{y})^2$
13.2	13.3	-1.0	0.4	-0.4	1	0.16
14.9	12.1	0.7	-0.8	-0.56	0.49	0.64
13.3	13.9	-0.9	1.0	-0.9	0.81	1.0
15.3	12.3	1.1	-0.6	-0.66	1.21	0.36
13.6	13.2	-0.6	0.3	-0.18	0.36	0.09
15.5	11.6	1.3	-1.3	-1.69	1.69	1.69
14.0	13.1	-0.2	0.2	-0.04	0.04	0.04
14.3	12.3	0.1	-0.6	-0.06	0.01	0.36
12.8	13.9	-1.4	1.0	-1.4	1.96	1.0
14.8	13.1	0.6	0.2	0.12	0.36	0.04
		$\sum(x_i - \bar{x})$	$\sum(y_i - \bar{y})$	$\sum\{(x_i - \bar{x}) \times (y_i - \bar{y})\} = -5.77$	$\sum(x_i - \bar{x})^2 = 7.93$	$\sum(y_i - \bar{y})^2 = 5.38$

Finally, we calculate the correlation coefficient ( $r$ ):

$$r = \frac{\sum\{(x_i - \bar{x}) \times (y_i - \bar{y})\}}{\sqrt{(\sum(x_i - \bar{x})^2) \times (\sum(y_i - \bar{y})^2)}}$$

$$r = \frac{-5.77}{\sqrt{7.93 \times 5.38}} \approx -0.88$$

The correlation coefficient ( $r$ ) is approximately  $-0.88$ , indicating a strong negative linear relationship between Age and Time. As Age increases, Time tends to decrease, and vice versa.

- f) No, it is highly unreliable as it lies outside the range as there are no children with age less than 10 years.

### Exercise 12.2


1. A mechanical engineer wants to analyze the relationship between the mass of a car and its fuel efficiency. Show the data by box and whisker plots.

variables	Min/Max	Q1	median	Q3	Min/Max
Mass (kg)	1100	1200	1500	1700	2000
Fuel efficiency (Km/l)	24	20	14	12	10

- a. Find IQR      b. What is type of correlation?  
c. What conclusion can be drawn from this?

2. Adjoining box & whisker plot shows wages for part time and full-time employees:


**Hourly rate for Full and Part-time employees**



- a) What is the amount of median wages paid to full time employees?
  - b) What are minimum and maximum wages paid to part time employees?
  - c) What is IQR for the dataset of fulltime employees?
  - d) What is IQR for the dataset of part time employees?
  - e) What type of correlation is observed in both datasets?
3. Maryam wants to investigate the relationship between the amount of rainfall and the number of flowers blooming in her garden. Draw the scatter diagram.

Rain (inch)	2	4	6	8	10
No. of flowers	10	22	55	70	80

- a. Draw a line of best fit.      b. Comment on type of correlation
4. Given scatter diagram is showing relation between practice time and marks obtained in math's test.
- a. Find type of correlation.
  - b. Draw line of best fit.



5. Draw the scatter diagram for the data showing the relationship between amount of pollution (measured in Air Quality Index (AQI)) and number of sick days reported by a locality:

AQI (Pollution Level)	20	40	60	80	100	120	10	30	50	70
No. of Sick Days	5	18	25	40	60	80	3	15	20	27

Find correlation and what does this suggest?

6. A teacher wants to analyze the test scores of her students. Draw box and whisker plot to show a median score of 75, a first quartile of 60, and a third quartile of 85. Find:
- What percentage of students scored above 85?
  - IQR

## Measures of Dispersion

The single value which can represent a data is known as average or measure of central tendency. In grade 9, different types of averages e.g. mean, median and mode are studied. Let us find the arithmetic mean of scores of five participants of a math quiz:

$$V \text{ (Bilal's score)} = 5, 5, 5, 5, 5 \Rightarrow \bar{V} = \frac{\Sigma V}{n} = \frac{25}{5} = 5$$

$$W \text{ (Umer's score)} = 1, 3, 5, 8, 8 \Rightarrow \bar{W} = \frac{\Sigma W}{n} = \frac{25}{5} = 5$$

$$X \text{ (Usman's score)} = 2, 4, 5, 6, 8 \Rightarrow \bar{X} = \frac{\Sigma X}{n} = \frac{25}{5} = 5$$

$$Y \text{ (Hassan's score)} = -10, 0, 10, 12, 13 \Rightarrow \bar{Y} = \frac{\Sigma Y}{n} = \frac{25}{5} = 5$$

$$Z \text{ (Abdullah's score)} = -15, -5, 10, 15, 20 \Rightarrow \bar{Z} = \frac{\Sigma Z}{n} = \frac{25}{5} = 5$$

We have seen that all of the above five data have same measure of central tendency (arithmetic mean) but they vary in different ways. The values of above data are scattered away from their average in different ways. It is a confusing situation because we are unable to compare the data on the basis of their average. We can say that the average of a data is not more significant unless the scatterings of values around the average are measured.

The techniques or formulas which are used to measure, how the values of a data are scattered around its central value, are called measures of dispersion.

### Types of Measures of Dispersion

#### (a) Range

Range is the difference between the maximum and the minimum values of the data. i.e.

$R = x_m - x_o$ , where  $x_m$  = maximum value and  $x_o$  = Minimum value

#### (b) Variance

Variance is the mean of squared deviation taken from mean.

Usually it is denoted by  $S^2$ . The formula for variance of grouped data:

$$S^2 = \frac{\sum f(x_i - \bar{x})^2}{\sum f} \quad (\text{Proper mean formula})$$

$$\text{or } S^2 = \frac{\sum fx^2}{\sum f} - \left(\frac{\sum fx}{\sum f}\right)^2 \quad (\text{Direct formula})$$

#### (c) Standard Deviation

Standard deviation is a positive square root of variance. It is denoted by  $S$  and given by following formulae:

$$S = \sqrt{\frac{\sum f(x_i - \bar{x})^2}{\sum f}} \quad (\text{Proper mean formula})$$

$$\text{or } S = \sqrt{\frac{\sum fx^2}{\sum f} - \left(\frac{\sum fx}{\sum f}\right)^2} \quad (\text{Direct formula})$$

**Example:** Data shows scores of Huffaz in a Hifz Quran test:

scores	41 - 50	51 - 60	61 - 70	71 - 80	81 - 90	91 - 100
f	3	17	20	30	18	2

Find Range, variance and standard deviation of the data.

**Solution:** Range =  $x_m - x_o = 100 - 41 = 59$

For the variance and SD, we need to construct the following table:

scores	x	$x^2$	f	$fx$	$fx^2$
41-50	45.5	2070.25	3	136.5	6210.75
51-60	55.5	3080.25	17	943.5	52364.25
61-70	65.5	4290.25	20	1310	85805
71-80	75.5	5700.25	30	2265	171007.5
81-90	85.5	7310.25	18	1539	131584.5
91-100	95.5	9120.25	2	191	18240.5
		$\sum f = 90$		$\sum fx = 6385$	$\sum fx^2 = 465212.5$

$$\text{Variance: } S^2 = \frac{\sum fx^2}{\sum f} - \left(\frac{\sum fx}{\sum f}\right)^2 = \frac{465212.5}{90} - \left(\frac{6385}{90}\right)^2 = 135.9 \text{ app.}$$

$$\text{Standard deviation: } S = \sqrt{\frac{\sum fx^2}{\sum f} - \left(\frac{\sum fx}{\sum f}\right)^2} = \sqrt{\frac{465212.5}{90} - \left(\frac{6385}{90}\right)^2} = \sqrt{135.9} = 11.6$$

### Coefficient of Variation (C.V.)

The percentage ratio between the standard deviation and mean of a data is called coefficient of variation i.e.

$$C.V. = \frac{SD}{\text{Mean}} \times 100$$

$$\text{In above example, } CV = \frac{11.6}{70.5} \times 100 = 16.4 \text{ app.}$$

#### Key Fact:

C.V. is used to compare the dispersion of different data which differ in their means and units. It is also used as a criterion for consistent performance. The data having smaller C.V is more consistent than the data having larger C.V.

## Comparison of 2 Data Sets using Mean and Standard Deviation

To compare two data sets using mean and standard deviation, follow these steps:

1. Calculate the arithmetic mean of each data set.
2. Calculate the standard deviation of each data set.
3. Compare the means which indicates a difference in the central tendency of the two data sets.
4. Compare the standard deviations which indicate a difference in the variability or spread of the two data sets.

**Example:** Following are scores of 2 girl sections orchid and teal in Math quiz.

Score chart for Teal:

scores	41-50	51-60	61-70	71-80	81-90	91-100
No. of students	3	8	6	7	4	2

Score chart for orchid:

scores	41-50	51-60	61-70	71-80	81-90	91-100
No. of students	3	6	4	9	1	7

- i. Calculate the arithmetic mean of each data set.
- ii. Calculate the standard deviation of each data set.
- iii. Compare the means. What does this indicate?
- iv. Compare the standard deviations. What does this indicate?
- v. Calculate CV for both data sets.

**Solution:**

Teal section				
scores	x	f	fx	$f(x - \bar{x})^2$
41-49	45.0	4	180	2116
50-60	55.0	6	330	1014
61-70	65.5	6	393	37.5
71-80	75.5	8	604	450
81-90	85.5	4	342	1225
91-100	95.5	2	191	1512.5
		$\sum f = 30$	$\sum fx = 2040$	$\sum f(x - \bar{x})^2 = 6355$

Orchid section				
scores	y	f	fy	$f(y - \bar{y})^2$
41-49	45.0	3	135	2187
50-60	55.0	6	330	1734
61-70	65.5	4	262	169
71-80	75.5	9	679.5	110.25
81-90	85.5	1	85.5	182.25
91-100	95.5	7	668.5	3865.75
		$\sum f = 30$	$\sum fy = 2160.5$	$\sum f(y - \bar{y})^2 = 8248.25$

$$\text{i) } \bar{x} = \frac{\sum fx}{\sum f} = \frac{2040}{30} = 68$$

ii) SD teal :

$$S_{\text{teal}} = \sqrt{\frac{\sum f(x - \bar{x})^2}{\sum f}} = \sqrt{\frac{6355}{30}} = 14.6$$

$$\bar{y} = \frac{\sum fy}{\sum f} = \frac{2160.5}{30} = 72.0$$

SD orchid:

$$S_{\text{orchid}} = \sqrt{\frac{\sum f(y - \bar{y})^2}{\sum f}} = \sqrt{\frac{8248.25}{30}} = 16.6$$

**Comparison:**

- iii. The mean score of Orchid section (72.0) is higher than the mean score of Teal section (68.0), indicating that the average performance of Orchid is better than Teal.
- iv. The standard deviation of Orchid (16.6) is greater than the standard deviation of Teal (14.6), indicating that the performance of students in teal is more consistent and less spread out than in Orchid. Such performance is more reliable.

v.  $CV_{teal} = \frac{14.6}{68} \times 100 = 21.47$

$$CV_{orchid} = \frac{16.6}{72} \times 100 = 24.41$$

**Real Life Application of Variance and Standard Deviation**

Variance and standard deviation have numerous real-life uses in various fields, e.g. Finance, Quality Control, Healthcare, Sports, Education, Environmental Science & Engineering.

**Exercise 12.3**

- What is dispersion? Write names and formulae of 3 measures of dispersion.
- The energy consumption for a month is given below for Rabee's home.

Units consumed	101-200	201-300	301-400	401-500	501-600	601-700
No. of months	2	1	3	1	3	2

Find Range and variance of the energy consumption.

- Maria wants to analyze the water supply in her area, given in the data as shown:

Water supply				
Water supply (gallons)	501-700	701-900	901-1100	1101-1300
No. of hours	4	5	2	1

Find i. mean supply of water in the area. ii. Variance of the supply of water.  
iii. SD of the supply of water. iv. CV of the supply of water.

- An organization wants to analyze salaries, grouped by department as shown below:

Department	Salary Range	Number of Employees
Elementary	40,000 - 60,000	10
Secondary	60,000 - 80,000	12
Higher secondary	80,000 - 100,000	8

Find the range, variance and standard deviation of the salaries across all departments. Also calculate CV of the data.

- A teacher wants to analyze the scores of students in different sections of class 10 as shown:

Class	Scores Range	Number of Students
10 Mauve	70 - 80	15
10 teal	60 - 70	12
10 hazel	50 - 60	10

Find the variance and standard deviation of the grades of all sections .

6. Mutahhir wants to analyze the purchases made by customers, grouped by age as shown:

Age Group	Purchase worth (in Rs.)	Number of Customers
kids	0 - 500	20
juniors	500 - 1000	30
seniors	1000 - 2000	25

Find the range, variance and standard deviation of the purchases across all age groups.

7. Namra wants to analyze blood pressure of patients in OT & OPD grouped by age as shown:

Patients in OT						
BP	100-110	110-120	120-130	130-140	140-150	150-160
No. of patients	3	8	6	7	4	2

Patients in OPD						
BP	100-110	110-120	120-130	130-140	140-150	150-160
No. of patients	9	1	5	7	6	2

Find the Range, Mean, variance and standard deviation of the blood pressure of both datasets. Also find CV and comment on results.

8. Khola wants to forecast the temperature for the next week for Jhelum and Indiana. She has the history of a week for both of the cities as given:

Jhelum				Indiana					
Temperature(F°)	96 - 100	101 - 105	106 - 110	111 - 115	Temperature(F°)	51 - 60	61 - 70	71 - 80	81 - 90
No. of days	2	3	1	1	No. of days	3	1	2	1

Help her to find:

- i. Mean of both data sets.                    ii. Variance of both datasets

Also comment on the possible predictions for next week temperature for both cities.

(Hint: In weather forecasting, variance is used to measure the spread or uncertainty in temperature predictions. A higher variance indicates a wider range of possible temperatures, making it more challenging to predict the exact temperature.)

# Probability

Probability is a measure of the likelihood of an event occurring. It is a number between 0 and 1 both inclusive that represents the chance of an event happening. Here are some key concepts in probability:

Experiment:	An action or process that can produce a set of outcomes.
Outcome:	A specific result of an experiment.
Sample Space:	The set of all possible outcomes of an experiment.
Event:	A set of one or more outcomes of an experiment.
Probability of an Event:	A number between 0 and 1 both inclusive that represents the likelihood of an event.

## Types of Probability

There are several types of probability such as:

- Theoretical Probability  
The number of favorable outcomes divided by the total number of possible outcomes.
- Experimental Probability  
Based on the results of repeated trials of an experiment.
- Conditional Probability  
The probability of an event occurring given that another event has occurred.
- Dependent Probability  
The probability of two or more events occurring dependently on each other.
- Mutually Exclusive Probability  
The probability of two or more events that cannot occur simultaneously.
- Joint Probability  
The probability of two or more events occurring together.

## Probability of Combined Events

The probability of combined events depends on whether the events are independent, dependent, or mutually exclusive.

### Probability of Independent Events

The probability of two or more independent events occurring is the product of their individual probabilities.

$$\text{i.e. } P(A \text{ and } B) = P(A) \times P(B)$$

### Probability of Dependent Events

The probability of two or more dependent events occurring is the product of their individual probabilities, considering the conditional probability of each event.

$$\text{i.e. } P(A \text{ and } B) = P(A) \times P(B|A)$$

### Probability of Mutually Exclusive Events

The probability of two or more mutually exclusive events is the sum of their individual probabilities.

$$\text{i.e. } P(A \text{ or } B) = P(A) + P(B)$$

### Conditional Probability

Probability of an event occurring given that another event has occurred.

$$\text{i.e. } P(A|B) = P(A \text{ and } B) / P(B)$$

For example:

1. Probability of drawing two Kings (with replacement):

$$P(\text{King}) \times P(\text{King}) = 1/13 \times 1/13 = 1/169 \approx 0.006 \text{ (or } 0.6\%)$$

Probability of drawing two Kings (without replacement):

$$P(\text{King}) \times P(\text{King}) = 4/52 \times 3/51 = 3/663 \approx 0.0045 \text{ (or } 0.45\%)$$

2. Probability of rolling two 6s:  $P(6) \times P(6) = 1/6 \times 1/6 = 1/36$

3. Probability of drawing a King and then a Queen:

$$P(\text{King}) \times P(\text{Queen} | \text{King}) = (4/52) \times (4/51) = 16/2652$$

Remember to consider the type of events and their relationships when calculating the probability of combined events.

### Probability of Combined Events using Sample Space Diagrams

A sample space diagram is a visual representation of all possible outcomes of an experiment. It can be used to calculate the probability of combined events by identifying the favorable outcomes and dividing them by the total number of possible outcomes.

To calculate the probability of combined events using a sample space diagram:

- Draw a sample space diagram showing all possible outcomes of the experiment.
- Identify the favorable outcomes that meet the conditions of the combined event.
- Count the number of favorable outcomes.
- Divide the number of favorable outcomes by the total number of possible outcomes in the sample space.

#### Example:

What is the probability that both coin A and coin B land heads up if both coins are flipped simultaneously?

#### Solution:

Favorable Outcomes: hH (both coins land heads up)

Number of Favorable Outcomes: 1

Total Number of Possible Outcomes: 4

Probability:  $1/4$  or  $0.25$

This means that the probability of both coins landing heads up is  $1/4$  or  $25\%$ .

Sample space diagram		Coin 1	
		H	T
Coin 2	h	hH	hT
	t	tH	tT

### Probability of Combined Events using Diagram

There are several types of sample space diagram, including:

Possibility Diagrams

Tree Diagrams

Venn Diagrams

By using an appropriate sample space diagram, we can easily visualize and calculate the probability of combined events. This method is especially helpful while dealing with complicated experiments and multiple events.

**(a) Probability of Combined Events using Possibility Diagram**

Possibility diagram is graphical representation of all possible outcomes of an experiment.

### **Example:**

Raabi and Jerry are hoping on their play circles of four sectors each painted in red, green, yellow and blue colours.

**Find the probability that they will jump on:**



**Solution:** We can associate the ordered pair  $(x, y)$ , where  $x$  represents hops of Raabi and  $y$  represents hops of Jerry to show an outcome in the sample space. Here (blue, green) means Raabi lands on blue while Jerry lands on green.

So the sample space is:

$$S = \{(red, red), (red, green), (red, yellow), (red, blue), (green, red), (green, green), (green, yellow), (green, blue), (yellow, red), (yellow, green), (yellow, yellow), (yellow, blue), (blue, red), (blue, green), (blue, yellow), (blue, blue)\}$$

Which seems complicated and time taking method of writing sample space for bigger problems.

We can express the same in more systematic and efficient way through possibility diagram..

There are 16 dots representing individual outcome,  
i.e.  $n(S) = 16$

Let  $E_1$  be the event that both girls jump on same colour.

And  $E_2$  be the event that they jump on different colours

The loop on the diagonal enclosing four dots is associated

With possible outcomes of  $E_1$

$$\therefore n(E_1) = 4 \quad \text{and} \quad P(E_1) = 4/16 = \frac{1}{4}$$

The triangles enclosing 6+6 dots not on the diagonal are Possible outcomes of E<sub>2</sub>.

$$\therefore n(E_2) = 12 \quad \text{and} \quad P(E_2) = 12/16 = 3/4$$

By using a possibility diagram, we can easily visualize and calculate the probability of combined events. This method is especially helpful when dealing with sequential experiments and conditional probability.


**(b) Probability of Combined Events using Tree Diagram**

**Tree diagram** is a graphical representation of a sequence of events or decisions, showing the possible outcomes and their probabilities in the form of:

**Root Node** (starting point)      **Branches** (representing possible outcomes or decisions)

**Leaf Nodes** (representing the final outcomes)

Tree diagrams are used to visualize probability problems, identify possible outcomes, calculate probabilities, analyze decision-making processes and show conditional probability.



### **Example:**

Maria and Haleema toss 2 coins Haleema after Maria.




**Solution:**

- (a) Tree diagram

(b) Let  $E_1$  be the probability of both heads up, then obviously there is only one such outcome among total of 4 outcomes. i.e.  $E_1 = \{\text{HH}\}$   
 $\therefore P(E_1) = 1/4$

(c) Let  $E_2$  be the probability of Maria's coin head up and Haleema's coin tail up, then obviously there is only one such outcome among total of 4 outcomes.  
i.e.  $E_2 = \{\text{HT}\}$   
 $\therefore P(E_2) = 1/4$



### (c) Probability of Combined Events using Venn Diagram

A Venn diagram is a pictorial representation of sets and their relationships, using overlapping /disjoint circles to represent the intersection of sets. It can be used to calculate the probability of combined events by identifying the regions that represent the favorable outcomes and calculating their probabilities.

**Example:**

Zainab randomly picks a book from a library shelf containing 50 books, as shown in fig:

Find probabilities:


- (a) It is grade 7 book      (b) It is an English book  
(c) It is an English book of grade 7

**Solution:**

- (a) The probability of picking a grade 7 book:  
 $P(\text{grade 7 book}) = 5/50 = 1/10$

(b) The probability of picking an English book  
 (given that it's a grade 7 book):  
 $P(\text{English book from grade 7 books}) = 2/5$

(c) To find the probability of picking a book that is both English and grade 7,  
 multiply the above probabilities:  
 $\therefore P(E) = (1/10) \times (2/5) = 1/25$



So, the probability of picking a book randomly from the 50 books that is a grade 7 English book is  $\frac{1}{25}$  or 4%.

By using a Venn diagram, we can easily visualize and calculate the probability of combined events. This method is especially helpful when dealing with multiple events and their intersections.

## Probability Rules

### Addition Rule

The probability of two mutually exclusive events is the sum of their individual probabilities.

### Multiplication Rule

The probability of two or more events occurring independently is the product of their individual probabilities.

### Complement Rule

The probability of an event not occurring is 1 minus the probability of the event occurring.

### Addition Rule of Probability for Mutually Exclusive Events

The addition rule of probability states that the probability of two or more mutually exclusive events occurring is the sum of their individual probabilities.

$$P(A \text{ or } B) = P(A) + P(B), \quad \text{where A and B are mutually exclusive events.}$$

Applying addition rule of probability to problems of rolling even and odd number on a dice

The addition rule of probability can be applied to solve problems involving even and odd numbers rolled on a dice by treating the outcomes as mutually exclusive events

#### Example:

Hajrah rolled a fair cubical dice once. Find the probability of rolling an even or an odd number.

#### Solution:

On a fair cubical dice, there are 3 even numbers: 2, 4, 6 and 3 odd numbers: 1, 3, 5

Since the dice has 6 faces, the total number of possible outcomes is 6.

The probability of rolling an even number or an odd number is:

$$P(\text{Even}) = 3/6 = 1/2 \quad \text{and} \quad P(\text{Odd}) = 3/6 = 1/2$$

Since the events are mutually exclusive (a roll can't be both even and odd), we can add the probabilities:

$$P(\text{Even or Odd}) = P(\text{Even}) + P(\text{Odd}) = 1/2 + 1/2 = 1$$

Applying addition rule of probability to problems involving left and right hand turns

The addition rule of probability can be applied to solve problems involving taking left or right turns, by treating the outcomes as mutually exclusive events.

#### Example:

Mr. Bilal can make either a left turn (L) or a right turn (R) at an intersection on his journey to Tiri Mengal. The probabilities of making a left turn and a right turn are:

$P(L) = 0.3$  (30%) &  $P(R) = 0.7$  (70%). Find the probability of making either a left or a right turn.

#### Solution:

Since making a left turn and making a right turn are mutually exclusive events (he can't make

both at the same time), so

we can apply the addition rule to find the probability of making either a left turn or a right turn:

$$P(L \text{ or } R) = P(L) + P(R) = 0.3 + 0.7 = 1$$

**Applying addition rule of probability to problems involving tossing a coin**

The addition rule of probability can be applied to solve problems involving tossing a coin by treating the outcomes as mutually exclusive events.

**Example:**

Sunayna tossed a fair coin with two outcomes: Heads (H) and Tails (T). The probabilities are:

$$P(H) = 0.5 \text{ (50\% chance of getting Heads)} \quad \text{and} \quad P(T) = 0.5 \text{ (50\% chance of getting Tails)}$$

find the probability of getting either Head or Tail.

**Solution:**

Since getting Head and getting Tail are mutually exclusive events (the coin can't land on both at the same time), so

we can apply the addition rule to find the probability of getting either Heads or Tails:

$$P(H \text{ or } T) = P(H) + P(T) = 0.5 + 0.5 = 1$$

**Applying addition rule of probability to problems involving winning or losing**

The addition rule of probability can be applied to solve problems involving winning or losing by treating the outcomes as mutually exclusive events.

**Example:**

Affan played a game with two possible outcomes L and W such that  $P(W) = 0.4$  (40%) and  $P(L) = 0.6$  (60%). Find the probability of either winning or losing the game.

**Solution:**

Since winning and losing are mutually exclusive events (he can't both win and lose at the same time), we can apply the addition rule to find the probability of either winning or losing:

$$P(W \text{ or } L) = P(W) + P(L) = 0.4 + 0.6 = 1$$

### Multiplication Rule of Probability for Independent and Dependent Events

The multiplication rule of probability states that the probability of two or more events occurring is the product of their individual probabilities.

(i) For independent events:

$$P(A \text{ and } B) = P(A) \times P(B)$$

(ii) For dependent events:

$$P(A \text{ and } B) = P(A) \times P(B|A); \text{ where } P(B|A) \text{ is the conditional probability of } B \text{ given } A.$$

**Example: (Independent Events)**

Haneen flipped a fair coin twice. What is the probability of getting heads both times?

**Solution:**

$$P(\text{Heads on first flip}) = \frac{1}{2} \quad \text{and} \quad P(\text{Heads on second flip}) = \frac{1}{2}$$

Since the coin flips are independent, we can apply the multiplication rule:

$$\therefore P(\text{Heads on both flips}) = P(\text{Heads on first flip}) \times P(\text{Heads on second flip}) = \frac{1}{2} \times \frac{1}{2} = \frac{1}{4}$$

**Example: (Dependent Events)**

Zeemal drew 2 cards from a standard deck of cards. What is the probability of drawing two

spades in a row?

**Solution:**

$$P(\text{First card is spade}) = 13/52 = 1/4$$

$$P(\text{Second card is spade} \mid \text{First card is spade}) = 12/51$$

Since the events are dependent, we can apply the multiplication rule with conditional probability

$$\begin{aligned} P(\text{Two spades in a row}) &= P(\text{First card is spade}) \times P(\text{Second card is spade} \mid \text{First card is spade}) \\ &= 1/4 \times 12/51 = 3/51 \end{aligned}$$

### Multiplication Rule of Probability Involving Trade

The multiplication rule of probability can be applied to solve problems involving trade by treating the outcomes as independent events.

**Example:**

Mahmil made 2 trades, as given:

Trade 1: Buying Mangoes (BM) and Trade 2: Selling Berries (SB)

With the probabilities:

$$P(BM) = 0.6 \quad \text{and} \quad P(SB) = 0.7$$

Find the probability of both trades being successful.

**Solution:**

Since the trades are independent (the outcome of one trade doesn't affect the other), we can apply the multiplication rule to find the probability of both trades being successful:

$$P(BM \text{ and } SB) = P(BM) \times P(SB) = 0.6 \times 0.7 = 0.42$$

This means there is a 42% chance of both trades being successful.

### Multiplication Rule of Probability Involving Flipping a Coin

The multiplication rule of probability can be applied to solve problems involving flipping a coin by treating the outcomes as independent events.

**Example:**

Azkaa flipped a biased coin twice with following outcomes:

Flip 1: Heads (H) or Tails (T) and Flip 2: Heads (H) or Tails (T)

With the probabilities:

$$P(H) = 0.7 \quad \text{and} \quad P(T) = 0.3$$

Find the probability of:

- i. Both heads
- ii. First head then tail
- iii. First tail then head
- iv. Both tails

**Solution:**

Since the coin flips are independent (the outcome of one flip doesn't affect the other), we can apply the multiplication rule to find the probability of specific outcomes:

- i.  $P(HH) = P(H) \times P(H) = 0.7 \times 0.7 = 0.49$  (49% chance of getting Heads both times)
- ii.  $P(HT) = P(H) \times P(T) = 0.7 \times 0.3 = 0.21$  (21% chance of getting Heads then Tails)
- iii.  $P(TH) = P(T) \times P(H) = 0.3 \times 0.7 = 0.21$  (21% chance of getting Tails then Heads)
- iv.  $P(TT) = P(T) \times P(T) = 0.3 \times 0.3 = 0.09$  (9 % chance of getting Tails both times)

### Multiplication Rule of Probability for Drawing Cards with Replacement

The multiplication rule of probability can be applied to solve problems involving drawing 2

cards with replacement by treating the outcomes as independent events.

### Example:

Romaiza draws 2 cards from standard deck with replacement as given :

First draw → Any card (52 possibilities)

Second draw → Any card (52 possibilities, since the first card is replaced)

Find the probabilities of:

- i. Both black cards
- ii. Both hearts
- iii. First spade then heart
- iv. First Queen of Spades then King of Hearts

### Solution:

Since the events are independent (the outcome of the second draw does not depend on the first draw), we can apply the multiplication rule to find the probability of specific outcomes:

$$\begin{aligned} \text{i. } P(\text{Both black cards}) &= P(\text{black}) \times P(\text{black}) = 26/52 \times 26/52 = 1/4 \\ \text{ii. } P(\text{Both hearts}) &= P(\text{heart}) \times P(\text{heart}) = 13/52 \times 13/52 = 1/16 \\ \text{iii. } P(\text{spade then heart}) &= P(\text{spade}) \times P(\text{heart}) = 13/52 \times 13/52 = 1/16 \\ \text{iv. } P(\text{Queen of Spades then King of Hearts}) &= P(\text{Queen of Spades}) \times P(\text{King of Hearts}) \\ &= 1/52 \times 1/52 = 1/1704 \end{aligned}$$

### Multiplication Rule of Probability for Drawing Cards without Replacement

The multiplication rule of probability can be applied to solve problems involving drawing 2 cards without replacement by treating the outcomes as dependent events.

### Example:

Sundas draws 2 cards one after the other from a standard pack of playing cards, such that:

First draw → Any card (52 possibilities)

Second draw → Any card (51 possibilities, since one card has already been drawn)

then find the probabilities of:

- i. Both black cards
- ii. Both hearts
- iii. First spade then heart
- iv. First Queen of Spades then King of Hearts

### Solution:

Since the events are dependent we can apply the multiplication rule to find the probability :

$$\begin{aligned} \text{i. } P(\text{Both black cards}) &= P(\text{black}) \times P(\text{black/black}) = 26/52 \times 25/51 = 25/102 \\ \text{ii. } P(\text{Both hearts}) &= P(\text{heart}) \times P(\text{heart/heart}) = 13/52 \times 12/51 = 3/51 \\ \text{iii. } P(\text{spade then heart}) &= P(\text{spade}) \times P(\text{heart/spade}) = 13/52 \times 13/51 = 13/204 \\ \text{iv. } P(\text{first Queen of Spades then King of Hearts}) &= P(\text{Queen of Spades}) \times P(\text{King of Hearts/ Queen of Spades}) = 1/52 \times 1/51 = 1/2652 \end{aligned}$$

### Exercise 12.4

1. Musaab picked a card randomly from a box containing cards bearing numbers 1, 2 and 3. He then replaced and picked another after well shuffling. He found product of two numbers he got in 2 trials. Draw the possibility diagram of this experiment and find probability of getting an even product.

2. Ashas flipped a coin thrice. Draw tree diagram to find the probability of getting three Tails?
3. A survey of 100 students shows that 40 like pizza, 45 like burgers. Draw venn diagram and find probability of students like either pizza or burgers, assuming none like both?
4. What is the probability of a patient testing positive for a disease with either Test A or Test B, given that the probability of testing positive with Test A is 0.3 and the probability of testing positive with Test B is 0.4, and the tests are independent?
5. What is the probability of getting either Job A or Job B, given that the probability of getting Job A is 0.5 and the probability of getting Job B is 0.4 and the offers are independent?
6. What is the probability of taking either Volvo bus or Bedford bus to work, given that the probability of taking Volvo bus is  $\frac{1}{2}$  and the probability of taking Bedford bus is  $\frac{1}{3}$ , and the buses are to be chosen independently?
7. What is the probability of either Team Sudaisee or Team Huzaifee winning the championship, given that the probability of Team Sudaisee winning is  $\frac{12}{27}$  and the probability of Huzaifee team winning is  $\frac{13}{27}$ , and the teams play independently?
8. HJ produces party abayaas. In the products, probability of colour damage is 0.1 and the probability of bad stitching is 0.2. Find the probability of a product having either colour damage or bad stitching.
9. Fakaha is tested for a disease with Test A and Test B and the probability of testing positive with A is 0.9 and the probability of testing positive with B is 0.7. Find the probability of Fakehaa for testing positive with both Test A and Test B, when the tests are independent?
10. Cocobakes produces festive cakes. The quality control department performs 2 quality checks A and B on an item. Find the probability of a product passing both Quality Control Test A and Quality Control Test B, given that the probability of passing Test A is 0.9 and the probability of passing Test B is 0.8, and the tests are independent?

### I have Learnt

- A table consisting of values of a data along with their frequencies is called a frequency table.
- In a group data, the table consisting of class intervals and their respective frequencies is called frequency table or a frequency distribution.
- The representation of data by a polygon in which class marks are plotted along x-axis and frequencies along y-axis is called a frequency polygon.
- In a group data, number of values less than or within the limits of a class is called cumulative frequency of the respective class.
- A polygon made by joining the points whose x-coordinates are class marks and y-coordinates are their cumulative frequencies is called a cumulative frequency polygon.

- An ogive is a smooth curve, while a cumulative frequency polygon is a polygon-shaped graph made by straight segments.
  - The three values which divide the whole data into four equal parts are called quartiles.
  - Deciles divide a dataset or a distribution into 10 equal parts, each representing 10% of the data.
  - Percentiles divide a dataset or a distribution into 100 equal parts, each representing 1% of the data.
  - $IQR = Q_3 - Q_1$ , where IQR represents the spread of the data within the middle half of the distribution.
  - A box-and-whisker plot is a graphical representation of a dataset that displays the distribution of values through 5 data points: min, max,  $Q_1$ ,  $Q_2$ ,  $Q_3$
  - A single value which can represent the whole data is called measure of central tendency, measure of location or an average.
  - The middle most value of an arranged data is called median average or simply median.
  - The measures which tell us how the values of a data are scattered around their average, are called measures of dispersion.
  - Range is the difference of largest and smallest values of the data.
  - Variance is the ratio between sum of squares of deviations from mean and the number of values.
  - Standard deviation is the positive square root of variance.
  - Probability is a measure of the likelihood of an event occurring.
  - The percentage ratio between the standard deviation and mean of a data is called coefficient of variation i.e.  $C.V. = \frac{SD}{Mean} \times 100$


## MISCELLANEOUS EXERCISE 12

**1. Encircle the correct option in the following.**



**5.** Given box and whisker plots show the marks of 4 boys of mauve section and 4 boys of hazel section in math test.

- a) Find Median scores of both sections
- b) Find IQR of scores of both sections



- 6.** Bareera and Hadiya draw cookie packs, Hadiya after Bareera from a bag of 2 choco cookies and 3 lemon cookies. Find the probability of drawing both lemon cookies
- 7.** A company has two machines, A & B, which produce 60% & 40% of the total products, respectively. Machine A produces 20% defective products & Machine B produces 15% defective products. What is the probability that a randomly selected product is defective?

**Maths Play Ground:**

Help students to make colour wheel/ number wheel and let them perform trials of moving the wheel to record the outcomes. Then ask them different related questions of probability.

# ANSWERS

## UNIT 1

**Exercise 1.1**

1. (i)  $\sqrt{3}i$  (ii)  $12i$  (iii)  $\frac{2}{3}i$  (iv)  $-2\sqrt{5}i$  (v)  $4 - 2\sqrt{15}i$  (vi)  $-4$
2. (i)  $9+4i$  (ii)  $2$  (iii)  $-9+5i$  (iv)  $-1$  (v)  $-6+8i$  (vi)  $-10-11i$   
 (vii)  $-16-16i$  (viii)  $5+12i$  (ix)  $45$  (x)  $16+30i$   
 (xi)  $1+\sqrt{6}+(\sqrt{2}-\sqrt{3})i$  (xii)  $3$
3. (i)  $i$  (ii)  $i$  (iii)  $1$  (iv)  $-i$  (v)  $-27i$  (vii)  $16$
4. (i)  $8+0=8$  (ii)  $-17+i$  (iii)  $1-13i$  (iv)  $22i$  (v)  $2i$  (vi)  $-2$
5. (i)  $\frac{12}{17} + \frac{3}{17}i$  (ii)  $-\frac{15}{16} + \frac{18}{61}i$  (iii)  $\frac{2}{7} - \frac{3\sqrt{5}}{7}i$  (iv)  $\frac{7}{5} - \frac{2}{5}i$  (v)  $-\frac{9}{41} + \frac{40}{41}i$   
 (vi)  $\frac{8}{5} + \frac{1}{5}i$  (vii)  $\frac{7}{5} - \frac{9}{5}i$  (viii)  $\frac{a^2-b^2}{a^2+b^2} + \frac{2ab}{a^2+b^2}i$  (ix)  $-\frac{1}{2} + \frac{1}{2}i$  (x)  $4$

**Exercise 1.2**

1. (a)  $4-5i$  (b)  $3+3i$  (c)  $-5+5i$  (d)  $-4i$
3. (a)  $\frac{1}{2} - \frac{1}{2}i$  (b)  $\frac{7}{58} + \frac{3}{58}i$  (c)  $\frac{5}{122} + \frac{3}{61}i$  (d)  $\frac{5}{2} - \frac{1}{2}i$  (e)  $3+2i$  (f)  $\frac{a}{a^2+b^2} + \frac{b}{a^2+b^2}i$
4. (a)  $16$  (b)  $2$  (c)  $49$  (d)  $40$  (e)  $181$  (f)  $137$
6. Do yourself 5. b. (i)  $\sqrt{10}$  (ii)  $\sqrt{34}$  (iii)  $1$
7. (a)  $\text{Re}=-1, \text{Im}=-2\sqrt{6}$  (b)  $\text{Re}=1, \text{Im}=2\sqrt{2}$  (c)  $\text{Re}=\frac{41}{10}, \text{Im}=-\frac{3}{10}$  (d)  $\text{Re}=-\frac{1}{2}, \text{Im}=\frac{\sqrt{3}}{2}$   
 (e)  $\text{Re}=-1, \text{Im}=1$  (f)  $\text{Re}=\frac{1}{2}, \text{Im}=0$  (g)  $\text{Re}=\frac{-8}{25}, \text{Im}=-\frac{6}{25}$

**Exercise 1.3**

1.  $\{\pm\sqrt{7}i\}$  2.  $\{\pm 3i\}$  3.  $\{\pm 10i\}$  4. yes 5. yes 6. No 7. Yes 8.  $(x-4i)(x+4i)$
9.  $(a-bi)(a+bi)$  10.  $(x-5yi)(x+5yi)$  11.  $W=1-i, Z=4-i$
12.  $W=3-2i, Z=1$  13. (a) (i)  $I = 9 + \frac{37}{4}J$  (ii)  $I = \frac{-37}{5} + \frac{2}{5}J$  (b) (i)  $Z = -\frac{5}{2} - 10J$   
 (ii)  $Z = -\frac{770}{73} - \frac{380}{73}J$  (c)  $\frac{225}{41} + \frac{200}{41}i$

**Miscellaneous Exercise-1**

1. (i) (c) (ii) (b) (iii) (a) (iv) (c) (v) (b) (vi) (c)  
 (vii) (d) (viii) (b) (ix) (c) (x) (d)
2. a.  $-10+9i$  b.  $1-3i$  3. (a)  $2x-3y+(3x+2y)i$  (b)  $-45i$
4. (a)  $3\sqrt{2}-\sqrt{7}i$  (b)  $\sqrt{2}i$  5. (a)  $\frac{5}{4}$  (b)  $245$  6. (a)  $\frac{-4}{5} - \frac{-3}{5}i$  (b)  $\frac{3\sqrt{5}}{5} - \frac{\sqrt{5}}{5}i$
7. a.  $2(x-3i)(x+3i)$  b.  $-(x-5i)(x+5i)$
8. a.  $3(x-\sqrt{5}i)(x+\sqrt{5}i)$  b.  $6(x-\sqrt{6}i)(x+\sqrt{6}i)$

## UNIT 2

**Exercise 2.1**

1. (i)  $x^2-x-11=0$  (ii)  $3x^2+26x-2=0$  (iii)  $x^2-2x=0$
2. (i)  $\{1, 4\}$  (ii)  $\{1\}$  (iii)  $\{-1, 8\}$  (iv)  $\{3, 5\}$  (v)  $\{3, -10\}$
3. (i)  $\{4, -8\}$  (ii)  $\{0, -8\}$  (iii)  $\{-3 \pm 3\sqrt{2}\}$  (iv)  $\left\{ \frac{-2\sqrt{3} \pm 2}{\sqrt{3}} \right\}$  (v)  $\left\{ \frac{-1 \pm \sqrt{-3}}{2} \right\}$  (vi)  $\left\{ \frac{5}{2}, -\frac{1}{2} \right\}$

4. (i)  $\{3, -3\}$  (ii)  $\left\{\frac{-5 \pm \sqrt{17}}{4}\right\}$  (iii)  $\{-1, 24\}$  (iv)  $\{0, 2\}$  (v)  $\left\{1, -\frac{3}{2}\right\}$  (vi)  $\left\{\frac{-11 \pm \sqrt{165}}{2}\right\}$

5.  $\{-3, -3\}$  6.  $-1 \pm \sqrt{3}i$  8. Only two solutions, degree is 2

9. The width of the strips added is approximately 16.85 meters.

The new dimensions of the playground are approximately 76.85 meters by 46.85 meters.

10. It takes approximately 5.65 seconds for the car to travel 145 meters.

### Exercise 2.2

1. (i)  $ay^2 + by + c = 0$  (ii)  $9y^2 - 3y + 7 = 0$  (iii)  $y^2 + 1 = 0$  (iv)  $z^2 - 4z + 3 = 0$   
 (v)  $3y^2 + 8y + 5 = 0$  (vi)  $3z^2 + 7z + 11 = 0$  (vii)  $az^2 + bz + c = 0$   
 (viii)  $8y^2 - 7y - 1 = 0$  (ix)  $y^2 - 16y - 42 = 0$  (x)  $4z^2 + 19z + 13 = 0$

2. (i)  $\{\pm 2, \pm 4\}$  (ii)  $\{\pm 3, \pm 5i\}$  (iii)  $\{-32, -243\}$  (iv)  $\left\{\pm 1, \pm \frac{2}{\sqrt{3}}\right\}$  (v)  $\left\{0, -\frac{2}{5}\right\}$

(vi)  $\left\{\pm \sqrt{\frac{14}{5}}, \pm \sqrt{\frac{-2}{5}}\right\}$  (vii)  $\left\{-1, 2, \frac{1}{2}\right\}$  (viii)  $\left\{3, -4, \frac{1}{3}, -\frac{1}{4}\right\}$

(ix)  $\left\{\frac{1 \pm \sqrt{5}}{2}, \frac{-3 \pm \sqrt{73}}{8}\right\}$  (x)  $\{1, 5\}$  (xi)  $\{1, 2\}$  (xii)  $\{2, 3\}$

(xiii)  $\left\{1, -8, \frac{-7 \pm \sqrt{97}}{2}\right\}$  (xiv)  $\{2, -4, -1 \pm 2i\}$  (xv)  $\{-1 \pm \sqrt{17}, -1 \pm \sqrt{41}\}$

3. (i)  $\left\{\pm 1, \pm \frac{1}{\sqrt{2}}\right\}$  (ii)  $\left\{1, -\frac{1}{2}\right\}$  (iii)  $\{\pm 1\}$  (iv)  $\{-1\}$

4. do yourself 5. Take  $z = y - 4$ , The solution  $y = 10, y = 5$  6.  $\left\{5, \frac{-5+5\sqrt{3}i}{2}, \frac{-5-5\sqrt{3}i}{2}\right\}$

### Exercise 2.3

1. i) 144 ii) 49 iii) -28 iv) 841  
 2. i) Rational and unequal ii) Rational and unequal  
 iii) Rational and unequal iv) Imaginary  
 v) Irrational and unequal

3.  $k = \pm 24$  4.  $k = \pm 6$

6. i)  $m = 9$  ii)  $m = -1/4$  iii)  $m = -1, -11/3$

### Exercise 2.4

1. i)  $S = 5, P = 2$  ii)  $S = -3/2, P = 1/2$  iii)  $S = 2/5, P = 2/5$   
 iv)  $S = -2, P = 9/4$  v)  $S = 17/16, P = -3/4$  vi)  $S = 25.67, P = 6.0$

2. i)  $3x^2 + 5x - 8 = 0$  ii)  $x^2 - 3\sqrt{3}x + 6 = 0$  iii)  $x^2 - 4x + 1 = 0$   
 iv)  $x^2 + 25 = 0$  v)  $x^2 - 14x + 53 = 0$

3. i)  $-5/4$  ii)  $-20/9$  iii)  $16/3$  iv)  $1/2$   
 v)  $-5/3$  vi)  $8/9$  vii)  $-80/27$  viii)  $25/3$

4. i)  $49x^2 - 2x + 49 = 0$  ii)  $7x^2 + 10x + 7 = 0$  iii)  $49x^2 - 2x + 49 = 0$   
 iv)  $49x^2 + 96 = 0$  v)  $7x^2 + 6x + 15 = 0$  vi)  $49x^2 - 2x + 49 = 0$   
 vii)  $49x^2 + 143x + 100 = 0$  viii)  $49x^2 + 100x + 100 = 0$  ix)  $x^2 + 68x/49 + 359/343 = 0$

5.  $x^2 - 60x + 864 = 0$  6.  $2x^2 + 38x - 7 = 0$  7.  $k = \pm \sqrt{5}$  8.  $k = 4$

### Exercise 2.5

1. S.S. =  $\{(-1, 3), (9/5, -13/5)\}$
2. S.S. =  $\{(1, 1), (-5, -8)\}$
3. S.S. =  $\{(2, -9), (-3, 6)\}$
4. S.S. =  $\{(1/2, 7/2)\}$
5. S.S. =  $\{(9/\sqrt{2}, 3/\sqrt{2}), (-9/\sqrt{2}, -3/\sqrt{2}), (3\sqrt{5}/2, 3\sqrt{5}/2), (-3\sqrt{5}/2, -3\sqrt{5}/2)\}$
6. S.S. =  $\left\{\left(\pm\sqrt{6}, \pm\sqrt{6}\right), \left(\sqrt{\frac{5}{2}}, -6\sqrt{\frac{5}{2}}\right), \left(-\sqrt{\frac{5}{2}}, 6\sqrt{\frac{5}{2}}\right)\right\}$
7. S.S. =  $(-1/2, -4), (2, 7/2)$
8. S.S. =  $\{(-1, -1), (2, -4)\}$
9. S.S. =  $\{\pm 1/\sqrt{2}, \pm 3/\sqrt{2}\}$
10. S.S. =  $\{(\pm 3, \pm\sqrt{2})\}$

### Exercise 2.6

1. 10, 12
2. 20
3. 6, 5
4. (3, 6) (6, 3)
5. 5, 6
6. 3, 5
7. Son's age = 13, Father's age = 37
8. 83
9. (24, -4)(15, 5)
10. 4, 6
11. 625, 400
12. 8, 9
13. Appro 4.1 seconds
14. 20.82 Km/h
15. 11 cm long and 10 cm wide

### Miscellaneous Exercise-2

1. (i) (c) (ii) (d) (iii) (b) (iv) (b) (v) (a) (vi) (a)  
(vii) (d) (viii) (b) (ix) (c) (x) (b) (xi) (c) (xii) (d) (xiii) (a)
2.  $\left\{1, -\frac{1}{2}, \frac{-1 \pm \sqrt{-3}}{2}, \frac{-1 \pm \sqrt{-3}}{4}\right\}$
3.  $m = 3/2, x = -3$
4.  $\frac{c+3(b+3a)}{a}$
5.  $a = \pm 2\sqrt{2}, b = -2$
6.  $L = x + 1$ , then  $W = x + 3$

### Exercise 3.1

1. (a) 1, 2, 1-by-2 (b) 2, 2, 2-by-2 (c) 1, 1, 1-by-1 (d) 1, 1, 1-by-1
2. (a) not equal (b) not equal (c) not equal (d) equal
3.  $R = \begin{bmatrix} 7 & 11 & 10 \end{bmatrix}, C = \begin{bmatrix} 7 \\ 11 \\ 10 \end{bmatrix}$
4. (a) scalar matrix (b) square matrix (c) unit matrix (d) unit matrix (e) scalar matrix
5. (a) symmetric matrix (b) skew symmetric matrix (c) skew symmetric matrix  
(d) skew symmetric matrix (e) symmetric matrix

### Exercise 3.2

1. (a)  $x = 2, y = 9$  (b) impossible, corresponding elements are not equal. (c) impossible; orders are not equal  
(d)  $x = -4, y = -2, z = -8$  (e)  $x = 3, y = 3, z = 6$  (f) impossible; orders are not equal.  
(g)  $x = 6, y = 5$  (h)  $x = 2, y = 5, z = 10$  (i)  $x = 4, y = 2$  (j)  $x = -10, y = -6, z = -4$

2.  $-R = \begin{bmatrix} -5 & 0 & -3 \\ -7 & 9 & 1 \\ 8 & -5 & -6 \end{bmatrix}, -S = \begin{bmatrix} 5 & -2 \\ -3 & 6 \\ 9 & -4 \end{bmatrix}, -T = \begin{bmatrix} -5 & 6 & -1 \end{bmatrix}$
3. (i)  $\begin{bmatrix} 20 & 12 \\ -8 & 11 \end{bmatrix}$  (ii)  $\begin{bmatrix} 0 & -1 \\ 2 & 3 \end{bmatrix}$

4. (i) yes (ii) no (iii) no (iv) no
8. (i)  $Z = \begin{bmatrix} 1 \\ 7 \end{bmatrix}$  (ii)  $Z = \begin{bmatrix} 5 \\ -2 \end{bmatrix}$

9. (a) (i) impossible (ii) 1-by-2 (iii) 2-by-1 (iv) impossible (v) impossible (vi) impossible  
(vii) 2-by-1 (viii) impossible (ix) 1-by-2 (x) impossible (xi) 1-by-1 (xii) 1-by-1  
(xiii) 2-by-2 (xiv) 2-by-2 (xv) 2-by-1

- (b) (ii)  $[8 \ 0]$  (iii)  $\begin{bmatrix} 8 \\ 0 \end{bmatrix}$  (vii)  $\begin{bmatrix} 5 \\ 15 \end{bmatrix}$  (ix)  $[45 \ 15]$  (xi)  $[50]$  (xii)  $[20]$  (xiii)  $\begin{bmatrix} 10 & 20 \\ 15 & 30 \end{bmatrix}$  (xiv)  $\begin{bmatrix} 2 & 5 \\ 4 & 3 \end{bmatrix}$  (xv)  $\begin{bmatrix} 19 \\ -2 \end{bmatrix}$
10. (i)  $AB = \begin{bmatrix} 2 & 1 \\ 4 & -7 \end{bmatrix}, BA = \begin{bmatrix} 4 & 3 \\ -6 & -9 \end{bmatrix}, AB \neq BA$  (ii)  $AC = CA = \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$

- (vi)  $A^2 = \begin{bmatrix} 1 & 0 \\ 8 & 9 \end{bmatrix}, B^2 = \begin{bmatrix} 4 & -1 \\ 0 & 9 \end{bmatrix}, A + B = \begin{bmatrix} 3 & 1 \\ 2 & 0 \end{bmatrix}, A - B = \begin{bmatrix} -1 & -1 \\ 2 & 6 \end{bmatrix},$

- $(A + B)(A - B) = \begin{bmatrix} -1 & 3 \\ -2 & -2 \end{bmatrix}, A^2 - B^2 = \begin{bmatrix} -3 & 1 \\ 8 & 0 \end{bmatrix}$  (vii) no (viii) no (ix) no (x) yes

### Exercise 3.3

1. (a) A is non singular, B is non singular, C is singular, D is non singular and E is singular. (b)  $k = 1$  (c)  $x = \frac{7}{3}$

2. (a)  $R^{-1} = \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 3 & 2 \end{bmatrix}$ , but  $S^{-1}$ ,  $T^{-1}$  &  $U^{-1}$  are not possible.

4. Do yourself. 5. (i)  $\begin{array}{l} 4x + 2y = 6 \\ 3x + y = 4 \end{array}$  (ii)  $\begin{array}{l} 5x = 10 \\ 4y = 20 \end{array}$  (iii) Not possible (iv)  $5x + 3y = 2$

6. (a) (i)  $\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$  (ii)  $\begin{bmatrix} 2 & 1 \\ -1 & 5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 90 \\ 10 \end{bmatrix}$  (iii)  $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} y \\ x \end{bmatrix} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$  or  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$

(iv)  $\begin{bmatrix} \frac{5}{2} & -3 \\ -4 & \frac{1}{2} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$  (b)  $K = 8$  7. (a) (i)  $\{(-3, 2)\}$  (ii)  $\left\{0, \frac{3}{4}\right\}$  (iii) impossible

(iv) impossible (b) (i)  $\{(4, -1)\}$  (ii)  $\left\{\left(\frac{5}{3}, -\frac{1}{2}\right)\right\}$  (iii)  $\{(20, 10)\}$  (iv) impossible

### Exercise 3.4

1.  $50^\circ, 40^\circ$  2. literature = 5, science = 10 3. Note book = Rs.100, Book = Rs.150

4. Sundas = 15 years, Zenab = 7 years 5. The first truck (with a capacity of 10 tons) made 7 round trips.

The second truck (with a capacity of 12 tons) made 13 round trips. 6. 8 kg of the metal with 55% and 12 kg of the metal with 80% aluminum content. 7. 150 pounds of soybean and 200 pounds of cornmeal

8. Length = 76 meters Width = 19 meters

### Miscellaneous Exercise-3

1. (i) (b) (ii) (d) (iii) (b) (iv) (d) (v) (d) (vi) (d)  
 (vii) (a) (viii) (a) (ix) (d) (x) (c) (xi) (c) (xii) (b)  
 (xiii) (d) (xiv) (c) (xv) (b) (xvi) (d)


3. impossible, matrix of coefficient is singular.

5. impossible, matrix of coefficient is singular.


### UNIT 4

#### Exercise 4.1


1.  $x \leq 4$




3.  $x > -10$



5.  $x \geq -20$



7.  $3x + 4 < 40, x < 12$



9.  $x > 7$  10.  $x \leq \frac{9}{2}$  11.  $x < -1$  or  $x > 3$  12.  $-1 < x < 9$

13.  $x < -9$  or  $x > 10$  14.  $-7 < x < 3$  15.  $x \neq 4$  16.  $4 - 15 < x < 15 + 4$


17.  $10 \leq W \leq 24$  18. There are no solutions except  $-1$  19.  $0 \leq x \leq 24.54$  20.  $L: x \leq 10m, W: 2x \leq 20m$

2. No solution


4.  $x < \frac{55}{7}$



6.  $x < 20$










8.  $2(x + 8) \leq -2, x \leq -9$




**Exercise 4.2**

1. No 2. Yes 3. Yes 4. No 5. Yes 6. No




15. Do Yourself.


16.  $x - 3 \geq y$



17.  $-2y \leq x + 6$




18.  $x + 4y < -2$


**Exercise 4.3**

1.  $\{-8, 14\}$  2.  $\{-\frac{39}{2}, \frac{21}{2}\}$  3. No solution 4.  $\{4\}$  5.  $\{1\}$  6.  $\{-\frac{38}{5}, \frac{46}{5}\}$  7. Absolute must be positive 8. Absolute Deviation =  $\pm 0.5$  psi, Minimum Pressure = 8 psi - 0.5 psi = 7.5 psi Maximum Pressure = 8 psi + 0.5 psi = 8.5 psi


9.  $\{-1 \leq x \leq \frac{11}{9}\}$




10.  $\{3 \leq x \leq 4\}$



11.  $\{x < -8 \text{ or } x > -4\}$




12.  $\{x < 12 \text{ or } x > 16\}$ ,



13.  $\{-\frac{8}{3} \leq x \leq 4\}$


14.  $\{x \leq -\frac{2}{15} \text{ or } x \geq \frac{14}{15}\}$



15.  $\{470 \leq W \leq 530\}$


#### Exercise 4.4

2. a.  $(0,0)$  and  $(-3,1)$  are solutions    b.  $(3,2)$  is solution




c. Both lines are parallel and superpose each other

d. Do Yourself.



f. Do Yourself.




g. Do Yourself.

4. No solution

#### Miscellaneous Exercise-4


1. (i) (d) (ii) (c)  
(vii) (b) (viii) (c)

2.  $x \geq \frac{15}{2}$










3.  $x \geq 3$     4. No solution

5.  $x \geq -1$     6.



7.



14.  $2 \leq x \leq 4$ ,  $1 \leq y \leq 4$ , 15.  $x \geq 3$ ,  $y \geq \frac{2}{3}x - 2$ ,  $y \leq \frac{4}{3}x - 2$

## UNIT 5

### Exercise 5.1

1. (i)  $\frac{3x^2}{5ay^4}$  (ii)  $\frac{2p^2m^2}{3k}$  (iii)  $\frac{nq}{mp^3}$  (iv)  $\frac{1}{5ab}$  (v)  $\frac{2lmn}{3}$  (vi) -1

(vii)  $x - 9$  (viii)  $\frac{r+3}{r-4}$

2. (i) 9 (ii) 294 (iii) 96.6 (iv) -7.5 (v) 1.002 (vi)  $-3\frac{17}{56}$

3. 5050 5.  $-11\frac{2}{3}$  6. 21 cm

### Exercise 5.2

1. (i)  $\frac{7x}{10}$  (ii)  $\frac{11x+2}{18}$  (iii)  $\frac{81x+24}{28}$  (iv)  $\frac{3x+2}{x+5}$  (v)  $\frac{7x+3}{x-3}$

2. (i)  $\frac{x+12}{5}$  (ii)  $\frac{17x-101}{21}$  (iii)  $\frac{-2x^3+2x^2+1}{x}$

3. (i)  $x^2 + 2x - 3$  (ii)  $x^2 - 2xy + y^2$  (iii)  $\frac{6(x-2)}{x}$  (iv)  $\frac{x}{x-5}$

4. (i)  $\frac{47x}{60}$  (ii)  $\frac{29-2x}{6}$  (iii)  $\frac{-4}{x^2-1}$  (iv)  $\frac{3x^2-100}{15}$  (v)  $\frac{9b^2cz^2}{4x^3y}$  (vi)  $\frac{p^2q^2y}{x^2}$  (vii)  $\frac{x+1}{x-3}$   
 (viii)  $\frac{4}{3}$  (ix)  $\frac{12a^2-4a+7}{3(4a^2-9)}$  (x)  $\frac{23x}{(1+2x)(2+x)(5-9x)}$  (xi)  $\frac{1}{8m}$  (xii)  $\frac{1-q}{p}$  (xiii)  $\frac{1}{1+x}$

### Exercise 5.3

1. {2} 2. {2, 3} 3. No solution 4. only 4 (-1 is not solution) 5. {-1, -5}  
 6. {-6, 8}

7. No solution 8. approximately  $2\frac{2}{9}$  or about 2 hours and 13 minutes

9. add 12 pints of yellow paint to the mixture to achieve a paint mixture that is 80% yellow.

10. Wasi takes 36 hours and Waqar takes 45

### Miscellaneous Exercise-5

1.(i)	(b)	(ii)	(a)	(iii)	(d)	(iv)	(c)	(v)	(c)
(vi)	(d)	(vii)	(b)	(viii)	(d)	(ix)	(a)	(x)	(c)
(xi)	(d)	(xii)	(d)	(xiii)	(c)				

2. 78.571 cm 3. (i) 24.64 square inches (ii) 1.75 feet 4. (i), (ii) and (iv)

5. 20 6.  $3x - 9$  7.  $x - 2$  8. (i)  $\frac{7x^2+30x+16}{8-2x^2}$  (ii)  $\frac{1}{3}$

9. (i) {-2, -6} (ii) {5, -6} 3 is not solution 10. Essential Step: Ensure that solutions do not make any denominator zero, as such values are not valid for rational equations.

## UNIT 6

### Exercise 6.1

1. i. Yes	ii. No	iii. No	iv. Yes	2. i. Yes, Bijective	ii. No	iii. Yes, Into
3. i. 6	ii. 9/2	iii. 28/9		iv. $\frac{1}{2}(2t^2 + 3t + 8)$		
4. Injective.	5. $a = t = 1$	6. Yes				


### Exercise 6.2

1. i.  $x^2 + 2x - 3$
- ii.  $-x^2 + 6x - 5$
- iii.  $4(x - 1)^3$
- iv.  $\frac{4}{x-1}$
2. i.  $4(x + 1)$
- ii.  $4x + 1$
- iii.  $16x$
- iv.  $x + 2$
3. i.  $f \circ g(x) = -2x + 1$
- ii.  $f \circ g(x) = \frac{x-1}{x}$
- iii.  $f \circ g(x) = \frac{6}{\sqrt{x-1}}$
- iv.  $f \circ g(x) = x - 2$
- $g \circ f(x) = -2x + 4$
- $g \circ f(x) = \frac{4}{2-x}$
- $g \circ f(x) = \frac{2}{\sqrt{3x-1}}$
- $g \circ f(x) = \sqrt{x^2 - 2}$
4. i.  $x = 0, -2$
- ii.  $x = 1 \pm \sqrt{2}$
- iii.  $x = \pm 2$
5. i.  $f^{-1}(x) = \frac{1+x}{2}$
- ii.  $f^{-1}(x) = \frac{2+3x}{x}$
- iii.  $f^{-1}(x) = x^2 - 5$
- iv.  $3 \pm \sqrt{x}$
6. i.  $f^{-1}(1) = 8$ , ii.  $g^{-1}\left(\frac{1}{2}\right) = -1$


### Exercise 6.3

1. (i), (iii), (iv) and (iv) are graphs of functions.


2. i. linear




i. linear




iii. not linear



3. (i)



(iv)



(ii), (iii), (v) draw yourself

4. i. upwards, x-intercept: 1, y-intercept: 1  
intercept: 2


ii. downwards, x-intercepts:  $\pm\sqrt{2}$ , y-

iii. downwards, x-intercept:  $2 \pm \sqrt{3}$ , y-intercept: 2  
intercept: 3


iv. downwards, x-intercepts: -1, 3, y-

5.  $h=4/9$ ,  $k=2$

6. i. draw yourself ii.



7.



8. 63814 (approx.)


Supply will become equal to demand at  $x = 50$ . 9. draw yourself 10. 175 dollars

### Miscellaneous Exercise-6

1. i. b ii. c iii. c iv. a v. c vi. c vii. b viii. c ix. d x. b

2.  $f^{-1}(x) = \frac{x}{2x-3}$ ,  $f^{-1}(x)$  is undefined at  $x = \frac{3}{2}$ ,  $f^{-1}(-1) = \frac{1}{5}$

3. downwards



4.  $y = -10x^2 + 10$


5.  $x = -2, 3$

6.  $F = \frac{9}{5}C + 32$ ,  $77^\circ F$

7.  $(1, -1), (4, 2)$

8. 17 hrs.(approx.)


9. 2 units




### UNIT 7

#### Exercise 7.1


1. (i)




(ii)




(iii)




(iv)



2.



3.



4.  $\vec{b} = -0.5\vec{a}$ ;  $\vec{c} = 2\vec{a}$ ;  $\vec{d} = -\vec{a}$ ;  $\vec{e} = 3\vec{a}$ ;  $\vec{f} = 0.5\vec{a}$ ;  $\vec{g} = -2\vec{a}$

5.  $(5, -10)$  6.  $[-1, 1]$

7. (i)  $A'(1, 6), B'(4, 4) C'(-1, 1)$  (ii)  $A'(-7, 2), B'(-4, 0), C'(-9, -3)$  8.  $[6, -5]$

9. (i)  $\vec{a}$  (ii)  $-b$  (iii)  $\vec{a} + b$  (iv)  $-\vec{a} + b$  (v)  $\frac{1}{2}(\vec{a} + b)$  (vi)  $\frac{1}{2}(-\vec{a} + \vec{b})$  (vii)  $2\vec{a} + \vec{b}$

10. (i)  $-2\vec{q}$  (ii)  $-\vec{q}$  (iii)  $\vec{q}$  (iv)  $\vec{p}$

11. (i)  $[3, 3] = 3\hat{i} + 3\hat{j}$  (ii)  $[4, -10] = 4\hat{i} - 10\hat{j}$  (iii)  $[-5, -2] = -5\hat{i} - 2\hat{j}$

12. (i)  $\frac{3}{5}\hat{i} + \frac{4}{5}\hat{j}$  (ii)  $\frac{5}{13}\hat{i} - \frac{12}{13}\hat{j}$  (iii)  $\frac{2}{5}\hat{i} - \hat{j}$

13.  $x = 2, y = -2$

14. (i)  $-6\hat{i} - 3\hat{j}$  (ii)  $-3\hat{i} - 2\hat{j}$  (iii)  $12\hat{i} - 7\hat{j}$  (iv)  $-\frac{3}{2}\hat{i} + 16\hat{j}$  (v)  $-5\hat{i} + 4\hat{j}$

15. (i)  $7\hat{i} - 3\hat{j}$  (ii)  $22\hat{j}$  (iii)  $40\hat{i} + 30\hat{j}$  (iv)  $\frac{1}{2}\hat{i} + \frac{9}{2}\hat{j}$  (v)  $\sqrt{26}$

(vi)  $\sqrt{26}$  (vii) 14.8 (viii) 1.4 (ix) 16.  $p = 2$

17. (i)  $\frac{1}{\sqrt{5}}\hat{i} - \frac{2}{\sqrt{5}}\hat{j}$  (ii)  $\frac{3}{\sqrt{130}}\hat{i} - \frac{11}{\sqrt{130}}\hat{j}$  (iii)  $\frac{1}{\sqrt{5}}\hat{i} - \frac{2}{\sqrt{5}}\hat{j}$  (iv)  $\frac{3}{\sqrt{10}}\hat{i} - \frac{1}{\sqrt{10}}\hat{j}$

(v)  $\hat{i}$  (vi)  $\frac{7}{\sqrt{130}}\hat{i} - \frac{9}{\sqrt{130}}\hat{j}$  (vii) 18. (i)  $-4\hat{j}$  (ii)  $-3\sqrt{2}(\hat{i} + \hat{j})$

19. 383.28 mph 20. 9.01 km away,  $82.5^\circ$

### Exercise 7.2

1. Parallelogram 2.  $\sqrt{13}$  units,  $\sqrt{17}$  units 4.  $\frac{11}{5}\hat{i} - \frac{2}{5}\hat{j}$  5. Y(5, 5)

6.  $\left(\frac{11}{3}, \frac{7}{3}\right)$  7. 1 8. D(-4, 7) 9. B(1, 3)

### Miscellaneous Exercise-7

1. (i) b (ii) a (iii) c (iv) d (v) c (vi) d (vii) a (viii) b (ix) d  
 (x) c (xi) a (xii) b (xiii) a (xiv) c

2.  $\vec{p} \neq \vec{q}$  3.  $x = 2$

4. (i)  $\frac{1}{2}\vec{a}$  (ii)  $-\frac{1}{2}\vec{a}$  (iii)  $\vec{b} - \frac{1}{2}\vec{a}$  (iv)  $\vec{a}$

5. Resultant force = 533.82 N, direction =  $44.05^\circ$  with horizontal

6. (i) First ball traveled 25.73 m farther than the second ball.

(ii) Distance between the two balls = 43.68 m

7. (i) 7.5 m/s west (ii) 4.5 m/s east (iii) 6.18 m/s at an angle of  $14.04^\circ$  north of east

8. (i) 158.11 km/h (ii) 1,581.1 km

**Exercise 8.1**

1. (i)  $55^\circ$  (ii)  $42^\circ$  (iii)  $69^\circ$  (iv)  $38^\circ$   
 2.  $144^\circ$  3.  $93^\circ$  4.  $\sin 120^\circ = \sin 60^\circ = \frac{\sqrt{3}}{2}$ ,  $\cos 120^\circ = -\cos 60^\circ = -\frac{1}{2}$ ,  $\tan 120^\circ = -\sqrt{3}$   
 5.  $\sin \frac{3\pi}{4} = \sin \frac{\pi}{4} = \frac{1}{\sqrt{2}}$ ,  $\cos \frac{3\pi}{4} = -\cos \frac{\pi}{4} = -\frac{\sqrt{2}}{2}$ ,  $\tan \frac{3\pi}{4} = -1$  6. (i) -0.559  
 (ii) 0.83 (iii) -1.485 (iv) -0.674 (v) -1.789 (vi) 1.204

**Exercise 8.2**

1. (i)  $\beta = 50^\circ$ ,  $a = 9.64m$ ,  $b = 11.49m$  (ii)  $\alpha = 54.5^\circ$ ,  $c = 86.1m$ ,  $a = 70.1m$   
 (iii)  $c = 19.5cm$ ,  $\alpha = 50.20^\circ$ ,  $\beta = 39.8^\circ$   
 2. (i)  $\alpha = 55^\circ$ ,  $H = 14.64cm$ ,  $P = 8.4cm$  (ii)  $\beta = 64.41^\circ$ ,  $a = 14.54cm$ ,  $c = 33.5cm$   
 (iii)  $c \cong 56cm$ ,  $\alpha \cong 63.43^\circ$ ,  $\beta \cong 26.57^\circ$  (vi)  $b \cong 26.46cm$ ,  $\alpha \cong 41.84^\circ$ ,  $\beta \cong 48.16^\circ$   
 (v)  $c \cong 40.5cm$ ,  $a \cong 32.6cm$ ,  $\beta \cong 53.75^\circ$  (vi)  $c \cong 38.7cm$ ,  $b \cong 37.32cm$ ,  $\beta \cong 14.12^\circ$   
 3.  $42.89m$  4.  $116.5m$  5.  $5.18\text{ ft}$  6.  $26.59$  7.  $48.6^\circ$  8.  $24\text{ ft}$  9.  $61^\circ$  10.  $32.47m$   
 11.  $814.3m$  12.  $4.7m$  13.  $23.2\text{ m}$  14.  $6.66\text{ m}$  15.  $9.33m$

**Exercise 8.3**

1. (i)  $\alpha = 36.1^\circ$ ,  $\beta = 85.4^\circ$ ,  $c = 10.2$  (ii)  $\gamma = 48.3^\circ$ ,  $\beta = 95.7^\circ$ ,  $c = 11.8$   
 (iii)  $\alpha = 86^\circ$ ,  $\gamma = 79^\circ$ ,  $b = 38.9$  (iv)  $\alpha = 71.57^\circ$ ,  $\gamma = 63.43^\circ$ ,  $b = 30$   
 (v)  $\alpha = 37.7^\circ$ ,  $\beta = 58.3^\circ$ ,  $\gamma = 88.0^\circ$  (vi)  $\alpha = 33.33^\circ$ ,  $\beta = 33.33^\circ$ ,  $\gamma = 112.89^\circ$   
 (vii)  $\alpha = 36.3^\circ$ ,  $\beta = 62.7^\circ$ ,  $\gamma = 81.0^\circ$  (viii)  $\alpha = 60^\circ$ ,  $\beta = 60^\circ$ ,  $\gamma = 60^\circ$   
 (ix)  $\alpha = 22.62^\circ$ ,  $\beta = 67.38^\circ$ ,  $\gamma = 90^\circ$   
 3. (i)  $\gamma = 80^\circ$ ,  $b \cong 13.5$ ,  $c \cong 15.3$  (ii)  $\gamma = 60^\circ$ ,  $a \cong 16.3$ ,  $c \cong 18.4$   
 (iii)  $\gamma = 60^\circ$ ,  $a \cong 9.8$ ,  $c \cong 13.4$  (iv)  $\alpha \cong 19.42^\circ$ ,  $a \cong 9.1$ ,  $c \cong 23.8$   
 (v)  $\gamma \cong 69.33^\circ$ ,  $b \cong 34.2$ ,  $c \cong 32.1$  (vi)  $\beta \cong 52.28^\circ$ ,  $a \cong 35.2$ ,  $b \cong 41.2$   
 5. (i)  $\alpha = 48.6^\circ$ ,  $\gamma \cong 51.4^\circ$ ,  $c \cong 10.4$  (ii)  $\alpha = 71.8^\circ$ ,  $\gamma \cong 39.2^\circ$ ,  $a \cong 20.1$   
 (iii)  $\alpha = 90^\circ$ ,  $\beta = 26^\circ$ ,  $a \cong 13.8$  (iv)  $\gamma = 101.3^\circ$ ,  $\beta = 23.6^\circ$ ,  $c = 43$   
 (v)  $\alpha \cong 57.4^\circ$ ,  $\beta \cong 35.1^\circ$ ,  $\gamma \cong 87.5^\circ$  (vi)  $\alpha \cong 83.6^\circ$ ,  $b \cong 36.2$ ,  $c \cong 88$   
 6. Distance =  $362.75\text{ km}$  7. Perimeter =  $1425\text{ m}$ , Area  $\cong 84852.8\text{ m}^2$   
 8. Largest Angle  $\cong 83.41^\circ$ , Smallest Angle  $\cong 44.64^\circ$  9.  $70.75\text{ cm}$ ,  $98.97\text{ cm}$   
 10. (i) BC =  $10.15\text{ inch}$  (ii)  $\angle BDC = 19.49^\circ$  11. (i) BC =  $11.49\text{ m}$  (ii)  $\angle EDG = 56.9^\circ$   
 12.  $284.51\text{ ft}$  13.  $9.51\text{ miles}$  14.  $25.98\text{ cm}$  15.  $36.71^\circ$

**Exercise 8.4**

1. (i) 52.27 square units (ii) 269.74 square units (iii) 63.3 square units (iv) 814.8 square units  
 (v) 5.13 square units (vi) Not possible (not triangle) (vii) 87.56 square units (viii) 254.73 square units  
 (ix) 136.04 square units (x) 154.55 square units (xi) 61.5 square units (xii) 295.1 square units  
 (xiii) 168 square units (xiv) 252 square units (xv) 0.033 square units (xvi) 74.53 square units  
 (xvii) 1.69 square units  
 2. 127.28 square units 3. 387.4 square units  
 4. (a)  $\cos \theta = \frac{1}{3}$  (b)  $\sin \theta = \frac{2\sqrt{2}}{3}$  (c) 5.66 sq. units (d) 6 sq. units  
 5.  $A \cong 233.75\text{ sq units}$  6.  $125\sqrt{3}\text{ sq units}$  7. Do yourself.  
 8. Length of side =  $22.3\text{ m}$ , Third Angle =  $75^\circ$  9.  $27\sqrt{3}\text{ cm}^2$   
 10. (a) Sine of angle B =  $\frac{2}{3}$  (b)  $\angle B = 41.8^\circ$  11. (a) Area =  $(ac) \sin \theta$  (b)  $\theta = 90^\circ$

**Exercise 8.5**

1. (i)  $r \cong 3.24$ ,  $R \cong 8.53$ ,  $r_1 \cong 6.48$ ,  $r_2 \cong 9.26$ ,  $r_3 \cong 21.6$   
 (ii)  $r \cong 6.87$ ,  $R \cong 15.18$ ,  $r_1 \cong 16.31$ ,  $r_2 \cong 18.64$ ,  $r_3 \cong 32.61$   
 (iii)  $r = 1$ ,  $R = 2.5$ ,  $r_1 = 2$ ,  $r_2 = 3$ ,  $r_3 = 3$   
 (iv)  $r \cong 16.33$ ,  $R \cong 35.75$ ,  $r_1 \cong 36.74$ ,  $r_2 \cong 48.99$ ,  $r_3 \cong 73.48$

10.  $r \cong 2.43 \text{ cm}$ ,  $R \cong 7.64 \text{ cm}$

11. (i)  $2\pi r_1 \cong 20.16 \text{ cm}$  (ii)  $2\pi r_3 \cong 70.56 \text{ cm}$  12.  $A \cong 2.72 \text{ cm}^2$ ,  $C \cong 5.85 \text{ cm}$

### Exercise 8.6

1. 16 cm 2.  $78.86^\circ$  3.  $6\sqrt{2} \text{ cm}$ ,  $6\sqrt{2} \text{ cm}$  4.  $13.9^\circ$  5. (i) 20cm (ii) 25 cm (iii)  $26.9^\circ$

6.  $46.7^\circ$  7. (i) 10.82m (ii) 12.44m (iii)  $53.07^\circ$  (iv)  $64.52^\circ$

8. (i) 29 km (ii) 29 km (iii) 56 km

### Miscellaneous Exercise-8

1. (i) a (ii) c (iii) b (iv) d (v) c (vi) d  
(vii) b (viii) a (ix) c (x) d (xi) b (xii) a

2. (i)  $\gamma = 145^\circ$ ,  $b = 10.6$ ,  $c = 17.7$  (ii)  $\beta = 32.4^\circ$ ,  $\gamma = 32.6^\circ$ ,  $c = 63.5$

(iii)  $\gamma = 56.9^\circ$ ,  $\alpha = 24.9^\circ$ ,  $a = 7.37$  (iv)  $b \approx 114.74$ ,  $a \approx 146.99$ ,  $\gamma = 20^\circ$

(v)  $b \cong 66.86$ ,  $\alpha = 28.88^\circ$ ,  $\gamma = 43.12^\circ$  (vi)  $\alpha = 53.13^\circ$ ,  $\beta = 36.87^\circ$ ,  $\gamma = 90^\circ$

3. 14.48 cm 4. a. 1.8 km b. 4.25 km 5. a. 33.73 inches b.  $57.5^\circ$  6.  $1717.62 \text{ m}^2$

7. a. 8.84 m b. 32.18 m 8. 137.3 meters and 121.6 meters

9. Yes, it is possible for Aamir to draw a parallelogram with one side of 12 cm, one diagonal of 10 cm, and one angle of  $120^\circ$ . The calculated length of the other diagonal is approximately 20.8 cm, which is consistent with the geometry of a parallelogram.

10. Approximately 73.8feet (higher observer) and 45.3 feet (lower observer). The distance from the pipe to the apartment complex (horizontal distance) is approximately 42.3 feet. 11. 5.22 miles 12. Both are 21.9 cm

16. 1702.3 meters 17.  $24.96^\circ$  18.  $21^\circ$


## UNIT 9

### Exercise 9.1

2. (i)  $x = 4\text{cm}$  (ii)  $y = 6\text{cm}$  3.  $CD = 26\text{cm}$ ,  $CE = 1\text{cm}$  4.  $CD = 24\text{cm}$

5.  $AB = 24\text{cm}$


6.



As the circle has only one centre O, therefore one and only one circle can pass through vertices of rectangle.

As the circle has only one centre B, therefore one and only one circle can pass through points A, C and D.


7.



8. (i)  $OU = 16 - r$  (ii)  $r^2 = 5^2 + (16 - r)^2$ ;  $r = 8.8\text{cm}$

9.  $r = 10\text{cm}$

10. Sides of triangle OCM satisfy Pythagorean triplet (3, 4, 5). Therefore,  $\angle OMC = 90^\circ$  proving the result.



11. Prove  $\triangle ACD \cong \triangle BCD$ , with the help of SAS postulate, then  $AC = BC$ .

### Exercise 9.2

1.  $\angle OCF = 15^\circ$ ,  $\angle CFD = 75^\circ$

2.  $2.24\text{cm}$ , Yes, they are equal.

3.  $7\text{cm}$

4. (i)  $a = 9\text{cm}$ ,  $y = 53^\circ$

(ii)  $a = 6\text{cm}$ ,  $x = 45^\circ$ ,  $y = 8.49\text{cm}$

5.  $16.53\text{cm}$

6.  $6\text{cm}, 4\text{cm}$

7.  $31\text{cm}$

8.  $\sqrt{10}\text{cm} = 3.16\text{cm}$

9.  $5.15\text{cm}$

**Exercise 9.3**

1. (i) yes (ii) yes (iii) chord AB < chord AD  
 4. (i) no (ii) both are proportional 5. (i) yes (ii) CD = 4 cm (iii) 2.24 cm  
 6.  $120^\circ$  8. 6 cm 11. (a)  $130^\circ$  (b)  $85^\circ$  (c)  $75^\circ$  (d)  $70^\circ$   
 13.  $\angle AOB = \angle COD = 100^\circ$ ; Equal chords of a circle subtend equal angles at the centre.  
 14. (i) All the four chords make central angle of  $90^\circ$ . (ii)  $\angle PCB = 45^\circ$ ,  $\angle ACB = 90^\circ$   
 (iii)  $\angle CBD$ ,  $\angle BDA$ ,  $\angle DAC$  (iv) square (v) one and only one  
 15. (i)  $\angle PTQ = 45^\circ$ ,  $\angle RTS = 135^\circ$  (ii) 1 : 3

**Exercise 9.4**

1. (i) Length of minor arc = 12.21 cm, length of major arc = 31.75 cm (ii) 43.96 cm; yes  
 2. (i) Area of minor sector = 33.51  $\text{cm}^2$ , area of major sector = 167.55  $\text{cm}^2$   
 (ii) 201.06  $\text{cm}^2$ ; yes 3. 26.2 cm  
 4. (i) 22 cm (ii) 10.5 cm (iii) 11.5 cm (iv)  $231\text{cm}^2$  (v)  $52.4\text{cm}^2$  (vi)  $178.6\text{cm}^2$   
 5. (i) 19.8 cm (ii) 22 cm (iii) 41.8 cm (iv)  $98\text{cm}^2$  (v)  $154\text{cm}^2$  (vi)  $56\text{cm}^2$   
 6. 12.57 cm 7. 4 cm 8. A =  $12.32 \text{ ft}^2$ ; P = 14.4 ft 9. 35.9 m 10. 28 m  
 11. P = 12.48 ft; A =  $1.85 \text{ ft}^2$  12. 29.3 ft 13. (i)  $12228.1 \text{ m}^2$  (ii) 65.1 m

**Miscellaneous Exercise-9**

1. (i) b (ii) d (iii) c (iv) c (v) b (vi) a (vii) d (viii) a (ix) d  
 (x) c (xi) b (xii) c (xiii) b (xiv) d (xv) b (xvi) a  
 (xvii) b (xviii) d

**UNIT 10****Exercise 10.1**

1. (i)  $r = 5\text{cm}$  (ii)  $d = 4\sqrt{7}\text{cm}$  (iii)  $x = 8\text{cm}, y = 4.62\text{cm}$   
 (iv)  $x = 25^\circ, y = 65^\circ$  (v)  $a = 30^\circ, b = 10\sqrt{3}\text{cm}$  (vi)  $a = b = 60^\circ$   
 (vii)  $a = 15^\circ$  (viii)  $x = 63^\circ, y = 60^\circ, z = 57^\circ$   
 2. (i)  $\angle OBA = 90^\circ$  (ii)  $\angle OCB = 30^\circ$  (iii)  $\angle BAC = 60^\circ$  (iv)  $\angle ABC = 60^\circ$   
 4. 4 ft 5. 2cm, 6cm, 10cm 6. 5.06 cm 7.  $\angle PGH = 35^\circ$  8.  $x = 48^\circ, y = 68^\circ$

**Exercise 10.2**

1.  $\angle Q = 30^\circ, \angle ROS = 60^\circ$  2.  $\angle ABO = 30^\circ, \angle AOB = 120^\circ, \angle ACB = 60^\circ$   
 3.  $\angle ABC = 90^\circ, \angle BCA = 50^\circ, \angle CAD = 30^\circ$  4.  $\angle O = 60^\circ, \angle C = 30^\circ$   
 5.  $\angle AEB = 110^\circ, \angle B = 40^\circ, \angle C = 40^\circ, \angle D = 30^\circ$   
 6. (i) 4 cm (ii)  $\angle ABE = 60^\circ$  (iii)  $\angle ABC = 90^\circ$  (iv)  $\angle EBD = 30^\circ$   
 7. (i) 2 (ii) 4 cm (iii)  $\angle BQC = \angle BPO = 90^\circ$  (iv)  $5\sqrt{5}\text{ cm}$  (v) 15 cm  
 8. (i)  $\angle QSR = 120^\circ$  (ii)  $\angle PRS = 30^\circ$  (iii)  $\angle PQS = 30^\circ$  (iv)  $\angle QTR = 60^\circ$   
 (v) Angle in a segment greater than semicircle is acute angle.  
 9. (i)  $\angle BAD = 65^\circ$  (ii)  $\angle BCD = 115^\circ$  (iii)  $\angle ABC + \angle ADC = 180^\circ$   
 10. (i)  $\angle QRS = 85^\circ$  (ii)  $\angle PQR = 95^\circ$  (iii)  $\angle PSR = 85^\circ$   
 11. (i)  $\angle BAC = 30^\circ$  (ii)  $\angle CAD = 30^\circ$   
 12. (i)  $\angle ECD = 65^\circ$  (ii)  $\angle CDE = 95^\circ$  (iii)  $\angle CED = 20^\circ$   
 13. (i) 2.2 cm (ii) 1.2 cm 14. 40 cm, 125.6 cm

**Miscellaneous Exercise-10**

1. (i) a (ii) c (iii) b (iv) b (v) d (vi) a (vii) d (viii) a (ix) b  
 (x) c (xi) d (xii) c (xiii) d (xiv) b (xv) d (xvi) a  
 (xvii) c (xviii) c (xix) b (xx) d  
 2.  $6\sqrt{2}\text{ cm}$  3. 8 cm 4. 4.75 cm

## UNIT 11

### Exercise 11.1

8. (i) He will take three points on circular path and join them to obtain a triangle. Point of intersection of right bisectors of sides of the triangle is the required point.  
 (ii) 25 cm      9. 22 tiles

### Exercise 11.2

11. 122 inches      12. 12451.51 km      13. 61.68 cm      14. 5 cm

### Miscellaneous Exercise-11

3. (i) d    (ii) a    (iii) b    (iv) c    (v) a    (vi) c    (vii) d    (viii) b    (ix) a  
 (x) d    (xi) c    (xii) d    (xiii) b    (xiv)

## UNIT 12

### Exercise 12.1

1.

Class Interval	1 – 10	11 – 20	21 – 30	31 – 40	41 – 50	51 – 60
Frequency	3	4	7	9	5	2
Cumulative freq	3	7	14	23	28	30



i. 30    ii. 21-30    iii. 7    iv. 31-40    v. 50.5

2.

3.    33

a. 26.5 app  
c. 34.5app

b. 37app





From the graph, mass of the baby at the age of  $4\frac{1}{2}$  months is 12kg app.

4. Draw ogive yourself. a. 90    b. 72.5    c. 45    d. 9    e. 27    f. 1    5. do yourself  
 6. a. 75 app    b. 45% app    c. 3 : 1 app    d. 25    e. 55m app    f. 80m app  
 7. Range = Rs.125,000 & IQR = Rs. 40,000    8. She performed above 75% of her class  
 9. Between Rs. 25,000,000 & Rs. 30,000,000    10. i. 38app    ii. 30app    iii. 50app    iv. 20app    v. 48app    vi. 32app


### Exercise 12.2

1.



- a.  $IQR_1 = 500$ ,  $IQR_2 = 8$ , b. negative  
 2. a. Rs. 500    b. 0 & 700    c. Rs. 400    c. lighter vehicles are more fuel efficient  
 d. Rs. 300    e. positive

3.



Positive correlation indicates increase in number of flowers with increased rain quantity.

4. a. Positive correlation indicates higher score with more practice time.

b. draw yourself

### Exercise 12.3

1. Measure of spread of data ; Range , Variance , Standard deviation

2. range = 599, Variance = 28886 app 3i. 758.8 ii. 19097 iii. 138.19 iv. 18.2

4. range = Rs.60,000 , Variance = 246437000app , SD = 15698.3 CV = 22861.5

5. mean = 66.35 , Variance = 67.57 , SD = 8.22

6. range = 2,000 , Variance = 243806app, SD = 493.8

7. OT: range = 60, mean = 127.3, Variance = 204.7app , SD = 14.31 CV = 11.24

OPD: range = 60, mean = 127, Variance = 292.4app , SD = 17.1 CV = 13.46

OPD patients BP data have more variation.

8. Mean : 103, 66.9 Variance: 33.3, 147.6

Next week average temp seems to be 103F app in Jhelum and 147.6F app in Indiana.

### Exercise 12.4

1.

trials	First trial			
	x	1	2	3
Second trial	1	1	2	3
	2	2	4	1
	3	3	6	9

P(Even) = 5/9

3. 0.85 4. 0.7 5. 0.9


6. 5/6 7. 25/27

8. 0.28

9. 0.63

10. 0.72

2.



### Miscellaneous Exercise-12

1. i. d ii. b iii. b iv. c v. d vi. d vii. b viii. a ix. d x. c  
xi. c xii. d xiii. a xiv. a xv. c

2. Draw the graph yourself , variance = 83.14 , SD = 9.12

3. a.  $IQR_1 = 30$ ,  $IQR_2 = 10/10$  b. negative c. more exercise time caused reduction in BP.

4. 25%, IQR = 20 5. Hazel: M = 65 app, IQR = 84-48 = 36,

- Mauve: M = 68 app, IQR = 86-45 = 41

6. 3/10 7. 0.18

**Absolute value of a complex number:** The absolute value of  $z$ , denoted by  $|z|$  is defined as  $|z| = \sqrt{a^2 + b^2}$  and is a real number.

**Absolute valued function:** The function defined by  $f(x) = |x|$ , is called the absolute valued function.

**Additive identity in matrices:** Additive identity is such a matrix which causes no change in any matrix  $A$  while 'adding to' or 'subtracting from' it.

**Additive inverse of a matrix:** Additive inverse of a matrix ' $A$ ' is such a matrix which when added to  $A$ , gives additive identity matrix of the same order.

**Additive inverse of complex number:** Complex number that is added to a given complex number in order to make the sum zero is known as the additive inverse.

**Algebraic expressions:** A statement in which variables or constants or both are connected by arithmetic operations.

**Imaginary Number:** An imaginary number is a number that can be written  $a + bi$ , where  $a$  and  $b$  are real numbers,  $b \neq 0$

**Angle in a segment:** An angle whose vertex is a point on the arc of a circle and whose arms pass through end points of chord.

**Arc of a circle:** An arc is a part of a circle. An arc  $AB$  is denoted by  $\widehat{AB}$ .

**Associative property of matrix addition:** If  $A$ ,  $B$  and  $C$  are three matrices of same order, then  $(A + B) + C = A + (B + C)$ , is called associative property of matrix addition.

**Associative property of matrix multiplication:** If  $A$ ,  $B$ , and  $C$  are matrices and products  $AB$  and  $BC$  are possible, then  $(AB)C = A(BC)$

**Bijective function:** A function which is both onto and 1-1 is called a bijective function.

**Binomial:** A polynomial having two terms is called a binomial.

**Box-and-whisker plot:** It is a graphical representation of a data set that displays the distribution of values through 5 data points i.e. min, max,  $Q_1$ ,  $Q_2$ ,  $Q_3$ .

**Central angle:** An angle whose vertex is the centre of circle and whose arms pass through end points of arc is known as central angle of arc.

**Chord of a circle:** A line segment joining two points of a circle is called chord of that circle.

**Circumcenter:** It is the point of intersection of right bisectors of sides of a triangle.

**Circumcircle:** A circle passing through the vertices of any triangle (polygon) is called circumcircle.

**Coefficient of variation:** The percentage ratio between the standard deviation and mean of a data.

**Column matrix:** A matrix having only one column is called a column matrix.

**Column:** Vertical arrangement of elements is in a matrix called a column.

**Cumulative frequency:** In a group data, number of values less than or within the limits of a class is called cumulative frequency of the respective class.

**Commutative property of matrix addition:** If  $A$  and  $B$  are two matrices of same order, then  $A + B = B + A$ , is called commutative property of matrix addition.

**Complex number:** A complex number is any number that can be written in the form of  $a + bi$ , where  $a$  and  $b$  are any real numbers and both  $a$  and  $b$  can be zero.

**Composition of functions:** The composition of  $f(x)$  and  $g(x)$  is a function denoted by  $f \circ g$  and is written as:

$(f \circ g)(x) = f(g(x))$

**Conditional probability:** The probability of an event occurring given that another event has occurred.

**Congruent arcs:** Two arcs of a circle are congruent if their central angles are congruent.

**Conjugates of complex numbers:** Conjugate of complex number is a complex number obtained by changing the sign of the imaginary part.

**Constant function:** A polynomial function of degree zero is called a constant function.

**Constant polynomial:** A polynomial having degree zero is called a constant polynomial.

**Coterminal angles:** Two angles having same terminal ray in standard position, are called coterminal angles.

**Cubic function:** A polynomial function of degree three is called a cubic function.

**Cubic polynomial:** A polynomial having degree three is called a cubic polynomial.

**Cumulative frequency polygon:** A polygon made by joining the points whose x-coordinates are class marks and y-coordinates are their cumulative frequencies is called a cumulative frequency polygon.

**Cyclic quadrilateral:** A quadrilateral inscribed in a circle is called cyclic quadrilateral.

**Deciles:** Deciles divide a data set into 10 equal parts, each representing 10% of the data.

**Dependent probability:** The probability of two or more events occurring dependently on each other.

**Determinant:** The determinant of its matrix  $\begin{bmatrix} a & c \\ b & d \end{bmatrix}$  is denoted by  $\begin{vmatrix} a & c \\ b & d \end{vmatrix} = ad - bc$ .

**Diagonal matrix:** A square matrix in which every element except the primary (principal) diagonal elements is zero is called a diagonal matrix.

**Domain of a function:** A set containing first element of each ordered pair in the function is called its domain.

**E-center:** It is the point of intersection of one internal bisector of angle and external bisectors of remaining angles of a triangle.

**Equal matrices:** Two matrices are said to be equal if and only if their order is same and corresponding elements are same.

**Equal vectors:** Two vectors  $\vec{u}$  and  $\vec{v}$  are said to be equal if both have the same magnitude and direction.

**Equality of complex numbers:** If two complex numbers  $a + bi$  and  $c + di$  are equal then  $a = c$  and  $b = d$ .

**Escribed circle:** A circle touching one of the sides of a triangle externally and the extensions of its two other sides internally, is called an escribed circle.

**Evaluation of a function:** It is a process of finding the value of dependent variable by substituting any specific value of the independent variable.

**Evaluation:** The process of getting a numerical answer by substituting numbers for each variable is called evaluation.

**Experimental probability:** Probability based on the results of repeated trials of an experiment.

**Exponential function:** The exponential function is of the form  $f(x) = a^x$ , where  $a > 0$  and  $a \neq 1$ .

**Frequency polygon:** The representation of data by a polygon in which class marks are plotted along x-axis and frequencies along y-axis is called a frequency polygon.

**Frequency table:** A table consisting of values of a data along with their frequencies.

**Function:** A function  $f$  from set A to set B is a relation, rule or mapping which maps each element of set A to a unique element of set B.

**Graph of inequality:** On a number line, the graph of an inequality in one variable is the set of points that represents all solutions of the inequality.

**Head to tail rule:** The method of adding vectors in which tail of the second vector is joined with the head of the first vector.

**Imaginary number:** An imaginary number is the product of a real number and the imaginary unit  $i$ , which is defined by its property  $i^2 = -1$ .

**Incentre:** It is the point of intersection of angle bisectors of a triangle.

**Incircle:** A circle drawn inside and touching the sides of a triangle (polygon) is called incircle.

**Injective function:** A function which is both into and 1-1 is called an injective function.

**Inter quartile range:** The IQR represents the range of values within which the middle 50% of the data points fall.

**Into function:** A function  $f : A \rightarrow B$  will be into if there is at least one element in set B which is not an image of any element of set A.

**Inverse of a function:** The inverse of any function  $f(x)$  is a function denoted by  $f^{-1}(x)$  which reverses the effect of  $f(x)$  and it undoes what  $f(x)$  does.

**Irrational expression:** An algebraic expression which cannot be expressed in the form  $\frac{P(x)}{Q(x)}$ , where  $P(x)$  and  $Q(x)$  are polynomials and  $Q(x) \neq 0$ .

**Joint probability:** The probability of two or more events occurring together.

**Linear function:** A polynomial function of degree one is called a linear function.

**Linear polynomial:** A polynomial having degree one is called a linear polynomial.

**Matrix:** A matrix is a rectangular arrangement of numbers.

**Measure of central tendency:** A single value which can represent the whole data.

**Measures of dispersion:** The measures which tell us how the values of a data are scattered around their average, are called measures of dispersion.

**Minor and major arcs:** Minor arc is the arc included in a semicircle while the arc which includes a semicircle is called major arc.

**Minor and major segments:** The region between chord and minor arc is called minor segment and the region between chord and major arc is called major segment.

**Monomial:** A polynomial having one term is called a monomial. e.g., 3,  $x$ , 0 etc.

**Multiplicative identity in matrices:** Multiplicative identity is such a matrix which causes no change in any matrix A when multiplied with A.

**Mutually exclusive probability:** The probability of two or more events that cannot occur simultaneously.

**Negative correlation:** If the boxes and whiskers of both variables tend to move in opposite directions it may indicate a negative correlation.

**Negative of a vector:** A vector having the same magnitude but opposite in direction of a given vector is called the negative of that vector.

**No correlation:** If the boxes and whiskers of both variables appear unrelated or don't show a clear pattern, it may indicate no correlation.

**Null matrix:** A matrix of any order, having all the elements equal to zero, is called a null matrix.

**Oblique triangle:** An oblique triangle is any triangle that is not a right triangle.

**One-to-one function:** A function f is one-to-one if distinct elements of set A have distinct images in set B.

**Onto function:** A function f : A → B will be onto if every element of set B is an image of at least one element of set A.

**Operations on functions:** Operations on functions are the ways of combining functions to create new functions.

**Order of matrix:** If 'A' is a matrix with 'm' number of rows and 'n' number of columns, then order of the matrix is m-by-n.

**Parallel vectors:** Two non-zero vectors  $\vec{u}$  and  $\vec{v}$  are said to be parallel if  $\vec{u} = \lambda\vec{v}$ .

**Percentiles:** Percentiles divide a data set into 100 equal parts, each representing 1% of the data.

**Polynomial function:** A function  $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots + a_{n-1}x^{n-1} + a_nx^n$ , where  $a_0, a_1, a_2, a_3, \dots, a_{n-1}, a_n$  are all real numbers and n is a non-negative integer.

**Polynomials:** Algebraic expressions in which exponents of the variables involved are non-negative integers.

**Position vector:** The vector used to specify the position of a point P with respect to origin O is called position vector of P.

**Positive correlation:** If the boxes and whiskers of both variables tend to move in the same direction it may indicate a positive correlation.

**Probability:** Probability is a measure of the likelihood of an event occurring.

**Product of matrices:** Two matrices A and B are conformable for product AB, if:

**Quadrantal angles:** Angles in standard position whose terminal side lies on an axis.

**Quadratic equation:** It is an equation that can be written in the form of  $ax^2 + bx + c = 0$ , where  $a \neq 0$ .

**Quadratic function:** A polynomial function of degree two is called a quadratic function.

**Quadratic inequality:**  $ax^2 + bx + c > 0$  or  $ax^2 + bx + c < 0$  is called quadratic inequality.

**Quadratic polynomial:** A polynomial having degree two is called a quadratic polynomial.

**Quartiles:** The three values which divide the whole data into four equal parts are called quartiles.

**Range of a function:** A set containing second element of each ordered pair in the function is called its range.

**Range:** Range is the difference of largest and smallest values of the data.

**Rational equations:** An equation that contains one or more rational expressions is called a rational equation.

**Rational expression:** Algebraic expression of the form  $\frac{P(x)}{Q(x)}$ , where P(x) and Q(x) are polynomials and  $Q(x) \neq 0$ .

**Reciprocal function:** The function defined by  $f(x) = \frac{1}{x}$ , where  $x \neq 0$  is called a reciprocal function.

**Rectangular matrix:** A matrix with unequal number of rows and columns is called a rectangular matrix.

**Row matrix:** A matrix having only one row is called a row matrix.

**Row:** Horizontal arrangement of elements in a matrix is called a row.

**Scalar matrix:** A diagonal matrix in which all the elements of primary diagonal are equal and non-zero is called a scalar matrix.

**Scalar multiplication of a matrix:** Scalar multiplication of a matrix means multiplication of a matrix with a constant.

**Scalar multiplication:** Multiplication of a vector by a scalar.

**Scalar:** A physical quantity that possesses magnitude only, is called a scalar quantity.

**Segment of circle:** A chord of a circle divides the circular region into two parts, called segment of the circle.

**Semicircle:** A diameter divides a circle into two equal parts. Each part is called semicircle.

**Singular and non-singular matrices:** A is singular matrix, if  $|A| = 0$  otherwise non-singular.

**Skew symmetric matrix:** A is skew Symmetric if,  $A = -A^t$ .

**Solution of a system of linear inequalities:** It is an ordered pair that is a solution of each inequality in the system.

**Solution of inequality:** A solution of an inequality in two variables  $x$  and  $y$  is an ordered pair  $(x, y)$  that produces a true statement when the values of  $x$  and  $y$  are satisfied into the inequality.

**Solution of the system:** The set of all the ordered pairs  $(x, y)$  which satisfies the system of equations is called the solution of the system.

**Square matrix:** A matrix with equal number of rows and columns is called a square matrix.

**Square root function:** The function defined by  $f(x) = \sqrt{x}$ , where  $x \geq 0$  is called a square root function.

**Standard deviation:** It is the positive square root of variance.

**Substitution:** The process of replacing the variables by numbers in an algebraic expression is called 'substitution'.

**Subtraction of two vectors:** To subtract a vector, form the other we find the negative vector of the vector to be subtracted and then add it to the other vector.

**Symmetric matrix:** A is symmetric if,  $A = A^t$ .

**System of linear inequalities:** A system of linear inequalities in two variables or simply a system of inequalities consists of two or more linear inequalities in the same variable

**System of simultaneous equations:** A system of equations having a common solution is called a system of simultaneous equations.

**Tangent line:** A line which touches the circle at only one point is called tangent line.

**Theoretical probability:** The number of favorable outcomes divided by the total number of possible outcomes.

**Three-D Trigonometry:** It is an application of the trigonometric skills developed for 3-dimensional shapes in order to find unknown sides and angles.

**Transpose of a matrix:** If rows (columns) of a matrix  $P$  are changed into columns (rows) then the resulting matrix is called transpose of the matrix  $P$ .

**Trigonometric ratio:** The ratio between any two sides of a right-angled triangle is called trigonometric ratio.

**Trinomial:** A polynomial having three terms is called a trinomial.

**Unit matrix:** A scalar matrix in which all the primary diagonal elements are equal to 1, is called a unit matrix.

**Unit vector:** A vector which is in the direction of non-zero vector  $\vec{u}$  and has magnitude 1 is called a unit vector of  $\vec{u}$ .

**Variance:** Variance is the ratio between sum of squares of deviations from mean and the number of values.

**Vector:** A physical quantity that possesses both magnitude and direction, is called a vector quantity.

**Zero or null vector:** If the initial and terminal points of a vector coincide then the vector has zero length and is called zero vector.

**Zero polynomial function:** A polynomial function of no degree is called a zero polynomial function.

**Zero polynomial:** '0' is called a polynomial of no degree.

**Zero product property:** For any real number  $a$  and  $b$ , if  $ab = 0$ , then either  $a = 0$  or  $b = 0$ .

# I N D E X

Imaginary number	07	Translation by vector	145
Complex number	08	Additions of vectors	147
Operations on complex numbers	09	Subtraction of vectors	148
Properties of complex numbers	13	Rectangular components of a vector	150
Graph of complex numbers	15	Applications of vectors in geometry	155
Modulus of complex numbers	16	Trigonometric ratios	161
Application of complex numbers	21	Solution of right triangles	163
Solution of complex equations	23	Law of cosines	167
Solution of quadratic equations	29	Law of sines	169
Quadratic formula	32	Half angles formulas	174
Equation reducible to quadratic form	34	Area of triangular regions	178
Nature of roots	41	Circles connected with triangles	182
Roots and coefficients	44	3-D trigonometry	187
Formation of quadratic equation	46	Chords of a circle	197
Simultaneous equations	48	Arcs of a circle	203
Types of matrices	57	Applications of chords and arcs	210
Operations on matrices	60	Tangents to a circle	216
Multiplicative inverse of matrix	67	Angle in segment of a circle	222
Application of matrices	74	Cyclic quadrilateral	224
Quadratic inequalities	81	Constructions of circles	233
Linear inequalities	84	Constructions of tangents to circles	235
Absolute valued equations	88	Cumulative frequency distribution	246
Absolute valued inequalities	90	Cumulative frequency polygon	247
Evaluation of algebraic expressions	101	Construction of frequency polygon	248
Operations on algebraic fractions	103	Measures of distribution	249
Solving rational equations	109	Bos & Visker correlation plot	255
Types of functions	117	Construction of scatter diagram	257
Evaluating a function	119	Measures of dispersion	262
Operations on functions	121	Applications of variance & S. D	265
Applications of functions & graphs	135	Probability of combined events	267
Cartesian coordinate system	142	Sample space diagram	268
Vectors in a plane	143	Probability rules	271

## **A b o u t A u t h o r s**

### ***Dr. Saleem Ullah***

Dr. Saleem Ullah Satti possesses Ph.D (Maths) along with different professional teaching degrees. Presently, he is working in mathematics department Air University, Islamabad. He has experience of two decades in teaching of mathematics at various levels. He authored and reviewed many books. He has been involved in the development of curriculum for mathematics upto higher level. As a resource person, he has conducted training programmes, workshops and seminars on the subject. Email: saleemullah314@hotmail.com

### ***Dr. Khalid Mahmood***

He has a long teaching experience of more than two decades at school and college level. He holds the degree of Ph.D (Maths) along with professional degree of M.Ed. He has authored and reviewed many books in the subject of mathematics at different level. He has worked as a resource person for imparting training to teachers at elementary and secondary levels. Email: khalidmeh2012@gmail.com

### ***Ms. Nabeela Azad***

She has been teaching mathematics at the high school and college levels since 2000. She holds an M.Phil in Applied Mathematics from Quaid-e-Azam University Islamabad, as well as a professional M.Ed degree. She has worked as a subject specialist in a variety of organisations and has played a pivotal role in preparing teachers and students for national and international mathematics olympiads such as IKMC, Mismo, and Sigma 7, and many others.

### ***Ms. Hafsa Jabeen***

She has been teaching mathematics at the high school and college levels for the past 20 years. She holds a master's degree in mathematics from the University of the Punjab-Lahore. She has worked as a secondary and higher secondary subject coordinator. She has made a substantial contribution to the design and review of elementary level mathematics books under the supervision of Mr. Obaid Ur Rehman (Late), who was known for his expertise and tremendous input in writing many books on mathematics at various levels.

*studyplusplus.com (study++)*