

Numerical study of the effect of secondary electron emission on the dynamics of electron clouds in gyrotron guns

S. Guinchard¹, G. Le Bars²

¹ Ecole Polytechnique Fédérale de Lausanne (EPFL), Physics Section (SPH), CH-1015 Lausanne, Switzerland

² Ecole Polytechnique Fédérale de Lausanne (EPFL), Swiss Plasma Center (SPC), CH-1015 Lausanne, Switzerland

November 24, 2022

- Recall: electronic yield $\gamma(E) = \Lambda_{exp} * \frac{dE_{ions}}{dx}$
- Ion Induced Electrons have been modeled using a Poisson Law

- $P(x = k) = e^{-\lambda} \frac{\lambda^k}{x^k} = e^{-\gamma(E_i)} \frac{\gamma(E_i)^k}{x^k}$

- Vertical slice of He ions impinging on stainless steel electrode
 - Potential bias: $\Delta\phi = 20kV$
 - Electrode radial positions: $r_a = 10^{-3}m$,
 $r_b = 10^{-2}m$
 - Acquired energy: $E \propto \Delta\phi \log(r/r_a) / \log(r_b/r_a)$




Figure: Particles number as function of time

- Horizontal slice of He ions impinging on stainless steel electrode
 - Potential bias: $\Delta\phi = 20kV$
 - Electrode radial positions: $r_a = 10^{-3}m$, $r_b = 10^{-2}m$
 - Acquired energy: $E \propto \Delta\phi \log(r/r_a) / \log(r_b/r_a)$
 - Yield prediction: Slice 1: $\gamma = 0.96$, Slice 2: $\gamma = 1.33$
 - Obtained yield: Slice 1: 967 e^- for 1000 lost ions
 - Slice 2: 1325 for 1000 ions : Relative error $\propto 1e-3$




Figure: Particles number as function of time