

HIGHER-FORM SYMMETRIES AND TOPOLOGICAL PHASES

Sal Pace (MIT)

SP & X-G Wen, PRB 106, 045145 (2022)

SP & X-G Wen, PRB 107, 075112 (2023)

Y-T Oh, **SP**, JH Han, Y You, H-Y Lee, arXiv:2301.04706

SP & X-G Wen, arXiv:2301.05261

THE MANY-BODY SAGA

Separation of scales

UV Scale



IR Scale

Short distance
physics

Large distance, low-
energy physics

Know UV degrees of freedom
(electrons, neutrons, etc)

Emergent IR degrees of
freedom (quasiparticles,
strings, etc)

A GUIDING PRINCIPLE

UV Scale



IR Scale

Too complicated:
describes all but
hides much

Reveals universal
properties but hides
their cause

It's desirable to have a set of guiding principles to
help us understand their interplay

A GUIDING PRINCIPLE

UV Scale

Too complicated:
describes all but
hides much



IR Scale

Reveals universal
properties but hides
their cause

It's desirable to have a set of guiding principles to
help us understand their interplay

Symmetry

THE SYMMETRY RENAISSANCE

Our understanding of symmetry has been recently revolutionized through modern generalizations

THE SYMMETRY RENAISSANCE

Our understanding of symmetry has been recently revolutionized through modern generalizations

Higher-form symmetry

Non-invertible symmetry

Dipole symmetry

Loop group symmetry

Subsystem symmetry

Higher-group symmetry

Biform symmetry


Harmonic symmetry

Framed symmetry

Categorical symmetry

THE SYMMETRY RENAISSANCE

Our understanding of symmetry has been recently revolutionized through modern generalizations



WHY ARE THESE SYMMETRIES

- There's a **symmetry operator** that commutes with the Hamiltonian
- The objects carrying the symmetry charge can condense, causing **spontaneous symmetry breaking**
- Are associated with **conservation laws** that constrain dynamics
- Can have **'t Hooft anomalies** that constrain possible phases of matter

WHY ARE THESE SYMMETRIES

- There's a **symmetry operator** that commutes with the Hamiltonian
- The objects carrying the symmetry charge can condense, causing **spontaneous symmetry breaking**
- Are associated with **conservation laws** that constrain dynamics
- Can have 't Hooft **anomalies** that constrain possible phases of matter



Passes the duck test!

*If it looks like a duck, swims like a duck,
and quacks like a duck, then it probably is
a duck.*

WHY ARE THESE SYMMETRIES

- There's a **symmetry operator** that commutes with the Hamiltonian
- The objects carrying the symmetry charge can condense, causing **spontaneous symmetry breaking**
- Are associated with **conservation laws** that constrain dynamics
- Can have 't Hooft **anomalies** that constrain possible phases of matter



Passes the duck test!

*If it looks like a duck, swims like a duck,
and quacks like a duck, then it probably is
a duck.*

GENERALIZED SYMMETRIES IN CMP



G_{UV} typically includes ordinary symmetries, but $G_{\text{mid-IR}}$ and G_{IR} can include emergent generalized symmetries

- Unifying perspective on different phases of matter
- New phases of matter characterized by generalized symmetries
- Classification scheme (generalized Landau paradigm)

THE PLAN FOR THIS TALK

Explore how higher-form symmetries arise in
topological phases of quantum matter

1. How higher-form symmetries emerge & exact emergent symmetries
[**SP** & X-G Wen, arXiv:2301.05261]
2. Symmetry Protected Trivial (SPT) phase protected by higher-form symmetries
[**SP** & X-G Wen, PRB 107, 075112 (2023)]
3. The rank-2 toric code, its symmetries, and UV/IR mixing
[**SP** & X-G Wen, PRB 106, 045145 (2022)]
[Y-T Oh, **SP**, JH Han, Y You, H-Y Lee, arXiv:2301.04706]

THE PLAN FOR THIS TALK

Explore how higher-form symmetries arise in
topological phases of quantum matter

1. How higher-form symmetries emerge & exact emergent symmetries
[**SP** & X-G Wen, arXiv:2301.05261]
2. Symmetry Protected Trivial (SPT) phase protected by higher-form symmetries
[**SP** & X-G Wen, PRB 107, 075112 (2023)]
3. The rank-2 toric code, its symmetries, and UV/IR mixing
[**SP** & X-G Wen, PRB 106, 045145 (2022)]
[Y-T Oh, **SP**, JH Han, Y You, H-Y Lee, arXiv:2301.04706]

HIGHER-FORM SYMMETRIES

Consider bosonic lattice model with Hilbert space $\mathcal{H} = \bigotimes \mathcal{H}_i$.

Ordinary symmetries act on entire lattice.

$$U = \prod_{i \in \text{lattice}} U_i \quad \text{and} \quad UH = HU$$

p -form symmetry acts on codimension p ($d - p$ dimensional)
closed hypersurface Σ_{d-p}

$$U(\Sigma_{d-p}) = \prod_{i \in \Sigma_{d-p}} U_i \quad \text{and} \quad U(\Sigma_{d-p})H = HU(\Sigma_{d-p}) \quad \forall \Sigma_{d-p}$$

► For a G p -form symmetry, $U : Z_{d-p}(\text{lattice}; G) \rightarrow \mathcal{U}(\mathcal{H})$

HIGHER-FORM SYMMETRIES


Ordinary (0-form) symmetries transform operators acting on points

➤ Ex) $U(1)^{(0)}$ symmetry: $U_\alpha b_i U_\alpha^\dagger = \exp[i\alpha] b_i$

p -form symmetry transform operators acting on a p dimensional hypersurface C_p

➤ Ex) $U(1)^{(p)}$ symmetry:

$$U_\alpha(\Sigma_{d-p}) W(C_p) U_\alpha^\dagger(\Sigma_{d-p}) = \exp[i\alpha \#(\Sigma_{d-p}, C_p)] W(C_p)$$



$\#(\Sigma_{d-p}, C_p) \neq 0$ only if Σ_{d-p} and C_p are non-contractible

Just as b_i creates a bosonic particle at site i , $W(C_p)$ creates a

p -dimensional excitation at C_p

HIGHER-FORM SYMMETRIES

Ordinary (0-form) symmetries transform operators acting on points

► Ex) $U(1)^{(0)}$ symmetry: $U = b, U^\dagger = \exp[i\alpha] b.$

p -form symmetries
hypersurface

$U(1)^{(p)}$ Conservation laws:

$p = 0$: Particle number conserved

$p = 1$: String flux conserved

$p = 2$: Membrane flux conserved

Just as b_i creates a bosonic particle at site i , $W(C_p)$ creates a

p -dimensional excitation at C_p

EXAMPLE: 2+1D \mathbb{Z}_2 TORIC CODE

Square lattice with one qubit on each edge with $\Delta_e, \Delta_m > 0$


$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e$$

Exact $\mathbb{Z}_N^{(1)} \times \mathbb{Z}_N^{(1)}$ symmetry:

$$U_m(\gamma) = \prod_{e \in \gamma} X_e$$

$$U_e(\hat{\gamma}) = \prod_{e \perp \hat{\gamma}} Z_e$$

$U_m(\gamma)$ and $U_e(\hat{\gamma})$ do not commute: manifestation of mixed 't Hooft anomaly



EXAMPLE: 2+1D \mathbb{Z}_2 TORIC CODE

Square lattice with one qubit on each edge with $\Delta_e, \Delta_m > 0$

$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e$$

Exact $\mathbb{Z}_N^{(1)} \times \mathbb{Z}_N^{(1)}$ symmetry:

$$U_m(\gamma) = \prod_{e \in \gamma} X_e$$

$$U_e(\hat{\gamma}) = \prod_{e \perp \hat{\gamma}} Z_e$$

$U_m(\gamma)$ and $U_e(\hat{\gamma})$ do not commute: manifestation of mixed 't Hooft anomaly

$$E \geq \Delta_m \quad \langle U_m(\gamma_1) \rangle = \langle U_m(\gamma_2) \rangle \text{ if } \gamma_1 = \gamma_2$$

Faithful 1-form symmetry

$$E < \Delta_m \quad \langle U_m(\gamma_1) \rangle = \langle U_m(\gamma_2) \rangle \text{ iff } [\gamma_1] = [\gamma_2] \in H_1(\text{lattice}; \mathbb{Z}_N)$$

Topological 1-form symmetry

$$\begin{aligned} U_m(\gamma_2 = \gamma_1 + \partial M) &= U_m(\gamma_1) \prod_{p \in M} \prod_{e \in \partial p} X_e \\ &= U_m(\gamma_1) \end{aligned}$$

EXAMPLE: 2+1D \mathbb{Z}_2 TORIC CODE

Square lattice with one qubit on each edge with $\Delta_e, \Delta_m > 0$

$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e$$

Exact $\mathbb{Z}_N^{(1)} \times \mathbb{Z}_N^{(1)}$ symmetry:

$$U_m(\gamma) = \prod_{e \in \gamma} X_e$$

$$U_e(\hat{\gamma}) = \prod_{e \perp \hat{\gamma}} Z_e$$

$U_m(\gamma)$ and $U_e(\hat{\gamma})$ do not commute: manifestation of mixed 't Hooft anomaly

Ground state subspace ($E < \Delta_e, \Delta_m$)

$U_e(\hat{\gamma})$ ($U_m(\gamma)$) is the charged operator of $U_m(\gamma)$ ($U_e(\hat{\gamma})$)

➤ $W_m(\hat{\gamma}) = U_e(\hat{\gamma})$ and $W_e(\gamma) = U_m(\gamma)$

$$\langle W_m(\hat{\gamma} = \partial \hat{M}) \rangle = \langle W_e(\gamma = \partial M) \rangle = 1$$

➤ $\mathbb{Z}_N^{(1)} \times \mathbb{Z}_N^{(1)}$ SSB phase

\mathbb{Z}_2 TORIC CODE IN TRANSVERSE FIELD

$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e - h_x \sum_e X_e$$

$$H U_m(\gamma) = U_m(\gamma) H$$

$$H U_e(\hat{\gamma}) \neq U_e(\hat{\gamma}) H$$

X_ℓ no longer excites e anyons on $\partial\ell$. The operator that does is $\tilde{X}_\ell = U X_\ell U^\dagger$, dressed due to quantum fluctuations

► U is a unitary such that \tilde{X}_ℓ is a local operator, acting on a neighborhood of edges near ℓ (fattened operator)

For $h_x \ll \Delta_e$, low-energy states satisfy $\prod_{e \in \delta s} \tilde{Z}_e = 1 \implies$ emergent

conservation law (\mathbb{Z}_N electric flux) \implies emergent topological $\mathbb{Z}_N^{(1)}$

symmetry


$$U_e(\hat{\gamma}) = \prod_{e \perp \hat{\gamma}} \tilde{Z}_e$$

\mathbb{Z}_2 TORIC CODE IN TRANSVERSE FIELD

$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e - h_x \sum_e X_e$$

$$H U_m(\gamma) = U_m(\gamma) H$$

$$H U_e(\hat{\gamma}) \neq U_e(\hat{\gamma}) H$$



EFFECTIVE HAMILTONIAN

$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e - h_x \sum_e X_e \quad 0 < h_x \ll \Delta_e$$

Effective Hamiltonian for $E < \tilde{\Delta}_e$ describes e anyon free sub-Hilbert space. Low-energy allowed operator: $W(\gamma) = \prod_{e \in \gamma} \tilde{X}_e$.

$$H_{e \text{ free}} = -\frac{\Delta_m}{2} \sum_p W(\partial p) - h_x \sum_{\gamma \in B_1(\text{lattice}; \mathbb{Z}_N)} \epsilon_\gamma W(\gamma) - h_x \sum_{\gamma \in H_1(\text{lattice}; \mathbb{Z}_N)} \epsilon_\gamma W(\gamma)$$

$$\epsilon_\gamma \sim (h_x / \Delta_e)^{|\gamma| - 1}$$

EFFECTIVE HAMILTONIAN

$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e - h_x \sum_e X_e \quad 0 < h_x \ll \Delta_e$$

Effective Hamiltonian for $E < \tilde{\Delta}_e$ describes e anyon free sub-Hilbert space. Low-energy allowed operator: $W(\gamma) = \prod_{e \in \gamma} \tilde{X}_e$.

$$H_{e \text{ free}} = -\frac{\Delta_m}{2} \sum_p W(\partial p) - h_x \sum_{\gamma \in B_1(\text{lattice}; \mathbb{Z}_N)} \epsilon_\gamma W(\gamma) - h_x \sum_{\gamma \in H_1(\text{lattice}; \mathbb{Z}_N)} \epsilon_\gamma W(\gamma)$$

$$\epsilon_\gamma \sim (h_x / \Delta_e)^{|\gamma|-1}$$

$$\mathcal{O}(e^{-L})$$

- Exact emergent topological $\mathbb{Z}_N^{(1)}$ symmetry: $\tilde{X}_e \rightarrow s_e \tilde{X}_e \quad \prod_{e \in \partial p} s_e = 1 \forall p$
- $\langle W \rangle = 1$ for contractible loops. $\mathbb{Z}_N^{(1)}$ SSB phase

EFFECTIVE HAMILTONIAN

$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e - h_x \sum_e X_e - h_z \sum_e Z_e$$

Effective Hamiltonian for $E < \tilde{\Delta}_e$ describes e anyon free sub-Hilbert space. Low-

Deconfined phase of \mathbb{Z}_2 gauge theory
is SSB phase of exact emergent

H_e free

$\mathbb{Z}_N^{(1)} \times \mathbb{Z}_N^{(1)}$ symmetry

$$\epsilon_\gamma \sim (h_x / \Delta_e)^{|\gamma|-1}$$

$$\mathcal{O}(e^{-L})$$

- Exact emergent topological $\mathbb{Z}_N^{(1)}$ symmetry: $\tilde{X}_e \rightarrow s_e \tilde{X}_e$ $\prod_{e \in \partial p} s_e = 1 \forall p$
- $\langle W \rangle = 1$ for contractible loops. $\mathbb{Z}_N^{(1)}$ SSB phase

EVIDENCE FROM NUMERICS


Precise form of dressed operators is not generally unique, depending on microscopic details

- Can be numerically constructed as a *matrix product operator* [Cian, Hafezi, & Barkeshli, arXiv:2209.14302]

Toric code with $h_x = 0.15$,
 $h_z = 0.05$ and $\Delta_e = \Delta_m = 2$

$$W_{\text{FP}} = \prod_{e \in \gamma} X_e \implies \langle W_{\text{FP}} \rangle \sim e^{-|\gamma|/\xi}$$

$$W = \left\langle \prod_{e \in \gamma} \tilde{X}_e \right\rangle \implies \langle W \rangle = 1$$



EMERGENCE WITHOUT SSB


$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e - h_x \sum_e X_e - h_z \sum_e Z_e$$

When $h_x \gg \Delta_e$, e anyons condense (Higgs phase)

- No e free low-energy subspace \implies no emergent conservation law (\mathbb{Z}_N electric flux) \implies no emergent $\mathbb{Z}_N^{(1)}$ symmetry
- There can be an m free low-energy subspace \implies emergent conservation law (\mathbb{Z}_N magnetic flux) \implies emergent $\mathbb{Z}_N^{(1)}$ symmetry

EMERGENCE WITHOUT SSB

$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e - h_x \sum_e X_e - h_z \sum_e Z_e \quad h_x \gg \Delta_e$$




EMERGENCE WITHOUT SSB

$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e - h_x \sum_e X_e - h_z \sum_e Z_e \quad h_x \gg \Delta_e$$

Exact emergent *trivial* without SSB:

- With PBC, unbroken exact emergent $\mathbb{Z}_N^{(1)}$ symmetry is **trivial**
- With a **spatial boundary**, unbroken exact emergent $\mathbb{Z}_N^{(1)}$ symmetry can be **nontrivial** (e.g., SPT phase with $\mathbb{Z}_N^{(1)}$ SSB on boundary)

No SSB (Higgs “phase”)




EMERGENCE WITHOUT SSB

$$H = -\frac{\Delta_e}{2} \sum_s \prod_{e \in \delta s} Z_e - \frac{\Delta_m}{2} \sum_p \prod_{e \in \partial p} X_e - h_x \sum_e X_e - h_z \sum_e Z_e \quad h_x \gg \Delta_e$$

Exact emergent *trivial* with SSB:

- **With PBC**, spontaneously broken exact emergent $\mathbb{Z}_N^{(1)}$ symmetry is **nontrivial**
- **On a disk**, spontaneously broken exact emergent $\mathbb{Z}_N^{(1)}$ symmetry is **trivial**




No SSB (Higgs “phase”)

GENERAL DISCUSSION

Typical Hamiltonians do not have exact higher-form symmetries.


- Consider $H = H_0 + H_1$ in $d = 3$. Suppose H_0 has a 1-form symmetry and H_1 is a generic local perturbation



GENERAL DISCUSSION

Typical Hamiltonians do not have exact higher-form symmetries.

- Consider $H = H_0 + H_1$ in $d = 3$. Suppose H_0 has a 1-form symmetry and H_1 is a generic local perturbation



General H_1 violates the flux conservation law of the 1-form symmetry, hence explicitly breaking H_0 's 1-form symmetry

EMERGENT HIGHER-FORM SYMMETRY

Ends of strings cost finite energy (Gauss law term in H_0)

- True gapped topological excitations are modified due to H_1 . Correspond to ends of modified (fattened) strings
- Below gap, only closed modified loops \implies emergent string flux conservation \implies emergent 1-form symmetry

General p -form symmetry: gapped $(p - 1)$ dimensional topological excitations. At energies below their gap, only modified (fattened) p -branes \implies emergent p -form symmetry.

- No 0-form symmetry analog ($p > 0$ only)

EXACT EMERGENT SYMMETRY

A conservation law implies a symmetry. This conservation law is violated by charged operators of the symmetry.

Emergent 0-form symmetries are approximate symmetries

- Emergent conservation laws *are* violated by local operators, which typically appear in a low-energy effective theory.

Emergent higher-form symmetries are exact symmetries

- Emergent conservation law *cannot* be violated by any local operators. The (local) low-energy effective theory will have the higher-form symmetry as an exact symmetry.

EXACT EMERGENT SYMMETRY

A conservation law implies a symmetry. This conservation law is violated in the SSB phase.

Emergent

Emergent **0-form symmetry** SSB phase:

- Discrete: no GSD
- Continuous: gapped goldstone modes

➤ Emergent

operator

theory.

etries

effective

Emergent

Emergent **higher-form symmetry** SSB phase:

➤ Eme

➤ Discrete: exact GSD

➤ Continuous: gapless goldstone modes

oper

the l

y local

ll have

EXACT EMERGENT HIGHER-FORM SYMMETRIES

Exact higher-form symmetries are rare, but **exact emergent higher-form symmetry** are common.

- Most ordinary SSB phases have **exact emergent higher-form symmetries** (To be posted on arXiv this spring... hopefully 🙏🙏🤞🤞)
- Known examples in $d = 3$: Ising ferromagnets $(\mathbb{Z}_2^{(0)} \times \mathbb{Z}_2^{(3)})$, bosonic superfluids $(U(1)^{(0)} \times U(1)^{(2)})$

THE PLAN FOR THIS TALK

Explore how higher-form symmetries arise in
topological phases of quantum matter

1. How higher-form symmetries emerge & exact emergent symmetries
[**SP** & X-G Wen, arXiv:2301.05261]
2. Symmetry Protected Trivial (SPT) phase protected by higher-form symmetries
[**SP** & X-G Wen, PRB 107, 075112 (2023)]
3. The rank-2 toric code, its symmetries, and UV/IR mixing
[**SP** & X-G Wen, PRB 106, 045145 (2022)]
[Y-T Oh, **SP**, JH Han, Y You, H-Y Lee, arXiv:2301.04706]

THE PLAN FOR THIS TALK

Explore how higher-form symmetries arise in
topological phases of quantum matter

1. How higher-form symmetries emerge & exact emergent symmetries
[SP & X-G Wen, arXiv:2301.05261]
2. Symmetry Protected Trivial (SPT) phase protected by higher-form symmetries
[SP & X-G Wen, PRB 107, 075112 (2023)]
3. The rank-2 toric code, its symmetries, and UV/IR mixing
[SP & X-G Wen, PRB 106, 045145 (2022)]
[Y-T Oh, SP, JH Han, Y You, H-Y Lee, arXiv:2301.04706]

2+1D ABELIAN BOSONIC TOPOLOGICAL ORDER

Deconfined phase of 2+1D \mathbb{Z}_N gauge theory is an exact emergent anomalous $\mathbb{Z}_N^{(1)} \times \mathbb{Z}_N^{(1)}$ SSB phase

➤ TQFT describing ground states: $S = \frac{iN}{2\pi} \int dt d^2x \epsilon^{\mu\nu\rho} a_\mu^1 \partial_\nu a_\rho^2$

Ground states of a general abelian bosonic topological order in 2+1D described by

$$S = \frac{iK_{IJ}}{4\pi} \int dt d^2x \epsilon^{\mu\nu\rho} a_\mu^I \partial_\nu a_\rho^J \quad K_{IJ} \in \begin{cases} \mathbb{Z}, & I \neq J \\ 2\mathbb{Z} & I = J \end{cases}, \quad K_{IJ} = K_{JI}$$

➤ Anomalous $\mathbb{Z}_{k_1}^{(1)} \times \mathbb{Z}_{k_2}^{(1)} \times \cdots$ symmetry

'T HOOFT ANOMALY AND SPT ORDER

The $\mathbb{Z}_{k_1}^{(1)} \times \mathbb{Z}_{k_2}^{(1)} \times \cdots$ symmetry is **anomalous**:

1. Theory has to be in a **gapless, SSB, or T.O.** phase.
2. The anomalous part of the symmetry **cannot be gauged**

$$Z[A] \rightarrow Z[A + d\omega] = e^{i\theta(A,\omega)} Z[A]$$

3. Theory is **perfectly well defined** by itself (with $A = 0$)
4. Can be gauged if it resides on the **boundary of an SPT**.

Question: What is the $\mathbb{Z}_{k_1}^{(1)} \times \mathbb{Z}_{k_2}^{(1)} \times \cdots$ SPT whose boundary has topological order described by the K matrix?

THE SPT INVARIANT

Turn on background fields \mathcal{A}^I : $S[\mathcal{A}] = \frac{iK_{IJ}}{4\pi} \int_{X_3} dt d^2x \epsilon^{\mu\nu\rho} a_\mu^I \left(\partial_\nu a_\rho^J - \frac{1}{2} \mathcal{A}_{\nu\rho}^J \right)$

$$\begin{aligned} a_\mu^I &\rightarrow a_\mu^I + \omega_\mu^I \\ \mathcal{A}_{\mu\nu}^I &\rightarrow \mathcal{A}_{\mu\nu}^I + \partial_\mu \omega_\nu^I - \partial_\nu \omega_\mu^I \end{aligned} \quad Z[X_3, \mathcal{A}] \rightarrow e^{-\frac{iK_{IJ}}{8\pi} \int_{X_3} dt d^2x \epsilon^{\mu\nu\rho} \omega_\mu^I \mathcal{A}_{\nu\rho}^J} Z[X_3, \mathcal{A}]$$

Extend \mathcal{A} from 2+1D X_3 to 3+1D Y_4 such that $\partial Y_4 = X_3$

$$Z_{\text{g-inv}}[Y_4, \mathcal{A}] = e^{\frac{iK_{IJ}}{16\pi} \int_{Y_4} dt d^3x \epsilon^{\mu\nu\rho\sigma} \mathcal{A}_{\mu\nu}^I \mathcal{A}_{\rho\sigma}^J} Z[X_3 = \partial Y_4, \mathcal{A}]$$

► Invertible theory characterizing the **SPT order**:

$$Z_{\text{SPT}}[Y_4, \mathcal{A}] = \exp \left[\frac{iK_{IJ}}{16\pi} \int_{Y_4} dt d^3x \epsilon^{\mu\nu\rho\sigma} \mathcal{A}_{\mu\nu}^I \mathcal{A}_{\rho\sigma}^J \right]$$

TWISTED $U(1)$ GAUGE THEORY

Z_{SPT} is the IR FP theory. What system is in this SPT phase?

► Trick: “ungauge” \mathcal{A}^I by $\mathcal{A}_{\mu\nu}^I \rightarrow \partial_\mu a_\nu^I - \partial_\nu a_\mu^I \equiv f_{\mu\nu}^I$

$$S_{\text{SPT}} = \frac{iK_{IJ}}{16\pi} \int_{Y_4} dt d^3x \epsilon^{\mu\nu\rho\sigma} f_{\mu\nu}^I f_{\rho\sigma}^J \equiv \frac{iK_{IJ}}{32\pi} \int_{Y_4} dt d^3x \vec{e}^I \cdot \vec{b}^J$$

► General twisted $U(1)$ gauge theory

$$S = \frac{1}{4g^2} \int_{Y_4} f_{\mu\nu}^I f^I{}^{\mu\nu} + \frac{iK_{IJ}}{16\pi} \int_{Y_4} dt d^3x \epsilon^{\mu\nu\rho\sigma} f_{\mu\nu}^I f_{\rho\sigma}^J$$


The confined phase ($g \rightarrow \infty$) of twisted $U(1)$ gauge theory is a

$$\mathbb{Z}_{k_1}^{(1)} \times \mathbb{Z}_{k_2}^{(1)} \times \dots \text{ SPT}$$

TWISTED $U(1)$ GAUGE THEORY


$U(1)$ gauge theory:

- $g > g_c$: monopoles condense



Twisted $U(1)$ gauge theory

- $g > g_c$: dyons condense



$(q_e, q_m) = (2, 1)$ dyon $\rightarrow \mathbb{Z}_2^{(1)}$ SPT

Boundary: $\nu = 1/2$ bosonic FHQ

$(q_e^{1,2}, q_m^{2,1}) = (2, 1)$ dyons $\rightarrow \mathbb{Z}_2^{(1)} \times \mathbb{Z}_2^{(1)}$ SPT

Boundary: \mathbb{Z}_2 topological order


Need magnetic monopoles (dyons), **why care?**

- **Quantum spin ice** on a breathing pyrochlore lattice

EMERGENT HIGHER-FORM SPT

$\mathbb{Z}_{k_1}^{(1)} \times \mathbb{Z}_{k_2}^{(1)} \times \dots$ symmetry operator (with $g \rightarrow \infty$)

$$U^{(I)}(\Sigma_2) = \exp \left[\frac{iK^{IJ}}{2\pi} \int_{\Sigma_2} da^J \right] = \exp \left[\frac{iK^{IJ}}{2\pi} \int_{\partial\Sigma_2} a^J \right]$$



- Nontrivial $\mathbb{Z}_{k_1}^{(1)} \times \mathbb{Z}_{k_2}^{(1)} \times \dots$ exact emergent symmetry (exact conservation law in low-energy bulk+boundary theory)
- Robustness of exact emergent SSB boundary phase (topological order) reflects exact emergent SPT order

THE PLAN FOR THIS TALK

Explore how higher-form symmetries arise in
topological phases of quantum matter

1. How higher-form symmetries emerge & exact emergent symmetries
[**SP** & X-G Wen, arXiv:2301.05261]
2. Symmetry Protected Trivial (SPT) phase protected by higher-form symmetries
[**SP** & X-G Wen, PRB 107, 075112 (2023)]
3. The rank-2 toric code, its symmetries, and UV/IR mixing
[**SP** & X-G Wen, PRB 106, 045145 (2022)]
[Y-T Oh, **SP**, JH Han, Y You, H-Y Lee, arXiv:2301.04706]

THE PLAN FOR THIS TALK

Explore how higher-form symmetries arise in
topological phases of quantum matter

1. How higher-form symmetries emerge & exact emergent symmetries
[**SP** & X-G Wen, arXiv:2301.05261]
2. Symmetry Protected Trivial (SPT) phase protected by higher-form symmetries
[**SP** & X-G Wen, PRB 107, 075112 (2023)]
3. The rank-2 toric code, its symmetries, and UV/IR mixing
[**SP** & X-G Wen, PRB 106, 045145 (2022)]
[Y-T Oh, **SP**, JH Han, Y You, H-Y Lee, arXiv:2301.04706]

2+1D RANK-2 TORIC CODE

Square lattice with two \mathbb{Z}_N quantum rotors on each site and one \mathbb{Z}_N quantum rotors on each plaquette

- **Toric code**: exactly solvable point in deconfined phase of vector \mathbb{Z}_N gauge theory
- **Rank-2 Toric Code (R2TC)**: exactly solvable point in deconfined phase of symmetric rank-2 tensor \mathbb{Z}_N gauge theory

2+1D RANK-2 TORIC CODE

Square lattice with two \mathbb{Z}_N quantum rotors on each site and one \mathbb{Z}_N quantum rotors on each plaquette


$$H_{\text{R2TC}} = - \sum_{x,y} \left(G_{x,y} + F_{x,y}^{(x)} + F_{x,y}^{(y)} + \text{h.c.} \right)$$

\mathbb{Z}_N clock operators:


$$Z_j X_i = \omega^{\delta_{i,j}} X_i Z_j$$

$$\omega = \exp[2\pi i/N]$$


$$X_i^N = Z_i^N = 1$$



● Z_1 ● Z_2 ● Z_3



● X_1 ● X_2 ● X_3




R2TC EXCITATIONS

$$H_{\text{R2TC}} = - \sum_{x,y} \left(G_{x,y} + F_{x,y}^{(x)} + F_{x,y}^{(y)} + \text{h.c.} \right)$$



- Exactly solvable because: $[G_{x,y}, F_{x',y'}^{(x)}] = [G_{x,y}, F_{x',y'}^{(y)}] = 0$
- A gapped excited states $|\psi\rangle$ satisfies:

$$G_{x,y} |\psi\rangle = \omega^{e_{x,y}} |\psi\rangle \quad F_{x,y}^{(x)} |\psi\rangle = \omega^{m_{x,y}^{(x)}} |\psi\rangle \quad F_{x,y}^{(y)} |\psi\rangle = \omega^{m_{x,y}^{(y)}} |\psi\rangle$$

\mathbb{Z}_N charge



\mathbb{Z}_N fluxes



SOME R2TC PROPERTIES

Restricted Mobility:

- e hops by N in all directions \rightarrow pseudo-fracton
- $m^{(i)}$ hops by 1 in i -direction, by N in transverse direction \rightarrow pseudo-lineon

Position-dependent braiding:

- Statistics of e_{x_e, y_e} and $m_{x_m, y_m}^{(i)}$ depend on $(x_e - x_m, y_e - y_m)$

UV/IR Mixing:

- Anyon counting on a $L_x \times L_y$ torus yields:

$$\text{GSD} = N^3 \gcd(L_x, N) \gcd(L_y, N) \gcd(L_x, L_y, N)$$

R2TC GENERALIZED SYMMETRIES

Given exotic features, it is desirable to have a **generalized symmetry** understanding


Can use SET machinery, so why generalized symmetry?

- Captures ground states' structures directly (**naturalness**)
- Easily generalizable to higher dimensions (**practical utility**)
- As we'll see, provides a **natural unifying picture** for intrinsic topological order and fracton topological order (**conceptually useful**)

R2TC GENERALIZED SYMMETRIES

There are six discrete 1-form symmetries, all **spontaneously broken**, and three types of mixed 't Hooft anomalies

- Three symmetries are **topological only within a sublattice**




UV/IR MIXING AND SYMMETRY

The UV/IR mixing originates entirely from the **sublattice topological 1-form symmetries**:

- Low-energy emergent symmetries (IR) are defined in terms of the lattice regularization (UV)

Conjecture: All UV/IR mixing in “topological phases” arises from SSB of subsystem generalized symmetries



SUMMARY

1. Emergent higher-form symmetries are exact emergent symmetries and are common.
[**SP** & X-G Wen, arXiv:2301.05261]
2. Emergent higher-form SPTs can appear in confined phases of twisted gauge theory
[**SP** & X-G Wen, PRB 107, 075112 (2023)]
3. UV/IR mixing in gapped phases is naturally understood using generalized symmetries
[**SP** & X-G Wen, PRB 106, 045145 (2022)]
[Y-T Oh, **SP**, JH Han, Y You, H-Y Lee, arXiv:2301.04706]