

Programme: RSD2 (Group 5)

Leader : Lim Jun Wei

Member : Chia Ming Yi

Member : Ong Yi Xin

Marks: _____/50

Answer ALL questions. All relevant and detailed workings must be shown.

Question 1

Let $A = \{18, 20, 26, 32, 38\}$. A relation R on set A is defined by xRy if and only if $x \equiv y \pmod{4}$.

- (a) Write R as a set of ordered pairs. (2 marks)
- (b) Find the in-degree and out-degree of each vertex. (2 marks)
- (c) Determine whether relation R is reflexive, irreflexive, symmetric, asymmetric, antisymmetric or transitive. Give a counterexample if your answer is "No". (9 marks)

- (d) Another relation S on set A whose matrix is given by $M_S = \begin{bmatrix} 0 & 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 & 1 \end{bmatrix}$. Find the matrix of $R \circ S^{-1}$. (4 marks)

Question 2

Let $f(x) = \lceil \frac{x^2+x-2}{x+1} \rceil$ be a function from $A = \{1, 2, 3, 4, 5, 6\}$ to $B = \{0, 1, 2, 3, 4, 5, 6, 7\}$.

- (a) Determine whether f is one-to-one, onto and everywhere defined. Explain your answers. (6 marks)
- (b) Is f bijective? Explain your answer. (2 marks)

Question 3

Without constructing truth table, obtain the principal conjunctive normal form (PCNF) of $A \equiv [(p \vee (q \rightarrow \sim r)) \vee \sim q] \rightarrow \sim(p \vee \sim r)$. (8 marks)

Question 4

Show the scope of each quantifier in the following expression $(\exists y)[\sim P(x, y) \rightarrow (\forall x)(Q(x) \vee \sim R(x, y))]$. Determine whether the given expression is a statement or not. If not, state the reason. (4 marks)

Question 5


The following arithmetic expression is given in Polish form.

$+ - \times 5 6 + 9 \div 8 2 - 7 \times 4 5$

- (a) Calculate the value represented by the expression. (3 marks)
- (b) Write the corresponding fully parenthesised expression and draw its binary tree. (6 marks)
- (c) Write the expression in postfix and infix form. (4 marks)


[End of Question Paper]

Question 1



a) $R = \{ (18,18), (18,26), (18,38), (20,20), (20,32), (26,18), (26,26), (26,38), (32,20), (32,32), (38,18), (38,26), (38,38) \}$

b)



	18	20	26	32	38
In-degree	3	3	3	2	2
Out-degree	3	3	3	2	2

c) - Relation R is reflexive

- Relation R is not irreflexive, $\because 18R18$

- Relation R is symmetric.

- Relation R is not asymmetric, $\because 18R18$ or $18R26$ but $26R18$.- Relation R is not antisymmetric, $\because 18R26$ and $26R18$ but $18 \neq 26$.

- Relation R is transitive.

d)

$$M_R = \begin{bmatrix} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \end{bmatrix}$$

$$M_S^{-1} = (M_S)^T$$

$$= \begin{bmatrix} 0 & 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 \end{bmatrix}$$

$$M_{R \circ S^{-1}} = M_S^{-1} \odot M_R$$

$$= \begin{bmatrix} 0 & 1 & 0 & 1 & 1 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 \end{bmatrix} \odot \begin{bmatrix} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 1 \end{bmatrix}$$


$$= \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 \end{bmatrix}$$

Question 2

a)

A

B



a) $\because f(1)=0, f(2)=2, f(3)=3, f(4)=4, f(5)=5, f(6)=6, \therefore f$ is one to one.

$\because \text{Ran}(f) = \{0, 2, 3, 4, 5, 6\} \neq B, \therefore f$ is not onto.

$\because \text{Dom}(f) = \{1, 2, 3, 4, 5, 6\} = A, \therefore f$ is everywhere defined.

b) f is not bijective because $\text{Ran}(f) = \{0, 2, 3, 4, 5, 6\} \neq B$ (not surjective) although f is injective.

Question 3

$$\begin{aligned}
 & [(p \vee (q \rightarrow \sim r)) \vee \sim q] \rightarrow \sim (p \vee \sim r) \\
 &= [(p \vee (\sim q \vee \sim r)) \vee \sim q] \rightarrow \sim (p \vee \sim r) \\
 &= [p \vee \sim q \vee \sim r \vee \sim q] \rightarrow \sim (p \vee \sim r) \\
 &= [p \vee \sim q \vee \sim r] \rightarrow \sim (p \vee \sim r) \\
 &= \sim (p \vee \sim q \vee \sim r) \vee (\sim p \wedge r) \\
 &= (\sim p \wedge q \wedge r) \vee (\sim p \wedge r) \\
 &= (\sim p \wedge q \wedge r) \vee [\sim p \wedge (q \vee \sim q) \wedge r] \\
 &= (\sim p \wedge q \wedge r) \vee (\sim p \wedge q \wedge r) \vee (\sim p \wedge \sim q \wedge r) \\
 &= (\sim p \wedge q \wedge r) \vee (\sim p \wedge \sim q \wedge r)
 \end{aligned}$$

$$\text{PDNF } A = (\bar{p}qr + \bar{p}\bar{q}r)$$

$$\text{PDNF } \sim A = pqr + pq\bar{r} + p\bar{q}r + p\bar{q}\bar{r} + \bar{p}q\bar{r} + \bar{p}\bar{q}\bar{r}$$

$$\text{PCNF } A = (\bar{p} + \bar{q} + \bar{r})(\bar{p} + \bar{q} + r)(\bar{p} + q + \bar{r})(\bar{p} + q + r)(p + \bar{q} + r)(p + q + r)$$

$$\text{PCNF } \sim A = (p + \bar{q} + \bar{r})(p + q + \bar{r})$$

Question 4

$$\exists y : \sim P(x, y) \rightarrow (\forall x) (Q(x) \vee \sim R(x, y))$$

$$\forall x : Q(x) \vee \sim R(x, y)$$

The given expression is not a statement because the variable x in $\sim P(x, y)$ is a free variable.

Question 5

a) Infix form :

$$[(5 \times 6) - (9 + (8 \div 2))] + [7 - (4 \times 5)]$$

$$= [30 - (9 + 4)] + [7 - 20]$$


$$= [30 - 13] + [-13]$$

$$= 17 - 13$$

$$= 4$$

b) Corresponding fully parenthesised expression :

$$[(5 \times 6) - (9 + (8 \div 2))] + [7 - (4 \times 5)]$$



c) Postfix form :

$$56 \times 982 \div + - 745 \times - +$$

Infix form :

$$[(5 \times 6) - (9 (8 \div 2))] + [7 - (4 \times 5)]$$