

Epipolar Geometry

Three Questions

1. Correspondence Geometry

- Given an image point x in the first view, how does this constrain the position of the corresponding point x' in the second image?

2. Camera Geometry (motion)

- Given a set of corresponding image points $x_i \leftrightarrow x'_i, i = 1, \dots, n$
- What are the cameras P and P' for the two views?
- What is the geometric transformation between the two views?

3. Scene Geometry (structure)

- Given corresponding image points $x_i \leftrightarrow x'_i$ and cameras P, P'
- What is the position of the point X in space?

Geometry

- Epipolar plane π is the plane formed by P, P', X
- Plane passes through camera planes and points x, x' lie on the edges of the plane
- Epipoles - Intersection of baseline with image plane
 - Projection of projection center in other image
 - Vanishing point of camera motion direction
- Baseline - Line formed by π and the epipoles \rightarrow Translation between cameras
- Cross Product

$$a \times b = \begin{bmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{bmatrix} b = [a_x]b$$

- Triple Scalar Product \rightarrow 3 points are coplanar

$$(a \times b) \cdot c = 0$$

- Equations

$$P' = RP + t$$

$$p = MP, M = [I|0]$$

$$p' = M'P', M' = [R|t]$$

P is the point in space, p, p' are points in image

$$\vec{O'p'}, \vec{OO'}, \vec{Op} \text{ are coplanar} \implies p' \cdot [t \times (Rp)] = 0, \begin{cases} p = (u, v, 1)^T \\ p' = (u', v', 1)^T \end{cases}$$

$$p' \cdot [t \times (Rp)] = 0 \implies p' \cdot [(t \times R)p] = 0 \implies p'([t_x]R)p = 0$$

$$\underbrace{E = [t_x]R}_{\text{Essential Matrix}} \implies \boxed{p'Ep = 0}$$

- Essential Matrix
 - E can be used to find rotation and translation
 - Computed from point correspondences
- Uncalibrated camera
 - $\hat{p} = M^{-1}p$ and $\hat{p}' = M'_{int}p'$
 - p and p' are points in pixel coordinates corresponding to \hat{p}, \hat{p}'
 - $p'^T F p = 0$
 - $F = M'^{-T}_{int} E M^{-1}_{int}$
- Properties of fundamental and essential matrix
 - With calibration parameters \rightarrow essential
 - Without \rightarrow fundamental
 - Matrix is 3x3
 - **Transpose:** If F is essential matrix of cameras P, P' $\Rightarrow F^T$ is essential matrix of camera P', P
 - **Epipolar Lines:** p and p' as points in the projective plane $\Rightarrow Fp$ is projective line in the right image
 - $l' = Fp, l = F^T p'$
 - **Epipole:** For any p the epipolar line $l' = Fp$ contains the epipole e' $\Rightarrow (e'^T F)p = 0$ for all p
 - $e'^T F = 0$ and $Fe = 0$
 - Epipoles are in the Null Space of the Essential Matrix

- Fundamental Matrix
 - Encodes information of the intrinsic and extrinsic parameters
 - F is of rank 2 since S has rank 2 (R and M and M' have full rank)
 - Has 7 DOF
 - 9 elements \rightarrow scaling is not significant & $\det(F) = 0$
- Essential Matrix
 - Encodes only extrinsic parameters
 - Rank 2 since S has rank 2
 - Its two nonzero singular values are equal
 - Has only 5 DOF \rightarrow 3 rotation, 2 translation
- Scaling Ambiguity

$$P' = RP + t$$

$$p = \frac{P}{z^T P} \quad p' = \frac{RP + t}{z^T (RP + t)}$$

- Depth Z and Z' and t can only be recovered to a scale factor
- Computing the Fundamental Matrix from Point Correspondences
 - Defined by the equation $x_i'^T F x_i = 0$
 - For n point matches \rightarrow A is an $n \times 9$ matrix
 - Solve $Af = 0$
 - Homogeneous set of equations
 - f can be determined only up to a scale, so there are 8 unknowns
 - 8 Point Algorithm
 - Least Squares solution is the singular vector corresponding the smallest singular value of A \rightarrow last column of V in the SVD
 - This has a lot of error since the matrix is almost singular
 - Non-Linear Least Squares Approach
- Rectification
 - Image Reprojection
 - Reproject image planes onto common plane parallel to line between optical centers
 - Rotate left camera so epipoles go to infinity along the horizontal axis
 - Apply same rotation to the right camera
 - Rotate the right camera by R
 - Adjust the scale
- 3D Reconstruction

- **Stereo:** We know the viewing geometry (extrinsic parameters) and the intrinsic parameters \Rightarrow Find correspondences exploiting epipolar geometry, then reconstruct
- **Structure from motion:** Find correspondences \rightarrow estimate extrinsic parameters (rotation and translation) \rightarrow reconstruct
- **Uncalibrated Cameras:** Find correspondences \rightarrow compute projection matrices \rightarrow reconstruct up to a projective transformation
 - Don't know the position of camera respective to origin
 - Can't find the orientation of camera plane
- **Triangulation**
 - If cameras are intrinsically and extrinsically calibrated \Rightarrow find P as the midpoint of the common perpendicular to the two rays in space
- Intrinsically calibrated cameras
 - Compute the essential matrix E using normalized points
 - Select $M = [I|0], M' = [R|T] \rightarrow E = [T_x]R$
 - Find T and R using SVD of E

$$E = U\Sigma V^T$$

$$[T_x] = UZU^T \quad R = UWV^T \text{ or } R = UW^TV^T$$

$$W = \begin{bmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } Z = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

- Multiple solutions \rightarrow One with positive values is correct
- Linear Triangulation
 - Homogenous System

$$x = MX \quad x' = M'X$$

$$x \times MX = 0 \quad x' \times M'X' = 0$$

$$A = \begin{bmatrix} xm_3^T & -m_1^T \\ ym_3^T & -m_2^T \\ x'm_3^T & -m_3'^T \\ y'm_3^T & -m_2'^T \end{bmatrix}$$

$$AX = 0$$

- X is the last column of V in the SVD of A
- Projective Reconstruction Theorem
 - Assume we determine matching points x_i and x'_i . Then we can compute a unique Fundamental matrix F
 - The camera matrices M, M' cannot be recovered uniquely
 - Thus the reconstruction X_i is not unique
 - There exists a projective transformation H
 - Reconstructive Ambiguity

$$x_i = MX_i = (MH_P^{-1})(H_P X_i)$$

- Two different objects produce the same image since the cameras are uncalibrated
 - Find the original distorted by a projective transformation $H \rightarrow$ *Projective Reconstruction*
 - If reconstruction is derived from real images \rightarrow there is a true reconstruction that can produce the actual points X_i of the scene
- Consequences
 - We can compute a projective reconstruction of a scene from 2 views based on image correspondences alone
 - Don't need anything about poses or calibration
- Projective \rightarrow Metric
 - Compute homography H such that $X_{Ei} = HX_i$ for 5 or more control points X_{Ei}
 - Rectification using 5 points
 - Compute projective reconstruction
 - Find 5+ control points
 - Use them to solve for projective transformation (9 DOF) and rectify the projective reconstruction
- Stratified Reconstruction
 - Begin with projective reconstruction
 - Refine it to affine reconstruction
 - Parallel lines are parallel, ratios along parallel lines are correct
 - Reconstructed scene is then an affine transformation of the actual scene
 - Refine to a metric reconstruction
 - Angles and ratios are correct

- Reconstructed scene is then a scaled version of the actual scene