



# A MIP approach to tackle the Optimal Power Flow problem with probabilistic constraints

---

Álvaro Porras, *Concepción Domínguez*, Juan Miguel Morales, Salvador Pineda  
June 29<sup>th</sup>, 2022

OASYS Group, Universidad de Málaga

Introduction to Chance-Constrained Problems

General chance-constrained SAA MIP reformulation

Tightening and screening

Valid inequalities

Computational Results: OPF

# Introduction to Chance-Constrained Problems

---

# Introduction to Chance-Constrained Problems (CCPs)

- Applications: areas where decisions have to be made dealing with **random parameters (uncertainty)**. CCs ensure feasibility of the system with a **tolerable probability of constraint violation**.

# Introduction to Chance-Constrained Problems (CCPs)

- Applications: areas where decisions have to be made dealing with **random parameters (uncertainty)**. CCs ensure feasibility of the system with a **tolerable probability of constraint violation**.
- **Optimal Power Flow**: minimize the expected operating cost whilst guaranteeing that the system withstands unforeseen peaks of electrical load due to stochastic demand.

# Introduction to Chance-Constrained Problems (CCPs)

- Applications: areas where decisions have to be made dealing with **random parameters (uncertainty)**. CCs ensure feasibility of the system with a **tolerable probability of constraint violation**.
- **Optimal Power Flow**: minimize the expected operating cost whilst guaranteeing that the system withstands unforeseen peaks of electrical load due to stochastic demand.
- General (linear) formulation:

$$\begin{aligned} \min_{\mathbf{x}} \quad & \mathbf{c}^\top \mathbf{x} \\ \text{s.t.} \quad & \mathbf{x} \in X \\ & \mathbb{P} \left\{ \mathbf{a}_j(\omega)^\top \mathbf{x} \leq b_j(\omega), \forall j \right\} \geq 1 - \epsilon. \end{aligned}$$

# Chance-constrained SAA MIP reformulation (with Big-Ms)

- Sample Average Approx.: assume a finite discrete distribution  $\Rightarrow$  MIP reformulation

# Chance-constrained SAA MIP reformulation (with Big-Ms)

- Sample Average Approx.: assume a finite discrete distribution  $\Rightarrow$  MIP reformulation
- $s \in \mathcal{S}$  scenarios (with equal probabilities)



# Chance-constrained SAA MIP reformulation (with Big-Ms)

- Sample Average Approx.: assume a finite discrete distribution  $\Rightarrow$  MIP reformulation
- $s \in \mathcal{S}$  scenarios (with equal probabilities)
- $y_s \in \{0, 1\}$  such that  $y_s = 0 \Leftrightarrow$  the scenario  $s$  is satisfied

# Chance-constrained SAA MIP reformulation (with Big-Ms)

- Sample Average Approx.: assume a finite discrete distribution  $\Rightarrow$  MIP reformulation
- $s \in \mathcal{S}$  scenarios (with equal probabilities)
- $y_s \in \{0, 1\}$  such that  $y_s = 0 \Leftrightarrow$  the scenario  $s$  is satisfied

$$\min_x \quad c^\top x$$

$$\text{s.t.} \quad x \in X$$

$$\mathbb{P} \left\{ a_j(\omega)^\top x \leq b_j(\omega), \quad \forall j \right\} \geq 1 - \epsilon.$$

$$\sum_{s \in \mathcal{S}} y_s \leq p$$

$$y_s \in \{0, 1\} \quad \forall s$$

# Chance-constrained SAA MIP reformulation (with Big-Ms)

- Sample Average Approx.: assume a finite discrete distribution  $\Rightarrow$  MIP reformulation
- $s \in \mathcal{S}$  scenarios (with equal probabilities)
- $y_s \in \{0, 1\}$  such that  $y_s = 0 \Leftrightarrow$  the scenario  $s$  is satisfied

$$\begin{aligned} \min_x \quad & c^\top x \\ \text{s.t.} \quad & x \in X \\ & \mathbb{P} \left\{ a_j(\omega)^\top x \leq b_j(\omega), \forall j \right\} \geq 1 - \epsilon. \\ & \sum_{s \in \mathcal{S}} v_s \leq p \\ & v_s \in \{0, 1\} \quad \forall s \end{aligned}$$

$$\begin{aligned} \min_x \quad & c^\top x \\ \text{s.t.} \quad & x \in X \\ & a_j(\omega)^\top x \leq b_j(\omega) + M_{js} y_s, \quad \forall j, s \\ & \sum_{s \in \mathcal{S}} y_s \leq p \\ & y_s \in \{0, 1\}, \quad \forall s. \end{aligned}$$

## Tightening and screening

---

---

## Algorithm 1 Iterative Coefficient Strengthening ( $\kappa$ iterations)

---

**Initialization:**  $k \leftarrow 0$ ,  $M_{js}^0 \leftarrow \infty$ .

**while**  $k < \kappa$  **do**

**for**  $j \in \mathcal{J}$  and  $s \in \mathcal{S}$  **do**

**if**  $M_{js}^k > 0$  **then**

            1) Tightening phase: Solve

$$M_{js}^{k+1} \leftarrow \arg \max_{x,y} a_{js}^\top x - b_{js}$$

s.t.  $x \in X$

$$x^\top a_j^0 + \Omega_s \hat{a}_j^\top x - b_{js} \leq M_{js}^k y_s, \quad \forall j, s$$

$$\sum_{s \in \mathcal{S}} y_s \leq p$$

$$0 \leq y_s \leq 1, \quad \forall s.$$

**end if**

**if**  $M_{js}^{k+1} < 0$  **then**

            2) Screening phase: Eliminate constraint  $(j, s)$  from the model.

**end if**

**end for**

    Set  $k \leftarrow k + 1$ .

**end while**

---

## Valid inequalities


---

## Valid inequalities I: generators

$$p_g - \Omega_s \beta_g \leq \bar{p}_g \quad \Rightarrow \quad L_{js} : f_{js}(z_j) = \Omega_s z_j + b_j \ (\leq 0)$$

# Valid inequalities I: generators


$$\rho_g - \Omega_s \beta_g \leq \bar{\rho}_g \quad \Rightarrow \quad L_{js} : f_{js}(z_j) = \Omega_s z_j + b_j (\leq 0)$$






# Valid inequalities I: generators

$$\rho_g - \Omega_s \beta_g \leq \bar{\rho}_g \quad \Rightarrow \quad L_{js} : f_{js}(z_j) = \Omega_s z_j + b_j (\leq 0)$$




## Valid inequalities II: lines

$$L_{js} : f_{js}(z_j) = \Omega_s z_j + b_{js}$$




## Valid inequalities II: lines

$$L_{js} : f_{js}(z_j) = \Omega_s z_j + b_{js}$$



# Valid inequalities II: lines

$$L_{js} : f_{js}(z_j) = \Omega_s z_j + b_{js}$$



## Proposition


For a fixed  $j \in \mathcal{J}$ , let  $U_j^{p+1}(\cdot)$  be the  $(p+1)$ -upper envelope of the set of lines  $\mathcal{L}_j$ , with  $p := \lfloor \epsilon |\mathcal{S}| \rfloor$ . Then the inequality

$$U_j^{p+1}(\hat{a}_j^\top x) + x^\top a_j^0 \leq 0, \quad x \in X$$

is valid.

# Valid inequalities II: lines


$$L_{js} : f_{js}(z_j) = \Omega_s z_j + b_{js}$$



Rider Algorithm


## Valid inequalities II: lines

$$L_{js} : f_{js}(z_j) = \Omega_s z_j + b_{js}$$




## Valid inequalities II: lines

$$L_{js} : f_{js}(z_j) = \Omega_s z_j + b_{js}$$



## Valid inequalities II: lines

$$L_{js} : f_{js}(z_j) = \Omega_s z_j + b_{js}$$




Lower hull: *Jarvis March, Graham scan*




## Valid inequalities II: lines

$$L_{js} : f_{js}(z_j) = \Omega_s z_j + b_{js}$$



## Valid inequalities II: lines

$$L_{js} : f_{js}(z_j) = \Omega_s z_j + b_{js}$$



- Relationship with *quantile cuts*.

## Computational Results: OPF

---

## Computational Results: OPF

- Approaches **T**, **TS**, **V** and **TS+V** using five standard power systems.
- Instance: IEEE-300 test system: 300 nodes, 57 generators, 411 lines.
- GUROBI 9.1.2 on a Linux-based server with CPUs clocking at 2.6 GHz, 6 threads and 32 GB of RAM.
- 1000 scenarios, 5% violation of the JCC ( $\epsilon = 0.05$ ,  $p = 50$ ).
- Time limit: 10 hours.
- Results averaged over ten instances.

# Computational Results

| IEEE-300 | BN        | T(3)      | TS(3)     | BN+V      | TS(1)+V    |
|----------|-----------|-----------|-----------|-----------|------------|
| #CON     | 936939    | 100%      | 8.0%      | 101.5%    | 3.30%      |
| LRgap    | 1.114%    | 0.264%    | 0.264%    | 0.3192%   | 0.1603%    |
| MIPgap   | 0.27% (0) | 0.07% (0) | 0.01% (4) | 0.08% (0) | 0.00% (10) |
| Time     | 36000     | 1.0x      | 1.2x      | 1.0x      | 8.5x       |

Á. Porras, C. Domínguez, J.M. Morales, and S. Pineda. (2022) Tight and compact sample average approximation for joint chance-constrained optimal power flow. *arXiv preprint arXiv:2205.03370*.

W. Xie, and S. Ahmed. (2018) On quantile cuts and their closure for chance constrained optimization problems. *Mathematical Programming*, **172**, 621–646.

M. Cheema, Z. Shen, X. Lin, and W. Zhang. (2014). A unified framework for efficiently processing ranking related queries. In *EDBT*, 427–438.

THANK YOU FOR YOUR ATTENTION