

Efficiently solving linear bilevel programming problems using off-the-shelf optimization software

EURO 2019

S. Pineda J. M. Morales

University of Malaga (Spain)

June 26, 2019

Bilevel programming

- Decentralized environments: multiple decisions maker with divergent objectives that interact with each other in a hierarchical organization
- Simplest case: one leader and one follower (Stackelberg game)
- A Stackelberg game can be mathematically formulated as a bilevel problem (BLP)

$$\min_x F(x, y) \quad (1a)$$

$$\text{s.t. } G_i(x, y) \geq 0, \quad \forall i \quad (1b)$$

$$\min_y f(x, y) \quad (1c)$$

$$\text{s.t. } g_j(x, y) \geq 0, \quad \forall j \quad (1d)$$

- Even if $F(x, y)$, $f(x, y)$, $G_i(x, y)$ and $g_j(x, y)$ are linear, the BLP is proven to be NP-hard¹

¹Jeroslow 1985; Bard 1991.

Bilevel programming applications

BLP is widely used in energy and power system applications:

- Electricity grid security analysis²
- Transmission expansion planning³
- Strategic bidding of power producers⁴
- Generation capacity expansion⁵
- Investment in wind power generation⁶
- Market equilibria models⁷

²Motto, Arroyo, and Galiana 2005.

³Garces et al. 2009; Jenabi, Fatemi Ghomi, and Smeers 2013.

⁴Ruiz and Conejo 2009; Zugno et al. 2013.

⁵Wogrin, Centeno, and Barquín 2011; Kazempour et al. 2011.

⁶Baringo and Conejo 2014; Maurovich-Horvat et al. 2015.

⁷Pozo and Contreras 2011; Ruiz, Conejo, and Smeers 2012.

Methods to solve bilevel programming

Methods to solve BLP can be divided into two main categories:

- Dedicated methods⁸
 - Efficient and globally optimal
 - Hard to implement in commercial optimization software
- Plug-and-play methods⁹
 - Straightforward implementation in commercial optimization software
 - High computational burden and locally optimal
 - Most common: reformulate as single-level using Fortuny-Amat (bigM)

⁸Bialas and Karwan 1984; Shi, Lu, and Zhang 2005; Calvete, Galé, and Mateo 2008; Li and Fang 2012; Sinha, Malo, and Deb 2013; Jiang et al. 2013; Bard and Falk 1982; Bard and Moore 1990; Hansen, Jaumard, and Savard 1992; Shi et al. 2006.

⁹Fortuny-Amat and McCarl 1981; Ruiz and Conejo 2009; Gabriel and Leuthold 2010; Siddiqui and Gabriel 2012; Scholtes 2001; Ralph and Wright 2004; White and Anandalingam 1993; Hu and Ralph 2004; Lv et al. 2007; Fletcher and Leyffer 2004;

Linear bilevel problem

If all functions are linear, the resulting linear bilevel problem (LBLP) can be generally formulated as

$$\min_x \quad c_1x + d_1y \quad (2a)$$

$$\text{s.t.} \quad A_1x + B_1y \leq b_1 \quad (2b)$$

$$\min_y \quad c_2x + d_2y \quad (2c)$$

$$\text{s.t.} \quad A_2x + B_2y \leq b_2 \quad (\lambda) \quad (2d)$$

We assume $B_1 = 0$ to avoid disconnected feasible regions¹⁰

¹⁰Colson, Marcotte, and Savard 2005; Shi, Zhang, and Lu 2005.

Linear bilevel problem

Since the lower-level optimization problem is linear, it can be replaced with its KKT optimality conditions

$$\min_{x,y,\lambda} \quad c_1x + d_1y \quad (3a)$$

$$\text{s.t.} \quad A_1x + B_1y \leq b_1 \quad (3b)$$

$$d_2 + \lambda B_2 = 0 \quad (3c)$$

$$b_2 - A_2x - B_2y \geq 0 \quad (3d)$$

$$\lambda \geq 0 \quad (3e)$$

$$\lambda (b_2 - A_2x - B_2y) = 0 \quad (3f)$$

Without complementarity conditions (3f), problem (3) would be linear. Thus, all methods differ on how to deal with this constraints.

Plug-and-play methods

- Special order sets type 1 (SOS1)
- Fortuny-Amat with bigM (FA)
- Regularization (REG)
- Proposed method (REG-FA)

- This method declares Special Ordered Sets of type 1 (SOS1)¹¹

$$s_j(1) = (b_2 - A_2x - B_2y)_j, \quad \forall j$$

$$s_j(2) = \lambda_j, \quad \forall j$$

- This method explores using a binary tree all combinations of the complementarity constraints and therefore ensures global optimality

¹¹Siddiqui and Gabriel 2012.

The complementarity constraints are reformulated as

$$\min_{x,y,\lambda,u} \quad F(x,y) = c_1x + d_1y \quad (5a)$$

$$\text{s.t.} \quad A_1x + B_1y \leq b_1 \quad (5b)$$

$$d_2 + \lambda B_2 = 0 \quad (5c)$$

$$b_2 - A_2x - B_2y \geq 0 \quad (5d)$$

$$\lambda \geq 0 \quad (5e)$$

$$b_2 - A_2x - B_2y \leq (1 - u)M_1 \quad (5f)$$

$$\lambda \leq uM_2 \quad (5g)$$

$$u \in \{0, 1\} \quad (5h)$$

where u is a vector of binary variables of appropriate size and M_1, M_2 are large enough scalars that need to be adjusted.

Fortuny-Amat with bigM

BigM are usually adjusted by the following trial-and-error procedure:

- 1 Select initial values for M_1 and M_2 .
- 2 Solve model (5) using MIP solver (CPLEX).
- 3 Find a j' such that $u_{j'} = 0$ and $(b_2 - A_2x - B_2y)_{j'} = M_{1j'}$. If such a j' exists, increase the value of $M_{1j'}$ and go to step 2). Otherwise, go to step 4).
- 4 Find a j' such that $u_{j'} = 1$ and $\lambda_{j'} = M_{2j'}$. If such a j' exists, increase the value of $M_{2j'}$ and go to step 2). Else, the solution to (2) *is assumed* to correspond to the optimal solution of the original bilevel problem (1).

This method may fail and provide highly suboptimal solutions!!

Let us consider the following linear bilevel problem:

$$\begin{aligned} \max_{x \in \mathbb{R}} \quad & z = x + y \\ \text{s.t.} \quad & 0 \leq x \leq 2 \\ & \min_{y \in \mathbb{R}} \quad y \\ & \text{s.t.} \quad y \geq 0 \quad (\lambda_1) \\ & \quad \quad x - 0.01y \leq 1 \quad (\lambda_2) \end{aligned}$$

It is easy to verify that the optimal solution to this problem is $z^* = 102, x^* = 2, y^* = 100, \lambda_1^* = 0, \lambda_2^* = 100$.

Fortuny-Amat with bigM

We can reformulate it using bigM as follows:

$$\max_{x,y} \quad z = x + y$$

$$\text{s.t.} \quad 0 \leq x \leq 2 \quad x - 0.01y \leq 1$$

$$1 - \lambda_1 - 0.01\lambda_2 = 0$$

$$y, \lambda_1, \lambda_2 \geq 0$$

$$\lambda_1 \leq u_1 M_1^D$$

$$y \leq (1 - u_1) M_1^P$$

$$\lambda_2 \leq u_2 M_2^D$$

$$-x + 0.01y + 1 \leq (1 - u_2) M_2^P$$

$$u_1, u_2 \in \{0, 1\}$$

$$\text{For } M_{1,2}^P = 200 \quad M_{1,2}^D = 50$$

Case	u_1	u_2	x	y	λ_1	λ_2	z
1	0	1	Infeasible				
2	1	1	1	0	Multiple ^(*)		1
3	1	0	1	0	1	0	1
4	0	0	Infeasible				

$$(*) \quad 0 \leq \lambda_1 \leq 50, 0 \leq \lambda_2 \leq 50, 1 - \lambda_1 - 0.01\lambda_2 = 0$$

- Case 2 includes $\lambda_1 = 0.5, \lambda_2 = 50$
- CPLEX always provides Case 3
- Since $\lambda_1 < 50, \lambda_2 < 50$, Case 3 is assumed to be global optimal!!

Adjusting big-M by trial-and-error may lead to suboptimal solutions¹²

¹²Pineda and Morales 2019.

Hanging out with mathematicians

- Good news: you can easily obtain valid bounds on λ_2 by formulating the dual of the lower-level problem

$$\begin{aligned} \max_{\lambda_2} \quad & \lambda_2(x - 1) \\ \text{s.t.} \quad & 0 \leq \lambda_2 \leq 100 \end{aligned}$$

- Bad news: in general, finding valid bounds for dual variables cannot be done in polynomial time unless $P = NP$ ¹³

¹³Kleinert et al. 2019.

Regularization approach

All feasible points of (3) are nonregular (nonlinear solvers fail even to find a local optimal solution). This problem can be regularized as follows¹⁴:

$$\min_{x,y,\lambda} \quad F(x,y) = c_1x + d_1y \quad (9a)$$

$$\text{s.t.} \quad A_1x + B_1y \leq b_1 \quad (9b)$$

$$d_2 + \lambda B_2 = 0 \quad (9c)$$

$$b_2 - A_2x - B_2y \geq 0 \quad (9d)$$

$$\lambda \geq 0 \quad (9e)$$

$$\lambda (b_2 - A_2x - B_2y) \leq t \quad (9f)$$

- Parameter t is iteratively decreased to 0
- Strong theoretical and empirical convergence properties
- Only guaranteed to provide local optimal solutions¹⁵

¹⁴Scholtes 2001; Ralph and Wright 2004.

¹⁵Dempe 2019.

Proposed approach

- The regularization method is fast, but only provides local optimal solutions
- The bigM method achieves global optimality provided that large constants are set to proper values
- The proposed method aims to combine both approaches. We propose to use information about the local optimal solution to set the **large constants** and find initial values of the **binary variables** as follows:
 - 1 Solve (3) using regularization to obtain a local optimal solution
 - 2 Select a value of $\mathcal{M} > 1$
 - 3 Set $M_1 \leftarrow \mathcal{M} \max_j \{(b_2 - A_2x - B_2y)_j\}$ and $M_2 \leftarrow \mathcal{M} \max_j \{(\lambda)_j\}$
 - 4 Set initial values of binary variables u as follows. If $(b_2 - A_2x - B_2y)_j > 0$, then $u_j = 0$. If $\lambda_j > 0$, then $u_j = 1$
 - 5 Solve mixed-integer problem (5)

Computational results

We compare the proposed method with existing ones using 300 randomly generated examples of different sizes:

$$\min_x \quad c_1x + d_1y$$

$$\text{s.t.} \quad A_1x + B_1y \leq b_1$$

$$\min_y \quad c_2x + d_2y$$

$$\text{s.t.} \quad A_2x + B_2y \leq b_2$$

	n	m	p	q	r
Small	50	50	25	25	25
Medium	100	100	50	50	50
Large	200	200	100	100	100

$$\begin{aligned} c_1 &= |\mathcal{N}(1, n)| \\ d_1 &= |\mathcal{N}(1, m)| \end{aligned} \quad A_1 = \begin{pmatrix} \mathcal{N}(p, n) \\ -I \end{pmatrix} \quad B_1 = \begin{pmatrix} \mathbf{0} \\ \mathbf{0} \end{pmatrix} \quad b_1 = \begin{pmatrix} \mathcal{N}(p, 1) \\ \mathbf{0} \end{pmatrix}$$

$$\begin{aligned} c_2 &= |\mathcal{N}(1, n)| \\ d_2 &= |\mathcal{N}(1, m)| \end{aligned} \quad A_2 = \begin{pmatrix} \mathcal{N}(q, n) \\ \mathbf{0} \\ \mathbf{0} \end{pmatrix} \quad B_2 = \begin{pmatrix} \mathcal{N}(q, m) \\ \mathcal{N}(r, m) \\ -I \end{pmatrix} \quad b_2 = \begin{pmatrix} \mathcal{N}(q, 1) \\ \mathcal{N}(r, 1) \\ \mathbf{0} \end{pmatrix}$$

Computational results

	Small (n=50)				Medium (n=100)				Large (n=200)			
	#opt	#inf	time (s)	gap(%)	#opt	#inf	time(s)	gap(%)	#opt	#inf	time(s)	gap(%)
SOS1	98	2	1	0.00	90	0	4656	0.27	27	0	17652	2.05
FA-5	11	24	12	7.65	8	7	5385	7.15	0	20	21415	4.57
FA-50	98	2	5	0.00	94	2	5495	0.04	53	1	16873	0.31
FA-100000	27	2	0	6.44	11	0	0	10.19	2	0	1	9.86
REG	61	2	0	0.55	41	0	1	0.52	30	1	14	0.29
REG-FA-10	98	2	5	0.00	99	0	2353	0.00	72	0	13258	0.10

- #opt: Number of problems solved to optimality (out of 100)
- #inf: Number of infeasible problems (out of 100)
- time: average time of 100 problems
- gap: average gap with respect to global optimal solution of 100 problems

Computational results

	Small (n=50)				Medium (n=100)				Large (n=200)			
	#opt	#inf	time (s)	gap(%)	#opt	#inf	time(s)	gap(%)	#opt	#inf	time(s)	gap(%)
SOS1	98	2	1	0.00	90	0	4656	0.27	27	0	17652	2.05
FA-5	11	24	12	7.65	8	7	5385	7.15	0	20	21415	4.57
FA-50	98	2	5	0.00	94	2	5495	0.04	53	1	16873	0.31
FA-100000	27	2	0	6.44	11	0	0	10.19	2	0	1	9.86
REG	61	2	0	0.55	41	0	1	0.52	30	1	14	0.29
REG-FA-10	98	2	5	0.00	99	0	2353	0.00	72	0	13258	0.10

- SOS1 works fine for small size problems
- For large problems, SOS1 reach the maximum time of 6 h

Computational results

	Small (n=50)				Medium (n=100)				Large (n=200)			
	#opt	#inf	time (s)	gap(%)	#opt	#inf	time(s)	gap(%)	#opt	#inf	time(s)	gap(%)
SOS1	98	2	1	0.00	90	0	4656	0.27	27	0	17652	2.05
FA-5	11	24	12	7.65	8	7	5385	7.15	0	20	21415	4.57
FA-50	98	2	5	0.00	94	2	5495	0.04	53	1	16873	0.31
FA-100000	27	2	0	6.44	11	0	0	10.19	2	0	1	9.86
REG	61	2	0	0.55	41	0	1	0.52	30	1	14	0.29
REG-FA-10	98	2	5	0.00	99	0	2353	0.00	72	0	13258	0.10

- FA-5 leads to infeasible problems since bigM are not large enough
- Numerical instabilities occur for FA-100000
- FA-50 provides the best results for this approach

Computational results

	Small (n=50)				Medium (n=100)				Large (n=200)			
	#opt	#inf	time (s)	gap(%)	#opt	#inf	time(s)	gap(%)	#opt	#inf	time(s)	gap(%)
SOS1	98	2	1	0.00	90	0	4656	0.27	27	0	17652	2.05
FA-5	11	24	12	7.65	8	7	5385	7.15	0	20	21415	4.57
FA-50	98	2	5	0.00	94	2	5495	0.04	53	1	16873	0.31
FA-100000	27	2	0	6.44	11	0	0	10.19	2	0	1	9.86
REG	61	2	0	0.55	41	0	1	0.52	30	1	14	0.29
REG-FA-10	98	2	5	0.00	99	0	2353	0.00	72	0	13258	0.10

- The computational time for FA approach dramatically increases with problem size

Computational results

	Small (n=50)				Medium (n=100)				Large (n=200)			
	#opt	#inf	time (s)	gap(%)	#opt	#inf	time(s)	gap(%)	#opt	#inf	time(s)	gap(%)
SOS1	98	2	1	0.00	90	0	4656	0.27	27	0	17652	2.05
FA-5	11	24	12	7.65	8	7	5385	7.15	0	20	21415	4.57
FA-50	98	2	5	0.00	94	2	5495	0.04	53	1	16873	0.31
FA-100000	27	2	0	6.44	11	0	0	10.19	2	0	1	9.86
REG	61	2	0	0.55	41	0	1	0.52	30	1	14	0.29
REG-FA-10	98	2	5	0.00	99	0	2353	0.00	72	0	13258	0.10

- Local optimal solutions are quite close to the global optimal solutions
- The size of the problem does not significantly affect computational time

Computational results

	Small (n=50)				Medium (n=100)				Large (n=200)			
	#opt	#inf	time (s)	gap(%)	#opt	#inf	time(s)	gap(%)	#opt	#inf	time(s)	gap(%)
SOS1	98	2	1	0.00	90	0	4656	0.27	27	0	17652	2.05
FA-5	11	24	12	7.65	8	7	5385	7.15	0	20	21415	4.57
FA-50	98	2	5	0.00	94	2	5495	0.04	53	1	16873	0.31
FA-100000	27	2	0	6.44	11	0	0	10.19	2	0	1	9.86
REG	61	2	0	0.55	41	0	1	0.52	30	1	14	0.29
REG-FA-10	98	2	5	0.00	99	0	2353	0.00	72	0	13258	0.10

- The proposed approach achieves the optimal solution in most problems and achieves the lower average gap at reasonable computational times

Computational results

For $n = 100$, we change the scaling and sparsity of matrices and vectors

	Full matrix, good scaled				Sparse matrix, good scaled				Full matrix, bad scaled			
	#opt	#inf	time (s)	gap(%)	#opt	#inf	time(s)	gap(%)	#opt	#inf	time(s)	gap(%)
SOS1	90	0	4656	0.27	86	0	4293	0.48	56	0	18419	7.02
FA-5	8	7	5385	7.15	7	12	4370	8.75	0	100	-	-
FA-50	94	2	5495	0.04	92	2	4283	0.02	0	100	-	-
FA-100000	11	0	0	10.19	10	0	0	10.58	3	0	1	202.40
REG	41	0	1	0.52	45	0	1	0.67	4	41	4	6.68
REG-FA-10	99	0	2353	0.00	97	1	1644	0.01	82	6	10702	0.16

- Having sparse matrices do not significantly affect the comparison
- FA works particularly bad for bad scaled problems for any value of bigM
- The proposed method outperforms existing ones in bad scaled problems

Conclusions

If you are solving a linear bilevel problem you have the following options:

- Dedicated methods: efficient but hard to code
- SOS1: global optimal, but computational time is extremely high
- REG: fast but only provides local optimal solutions
- FA with bigM: easy to implement, but setting bigM with trial-and-error method may provide suboptimal solutions
- Try to find better ways to set large constants as the one we propose

Thanks!! Any Questions?



website: oasys.uma.es

S. Pineda, H. Bylling and J. M. Morales,
“Efficiently solving linear bilevel programming
problems using off-the-shelf optimization
software”, in **Optimization and Engineering**,
19(1), 187-211, 2018.



S. Pineda and J. M. Morales, “Solving Linear
Bilevel Problems Using Big-Ms: Not All That
Glitters Is Gold”, in **IEEE Transactions on
Power Systems**, 34(3), 2469-2471, 2019.



References I

- Bard, J and J Moore (1990). "A Branch and Bound Algorithm for the Bilevel Programming Problem". In: *SIAM Journal on Scientific and Statistical Computing* 11.2, pp. 281–292.
- Bard, J. F. (1991). "Some properties of the bilevel programming problem". In: *Journal of Optimization Theory and Applications* 68.2, pp. 371–378.
- Bard, Jonathan F. and James E. Falk (1982). "An explicit solution to the multi-level programming problem". In: *Computers & Operations Research* 9.1, pp. 77–100.
- Baringo, Luis and Antonio J. Conejo (2014). "Strategic Wind Power Investment". In: *IEEE Trans. Power Syst.* 29.3, pp. 1250–1260.
- Bialas, W. F. and M. H. Karwan (1984). *Two-Level Linear Programming*.
- Calvete, Herminia I., Carmen Galé, and Pedro M. Mateo (2008). "A new approach for solving linear bilevel problems using genetic algorithms". In: *European Journal of Operational Research* 188.1, pp. 14–28.

References II

- Colson, Benoît, Patrice Marcotte, and Gilles Savard (2005). “Bilevel programming: A survey”. In: *4OR* 3.2, pp. 87–107.
- Dempe, Stephan (2019). “Computing Locally Optimal Solutions of the Bilevel Optimization Problem Using the KKT Approach”. In: Springer, Cham, pp. 147–157.
- Fletcher, Roger and Sven Leyffer (2004). “Solving mathematical programs with complementarity constraints as nonlinear programs”. In: *Optimization Methods and Software* 19.1, pp. 15–40.
- Fortuny-Amat, José and Bruce McCarl (1981). “A Representation and Economic Interpretation of a Two-Level Programming Problem”. In: *Journal of the Operational Research Society* 32.9, pp. 783–792.
- Gabriel, Steven A. and Florian U. Leuthold (2010). “Solving discretely-constrained MPEC problems with applications in electric power markets”. In: *Energy Economics* 32.1, pp. 3–14.

References III

- Garces, L.P. P et al. (2009). “A Bilevel Approach to Transmission Expansion Planning Within a Market Environment”. In: *IEEE Trans. Power Syst.* 24.3, pp. 1513–1522.
- Hansen, P., B. Jaumard, and G. Savard (1992). “New branch-and-bound rules for linear bilevel programming”. eng. In: *SIAM journal on scientific and statistical computing* 13.5, pp. 1194–1217.
- Hu, X. M. and D. Ralph (2004). “Convergence of a Penalty Method for Mathematical Programming with Complementarity Constraints”. In: *Journal of Optimization Theory and Applications* 123.2, pp. 365–390.
- Jenabi, Masoud, Seyyed Mohammad Taghi Fatemi Ghomi, and Yves Smeers (2013). “Bi-Level Game Approaches for Coordination of Generation and Transmission Expansion Planning Within a Market Environment”. In: *IEEE Trans. Power Syst.* 28.3, pp. 2639–2650.
- Jeroslow, Robert G. (1985). “The polynomial hierarchy and a simple model for competitive analysis”. In: *Mathematical Programming* 32.2, pp. 146–164.

References IV

- Jiang, Yan et al. (2013). "Application of particle swarm optimization based on CHKS smoothing function for solving nonlinear bilevel programming problem". In: *Applied Mathematics and Computation* 219.9, pp. 4332–4339.
- Kazempour, S. Jalal et al. (2011). "Strategic Generation Investment Using a Complementarity Approach". In: *IEEE Trans. Power Syst.* 26.2, pp. 940–948.
- Kleinert, Thomas et al. (2019). "There ' s No Free Lunch : On the Hardness of Choosing a Correct Big-M in Bilevel Optimization". In: pp. 11–13.
- Li, Hecheng and Lei Fang (2012). "An evolutionary algorithm for solving bilevel programming problems using duality conditions". In: *Mathematical Problems in Engineering* 2012.
- Lv, Yibing et al. (2007). "A penalty function method based on Kuhn-Tucker condition for solving linear bilevel programming". In: *Applied Mathematics and Computation* 188.1, pp. 808–813.

References V

- Maurovich-Horvat, Lajos et al. (2015). “Transmission and Wind Investment in a Deregulated Electricity Industry”. In: *IEEE Trans. Power Syst.* 30.3, pp. 1633–1643.
- Motto, A.L. L L, J.M. M M Arroyo, and F.D. D D Galiana (2005). “A Mixed-Integer LP Procedure for the Analysis of Electric Grid Security Under Disruptive Threat”. In: *IEEE Trans. Power Syst.* 20.3, pp. 1357–1365.
- Pineda, Salvador and Juan Miguel Morales (2019). “Solving linear bilevel problems using big-ms: not all that glitters is gold”. In: *IEEE Transactions on Power Systems* 34.3, pp. 2469–2471.
- Pozo, David and Javier Contreras (2011). “Finding Multiple Nash Equilibria in Pool-Based Markets: A Stochastic EPEC Approach”. In: *IEEE Trans. Power Syst.* 26.3, pp. 1744–1752.
- Ralph, Daniel and Stephen J. Wright (2004). “Some properties of regularization and penalization schemes for MPECs”. en. In: *Optimization Methods and Software* 19.5, pp. 527–556.

References VI

- Ruiz, C. and A.J. J J Conejo (2009). “Pool Strategy of a Producer With Endogenous Formation of Locational Marginal Prices”. In: *IEEE Trans. Power Syst.* 24.4, pp. 1855–1866.
- Ruiz, Carlos, Antonio J. Conejo, and Yves Smeers (2012). “Equilibria in an Oligopolistic Electricity Pool With Stepwise Offer Curves”. In: *IEEE Trans. Power Syst.* 27.2, pp. 752–761.
- Scholtes, Stefan (2001). “Convergence Properties of a Regularization Scheme for Mathematical Programs with Complementarity Constraints”. en. In: *SIAM Journal on Optimization* 11.4, pp. 918–936.
- Shi, Chenggen, Jie Lu, and Guangquan Zhang (2005). “An extended Kth-best approach for linear bilevel programming”. In: *Applied Mathematics and Computation* 164.3, pp. 843–855.
- Shi, Chenggen, Guangquan Zhang, and Jie Lu (2005). “On the definition of linear bilevel programming solution”. In: *Applied Mathematics and Computation* 160.1, pp. 169–176.

References VII

- Shi, Chenggen et al. (2006). “An extended branch and bound algorithm for linear bilevel programming”. In: *Applied Mathematics and Computation* 180.2, pp. 529–537.
- Siddiqui, S. and S. A. Gabriel (2012). “An SOS1-Based Approach for Solving MPECs with a Natural Gas Market Application”. In: *Networks and Spatial Economics* 13.2, pp. 205–227.
- Sinha, Ankur, Pekka Malo, and Kalyanmoy Deb (2013). “Efficient Evolutionary Algorithm for Single-Objective Bilevel Optimization”. In: *arXiv* 1303.3901.
- White, D. J. and G. Anandalingam (1993). “A penalty function approach for solving bi-level linear programs”. In: *Journal of Global Optimization* 3.4, pp. 397–419.
- Wogrin, Sonja, Efraim Centeno, and Julián Barquín (2011). “Generation capacity expansion in liberalized electricity markets: A stochastic MPEC approach”. In: *IEEE Trans. Power Syst.* 26.4, pp. 2526–2532.

Zugno, Marco et al. (2013). “Pool Strategy of a Price-Maker Wind Power Producer”. In: *IEEE Trans. Power Syst.* 28.3, pp. 3440–3450.