

Using Bézier extraction matrices

Péter Salvi

November 18, 2021

Given a B-spline curve of degree d with L segments in the form

$$\mathcal{C}(u) = \sum_{k=0}^{L+d-1} \mathbf{P}_k N_k^{d,U}(u),$$

where U is the clamped knot vector with $d+1$ identical values at both ends, and $N_k^{d,U}$ are the B-spline basis functions, the i -th Bézier component ($i = 0..L-1$) can be extracted as

$$\hat{\mathcal{C}}_i(u) = \sum_{j=0}^d \mathbf{Q}_j^i B_j^d(u),$$

where B_j^d are the Bernstein polynomials and

$$\mathbf{Q}_j^i = \sum_{k=0}^d \mathbf{P}_{k+s_i-d} C_{k,j}^i,$$

because of the extraction relation

$$N_k^{d,U}(u) = \sum_{j=0}^d C_{k+d-s_i,j}^i B_j^d(u),$$

\mathbf{C}^i being the i -th extraction matrix of the degree- d knot vector U , and s_i denoting the span index of the i -th segment.

Conversely, a B-spline control point \mathbf{P}_k can be restored from a suitable Bézier segment (i.e., one that is in the same span), using the formula

$$\mathbf{P}_k = \sum_{j=0}^d \mathbf{Q}_j^i \left(\mathbf{C}^{i-1} \right)_{j,k+d-s_i}.$$