Studies on interference effects of flavor changing neutral currents in the top sector ($tu\gamma$, $tc\gamma$ couplings)

Salvatore La Cagnina

Technische Universität Dortmund, Lehrstuhl EIV

October 24, 2017




Productions:

- Production mode sample:
- generate p p > t a \$\$ t \sim , add p p > t u a \$\$ t + anti-particle processes
- Decay mode sample:
- generate p p > t t \sim , t \sim > a u \sim (will be changed to p p > t t \sim > t a u \sim due to MadSpin errors)
- Interference Sample:
- generate p p > t a, add p p > t u \sim a + anti-particle processes




Creation of following plots:


- TopFCNC Model in MadGraph@NLO
- 50k Events for each generated sample
- Showering using Pythia 8
- DELPHES with ATLAS card as basic detector simulation
- Look at bJet, Photon, W Boson and Top Quark p_T , mass, η , ϕ distributions




BQuark_truth




BQuark_truth




BQuark_truth




BQuark_truth interference vs. decay+production




Photon_truth




Photon_truth




Photon_truth




Photon_truth interference vs. decay+production




TopQuark_truth




TopQuark_truth




TopQuark_truth




TopQuark_truth interference vs. decay+production




UQuark_truth




UQuark_truth




UQuark_truth




UQuark_truth interference vs. decay+production




WBoson_truth



WBoson_truth



WBoson_truth



WBoson_truth interference vs. decay+production

