

Clipboard Font Alignment Number Styles

Clipboard: Paste, Copy, Format Painter

Font: Aptos Narrow, 11, A, A, B, I, U, Font Color, Text Color

Alignment: Left, Center, Right, Justify, Merge & Center, Wrap Text

Number: General, Percentage, Decimals, Fractions, More Numbering

Styles: Normal, Good, Bad, Neutral

C8 Exclusions include: panobinostat and elotuzumab. Other potential therapies pending review of data may be considered

A	B	C	D	E	F	G	H	I	J	K	L
1											
2											
3											
4											

INCLUSION CRITERIA

At least one inpatient medical claim with a diagnosis code for MM in any position (any ICD-9-CM=203.0x or ICD-10-CM code=C90.0x) or ≥ 2 outpatient medical claims for MM in any position on the claim, on separate days within 90 days, during the study period

Aged ≥18 years at the time of MM diagnosis according to calendar year

Received an eligible or expected treatment for MM on or after MM diagnosis (other than belantamab), occurring on or after 01 Jan 2017 (eligible treatment period).

Exclusions include: panobinostat and elotuzumab. Other potential therapies pending review of data may be considered

CE of at least 12-months with medical and pharmacy benefits before the 1L cohort index date.

Patients with gaps in enrollment of ≤ 30 days are considered to be continuously enrolled

T1. Baseline demographics, comorbidities and clinical characteristics for NDMM and RRMM cohorts, overall and by LOT¹

Primary Objective / Secondary Baseline

1L (N=)																							2L (N=)						3L (N=)					
Overall		aCD38 Quad	aCD38 Triplet	Other Triplets	Doublet/Mono	Other	Overall	aCD38 Quad	aCD38 Triplet	Parm-based triplets (Non-aCD38)	Other Triplets	Doublet/Mono	BCMA	Bi-specific	Overall	aCD38 Quad	aCD38 Triplet	Parm-based triplets (Non-aCD38)	Other Triplets	Doublet/Mono	BCMA	Bi-specific												
N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=	N=													
Demographic characteristics at MM diagnosis (overall)/ at index																																		
Sex (N%)																																		
Male																																		
Female																																		
Age at Index																																		
Mean and SD																																		
Median and IQR																																		
Min and Max																																		
Age distribution at index (N%)																																		
<75 years																																		
18 to 44 years																																		
45 to 64 years																																		
65 to 74 years																																		
≥ 75 years																																		
Race (N%)																																		
White																																		
Black																																		
Asian																																		
Other																																		
Unknown/missing																																		
Ethnicity (N%)																																		
Hispanic/Latino																																		
Non-Hispanic/Latino																																		
Unknown/missing																																		
Insurance type at index (N%)																																		
Commercial Health Plan																																		
Medicare Advantage																																		
Region (N%)																																		
Midwest																																		
South																																		

Home Insert Draw Page Layout Formulas Data Review View Automate Help

Cut

Copy

Format Painter

Clipboard

Apptos Narrow

Font

Font

Align Left

Align Center

Align Right

Align Justify

General

Number

Conditional Formatting

Format as Table

Normal

Bad

Neutral

Insert

Delete

Format

Autosum

Fill

Clear

Editing

Sort & Find & Filter

Sensitivity

Add-ins

West

Alignment

Number

Conditional Formatting

Format as Table

Normal

Bad

Neutral

Insert

Delete

Format

Autosum

Fill

Clear

Editing

Sort & Find & Filter

Sensitivity

Add-ins

T1. Baseline demographics, comorbidities and clinical characteristics for NDMM and BRMM cohorts, overall and by LOT¹

Primary Objective / Secondary Baseline		1L (N=)										2L (N=)										3L (N=)									
West																															
Northeast																															
Unknown																															
Comorbidities and clinical characteristics during baseline ²																															
CCI score during baseline ³																															
Mean and SD																															
Median and IQR																															
Min and Max																															
CCI categorical score (N%) ⁴																															
0																															
1																															
2																															
3																															
4																															
5+																															
MM diagnosis and follow-up from 1L information																															
Age at diagnosis																															
Mean and SD																															
Median and IQR																															
Min and max																															
Age distribution at diagnosis (N%)																															
<75 years																															
18 to 44 years																															
45 to 64 years																															
65 to 74 years																															
≥ 75 years																															
Year of MM diagnosis (overall) / start of LoI (N%)																															
2015		na										na										na									
2016		na										na										na									
2017		na										na										na									
2018		na										na										na									
2019		na										na										na									
2020		na										na										na									
2021		na										na										na									
2022		na										na										na									
2023		na										na										na									

Cut

Copy

Format Painter

Clipboard

Font

Alignment

Number

Conditional Formatting

Format as Table

Styles

Cells

Editing

Sensitivity

Add-ins

Claude

Apptos Narrow

B I U

Font

Align Left

Align Center

Align Right

Align Justify

Wrap Text

Merge & Center

General

%

Number

Conditional Formatting

Format as Table

Normal

Good

Bad

Neutral

Insert

Delete

Format

Autosum

Fill

Clear

Sort & Filter

Find & Select

Sensitivity

Add-ins

Claude

B71

2022

A

B

C

D

E

F

G

H

I

J

K

L

M

N

O

P

Q

R

S

T

U

V

W

X

T1. Baseline demographics, comorbidities and clinical characteristics for NDMH and RRMH cohorts, overall and by LOT¹

Primary Objective Secondary Baseline

	T1 (N=)					2L (N=)					3L (N=)				
2022															
2023															
2024															
2025															
Total follow up time from diagnosis to each index															
Mean and SD															
Median and IQR															
Min and Max															
Time from diagnosis to T1 initiation (overall) / between each index (months) ⁴															
Mean and SD															
Median and IQR															
Min and Max															

Abbreviations: CCI, Charlson Comorbidity Index; IQR, interquartile range; LOT, line of therapy; SD, standard deviation;

[1] The NDMH cohort is defined as receipt of T1. There are two RRMH cohorts, represented by receipt of 2L and 3L. For each cohort, the index is defined as the initiation of the LOT. Each cohort has a 12-month baseline prior to index.

[2] The index date is the start of each LOT regimen. Patients in 2L could have received 2L+ lines, while 3L represents 3L+ lines.

[3] A 12-month baseline period prior to index was considered for each cohort. Codes to define comorbidities are defined in the appendix.

[4] The CCI was derived according to Quan et al. (2011).

[5] Time between each index= from 1L to 2L (under 1L), from 2L to 3L, and from 3L to T1; corresponds to Lot duration

</

Clipboard Font Alignment Number Styles Cells Editing Sensitivity

Font: 11, Arial, Bold, Italic, Underline, Color, Background Color, Merge & Center, Wrap Text

Number: General, Percentage, Decimals, Fractions, Text, Scientific, Custom

Styles: Normal, Good, Bad, Neutral

Cells: Insert, Delete, Format

Editing: Autosum, Fill, Clear, Sort & Filter, Find & Select

Sensitivity: Add-Ins

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U
38	Seizures																				
39	Infectious, N(%)																				
40	Severe infection resulting in hospitalization																				
41	Other, N(%)																				
42	Thrombocytopenia																				
43	Anemia																				
44	Inpatient hospitalizations																				
45	Number of inpatient hospitalizations by patient																				
46	Mean and SD																				
47	Median and IQR																				
48	Min and Max																				
49	Number of hospitalizations, categorically (N%)																				
50	1 hospitalization																				
51	2 hospitalizations																				
52	3 hospitalizations																				
53	4+ hospitalizations																				
54	Length of stay among patients with ≥1 hospitalization (N=)																				
55	Mean and SD																				
56	Median and IQR																				
57	Min and Max																				
58	Emergency department (ED) visits																				
59	Number of ED visits by patient																				
60	Mean and SD																				
61	Median and IQR																				
62	Min and Max																				
63	Number of hospitalizations, categorically (N%)																				
64	1 hospitalization																				
65	2 hospitalizations																				
66	3 hospitalizations																				
67	4+ hospitalizations																				
68																					
69																					
70																					
71																					
72																					
73																					
74																					
75																					
76																					
77																					

Cut

Copy

Paste

Format Painter

Clipboard

Agios Narrow

11

A

A

B

I

U

+

+

+

+

Font

Font

Alignment

Alignment

General

General

Conditional Formatting

Conditional Formatting

Normal

Bad

Good

Neutral

Insert

Delete

Format

Cells

Cells

Autosum

Autosum

Fill

Fill

Clear

Clear

Editing

Editing

Sensitivity

Sensitivity

Add-ins

Add-ins

Claude

Claude

A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S
Table 2. Real-world treatment outcomes used to proxy treatment effectiveness and tolerability during 2L baseline and treatment periods																		
Secondary objective																		

Background prevalence during 2L baseline										On-treatment incidence during 2L follow-up									
Rates of key safety events																			
Liver/hepatic events																			
Acute hepatitis																			
Abnormal liver function																			
Toxic liver disease																			
Hepatic failure																			
Chronic hepatitis																			
Fibrosis and cirrhosis																			
Non-alcoholic steatohepatitis																			
Renal impairment events																			
Acute kidney injury or acute kidney disease																			
Chronic kidney disease																			
Moderate to severe renal impairment or end stage renal disease																			
Ocular events																			
Corneal ulcer																			
Keratopathies (including ulcerative and infective)																			
Cardiovascular events																			
Myocardial infarction/unstable angina																			
Valvopathy																			
Pulmonary hypertension																			
Cerebrovascular events/stroke and transient ischemic attack (TIA)																			
Peripheral arterial thromboembolism																			
Deep venous thrombosis/pulmonary embolism																			
Peripheral arterial hypertension																			
Neurologic conditions																			

Cut

Copy

Paste

Format Painter

Clipboard

Aptos Narrow

11

A

A

B

I

U

Font

Alignment

General

Conditional Formatting

Normal

Good

Bad

Neutral

Cells

Σ Autosum

Fill

Clear

Editing

Sensitivity

Add-ins

Sensitivity

Add-ins

Comments

Wrap Text

Merge & Center

Number

Format as Table

Insert

Delete

Format

A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S
Peripheral neuropathy																		
Parkinson's disease																		
Cognitive impairment/dementia																		
Other movement disorders																		
Seizures																		
Infectious, N (%)																		
Severe infection resulting in hospitalization																		
Other, N (%)																		
Secondary malignancies																		
Thrombocytopenia																		
Anemia																		
Inpatient hospitalizations																		
Number of inpatient hospitalizations by patient																		
Mean and SD																		
Median and IQR																		
Min and Max																		
Number of hospitalizations, categorically (N%)																		
1 hospitalization																		
2 hospitalizations																		
3 hospitalizations																		
4+ hospitalizations																		
Length of stay among patients with ≥1 hospitalization (N=)																		
Mean and SD																		
Median and IQR																		
Min and Max																		
Emergency department (ED) visits																		
Number of ER visits by patient																		
Mean and SD																		
Median and IQR																		
Min and Max																		
Number of hospitalizations, categorically (N%)																		
1 hospitalization																		
2 hospitalizations																		
3 hospitalizations																		
4+ hospitalizations																		

Cut Copy Paste Format Painter Clipboard

Font: Aptos Narrow, 10, A, A, I, U, Bold, Italic, Underline, Color, Background Color, Merge & Center, Wrap Text


General: Number, Conditional Formatting, Format as Table, Styles: Normal, Good, Bad, Neutral

Cells: Insert, Delete, Format, Fill, Clear

D42

A

T1b. Baseline
Primary Obje



31	Unknown/
32	Insurance ty
33	Commerci
34	Medicare
35	Region (N%)
36	Midwest
37	South
38	West
39	Northeast
40	Unknown
41	Comorbiditie
42	CCI score d
43	Mean and
44	Median ar
45	Min and M
46	CCI categor
47	0
48	1
49	2
50	3
51	4
52	5+
53	MM diagnosis and follow-up from 1L information
54	Age at diagnosis
55	Mean and SD
56	Median and IQR

Cut Copy Paste Format Painter Clipboard

Aptos Narrow 10 A A Font Alignment

General % 00 00 Number

Conditional Formatting Table Normal Good Bad Neutral Styles

Insert Delete Format Cells

D42 fx

T1b. Baseline demographics, comorbidities and clinical characteristics for NDMM and RRM cohorts¹
Primary Objective / Secondary Baseline

	A	B	C	D	E	F	G	H	I	J	K
1											
2											
3											
4											
5											
31											
32											
33											
34											
35											
36											
37											
38											
39											
40											
41											
42											
43											
44											
45											
46											
47											
48											
49											
50											
51											
52											
53											
54											
55											
56											

File Home Insert Draw Page Layout Formulas Data Review View Automate Help

Cut Copy Paste Format Painter Clipboard

Aptos Narrow 10 A A B I U Font Alignment

Wrap Text Merge & Center

General % .00 -00

Conditional Formatting Table

Normal Good Bad Neutral

Insert Delete Format

Autosum Fill Clear

T1b. Baseline demographics, comorbidities and clinical characteristics for NDMM and RRM cohorts¹

Primary Objective / Secondary Baseline

		1L (N=)			2L (N=)			3L (N=)		
Min and max										
Age distribution at diagnosis (N%)										
< 75 years										
18 to 44 years										
45 to 64 years										
65 to 74 years										
≥ 75 years										
Year of MM diagnosis (overall) / start of LoT (N%)										
2016		na	na	na	na	na	na	na	na	na
2017										
2018										
2019										
2020										
2021										
2022										
2023										
2024										
2025										
Total follow up time from diagnosis to each index										
Mean and SD										
Median and IQR										
Min and Max										
Time from diagnosis to 1L initiation (overall) / between each index (months) ⁴										
Mean and SD										
Median and IQR										
Min and Max										

T1b. Baseline demographics, comorbidities and clinical characteristics for NDMM and RMM cohorts ¹			
Primary Objective / Secondary Baseline	1L (N=)	2L (N=)	3L (N=)
Abbreviations: CCI, Charlson Comorbidity Index; IQR, interquartile range; LOT, line of therapy; SD, standard deviation;			
[1] The NDMM cohort is defined as receipt of 1L. There are two RMM cohorts, represented by receipt of 2L and 3L. For each cohort, the index is defined as the initiation of the LOT. Each cohort has a 12-month			
[2] The index date is the start of each LOT regimen. Patients in 2L could have received 2L+ lines, while 3L represents 3L+ lines.			
[3] A 12-month baseline period prior to index was considered for each cohort. Codes to define comorbidities are defined in the appendix.			
[4] The CCI was derived according to Quan et al. (2011).			
[5] Time between each index= from 1L to 2L (under '1L'), from 2L to 3L and from 3L to end; corresponds to Lot duration			

Clipboard Font Alignment Number Styles Cells

Copy Paste Format Painter

Aptos Narrow 11 A A B I U A A Merge & Center

General % 0.00

Conditional Formatting Table

Normal Good Bad Neutral

Insert Delete Format

022

∨

:

×

∧

f_x

∨

A

B

C

D

E

F

G

H

I

J

K

L

M

N

O

Table 5c. Real-world treatment outcomes used to proxy treatment effectiveness and tolerability, by age, neuropathy, and frailty (1L and 2L only)

Secondary objective													
Time-to-next treatment (rwTNT)													
rwTNT from index date													
Events, n (%)													
Censored, n (%)													
Median, months (95% CI)													
Survival probability estimates, 95% CI													
3 months													
6 months													
9 months													
12 months													
18 months													
24 months													
Time to treatment discontinuation (rwTTD)													
rwTTD from index date													
Events, n (%)													
Censored, n (%)													
Median, months (95% CI)													
Survival probability estimates, 95% CI													
3 months													
6 months													
9 months													
12 months													
18 months													
24 months													
Overall survival (rwOS)													
rwTNT from index date													

Cut

Copy

Paste

Clipboard

Format Painter

Aptos Narrow

11

A

A

B

I

U

A

Font

≡

≡

≡

≡

≡

≡

2b

Wrap Text

Merge & Center

General

%

4.00

00

Conditional Formatting

Format as Table

Normal

Good

Bad

Neutral

Insert

Delete

Format

AutoSum

Fill

Clear

D57

fx

A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
	Events, n (%)													
	Censored, n (%)													
	Median, months (95% CI)													
	Survival probability estimates, 95% CI													
	3 months													
	6 months													
	9 months													
	12 months													
	18 months													
	24 months													

Ready

Figures>>

F1. Treatment Sankey

F2. Attrition

Subgroups>>

T1b. Baseline by neuropathy

T5b. Tx Outcomes by subg

EXTRA

T

...

+

FileHomeInsertDrawPage LayoutFormulasDataReviewViewAutomateHelp

CutCopyFormat PainterClipboard

Aptos Narrow11A^A^

IUN

Font

Alignment

General

Conditional Formatting

NormalBadNeutral

InsertDeleteFormat

Cells

U22

N (%)

A

B

U

V

W

X

Y

Z

AA

AB

AC

AD

AE

AF

AG

T2. Treatments received in each line, attrition, and common sequences

Secondary Objective

	LOT Stratified by Age				LOT Stratified by Baseline Neuropathy				LOT Stratified by Baseline Frailty			
	1L	≥75 years	<75 years	≥75 years	1L	No	Yes	No	1L	No	Yes	No
General information												
Patients eligible ¹												
Patients attrition ¹												
Treatment received (N and %)												
Quadruplets with anti-CD38 backbone												
Triplets with anti-CD38 backbone												
Other triplets												
Doublets/monotherapies												
Pomalidomide aCD38 backbone												
BCMA												
Bispecifics												
Other novel agents (e.g., Selinexor)												
Common sequences received												
Total number of unique regimens												
Regimen 1												
Regimen 2												
Regimen 3												
Regimen 4												
Regimen 5												