

A	B	C	D	E	F	G	H	I	J
+ BASELINE	MM_baseline_diag	Evidence of any MM during the baseline period	Binary	Evidence of any MM (each of newly diagnosed and R/R MM), during the baseline period =1; else =0;	Tab 42 & 49	Evidence of any MM here is defined use ≥1 non-diagnostic medical claim for MM (ICD-9-CM=203.0x) or ICD-10-CM code=C90.0x) during the baseline period. This flag is used to help better identify smoldering patients. From Optum: "When we refer to a non-diagnostic multiple myeloma (or any disease) claim, what we are referring to is a claim where one of the multiple myeloma diagnosis codes is present, but there is at least one service line on the claim that doesn't equal a diagnostic code. The idea is that a physician/facility might order labs for testing/diagnostics for multiple myeloma and on the claim they have to put something for the diagnosis so they will put what they ordered the lab to test for. e.g., diagnosis code for multiple myeloma. If the resulting facility claim had a diagnosis code for multiple myeloma and only one service line on the claim and that service line indicated a laboratory test, then we can't be certain that this is a true multiple myeloma diagnosis because the claim is suggesting that only testing was done. Alternatively, if the claim had the diagnosis codes for multiple myeloma and two service lines: the first line for testing and the second line a HCPCS code for medication or a CPT code for physician care management then we will accept that claim as a "non-diagnostic claim" indicating multiple myeloma."		OK19012026	
BASELINE	MM_baseline_other	Evidence of another cancer in the baseline period	Binary	Evidence of another cancer in the baseline period =1; else =0;	Tab 44	Patients with ≥ 2 ICD-9-CM or ICD-10-CM codes on separate days, within 30 days, for the same primary tumor type and/or metastatic cancer. Only the first of the 2 codes is required to occur inside the baseline period.		OK19012026	
ID_PD	Pregnant	Evidence of pregnancy	Binary	Evidence of pregnancy =1; else =0;	Tab 45	≥ 1 of medical claim with a diagnosis, procedure, or revenue code indicating pregnancy or childbirth during the patient identification period.		OK19012026	
ID_PD	CT_part	Clinical trial participation	Binary	Clinical trial participation =1; else =0;	Tab 46	Evidence of clinical trial participation during the patient identification will be flagged.		OK19012026	
FU_PD	Death_dt	Death Date	Date	Date patient died.		When date of death is only available at month-level granularity, the date will be generalized to the middle of the month or year, respectively (e.g., a date of death in June 2014 will be treated as 15 June 2014). For patients with only death year available, the death date is assumed to be July 15th year of the year they died. For patients with only death year available and their year of death is the same year as their index date, the death date is assumed to be July 15th of that year if the index date is on or before July 15th. The death date is assumed to be December 31st of that year if the index date is after July 15th.		OK19012026	
FU_PD	ENDDATE	Follow Up End Date	Date	min(Death_dt, study period end date)		Note: When the flag CE_3mos=1 is applied health plan disenrollment or the study period for patients who do not have have an assigned date of death <3mos after the index date.		OK19012026	
FU_PD	ENDDATE_CE	Follow Up End Date among CE pts	Date	min(Death_dt, date of health plan disenrollment, study period end date)		When the flag CE_3mos=1 is applied, follow up end date will be earliest of death date, health plan disenrollment or the study period for patients who do not have have an assigned date of death <3mos after the index date.		OK19012026	

A	B	C	D	E	F	G	H	I	J
ELUG_COH +	A patient level dataset containing those patients for whom we would like to run the multiple myeloma LOT algorithm on. It is used as an input into the multiple myeloma LOT algorithm.		The team will begin with the requirement that patients have at least one inpatient medical claim with a diagnosis code for multiple myeloma (any ICD-9-CM=203.0x or ICD-10-CM code=C90.0x) OR ≥ 2 outpatient medical claims for multiple myeloma (any ICD-9-CM=203.x or ICD-10-CM code=C90.x) in any position on claim, on separate days within <u>90 days</u> , during identification period. The team will apply all other IE criteria with the exception of the following exclusion criteria: 1) Evidence of another cancer (criteria 7) 2) Pregnancy or childbirth during the baseline or follow-up period (criteria 8) 3) Evidence of RCT participation during the baseline and/or follow-up periods (criteria 9) *Patients that meet exclusion criteria 1-3 indicated here will be included in this build of the algorithm.		Apply each criteria as a flag that can be carried forward or merged onto a later dataset. Helps make the cohort dynamic and flexible to change future study requirements. Suggest to establish global variables upfront when creating inputs for the study population, or other approaches to make the initial inputs easy to change. Suggest to implement IE criteria using a function where follow up period, allowable gaps, etc. can be modified. There is only 1 day of CE required in the follow up period for this initial cohort build. Patients with ≥ 2 outpatient medical claims for multiple myeloma (any ICD-9-CM=203.0x or ICD-10-CM code=C90.0x) in any position on claim, on separate days within 30 and 60 days during identification period will be flagged in the attrition table as well. ELUG_COH dataset should include patient ID, index date, death date, study end date, number of follow up days, and all IE criteria flags should be retained.			OK19012026	