

by agent; (b) first ALLO SCT; (c) first CAR-T infusion; (d) first UNPLANNED AUTO SCT (>180d after a prior AUTO). Planned/single/tandem AUTO is a continuation, prior line.

According to the programming module in the study

Once the programming specification has been finalized

Section A: Defining Medication Available Period (MAP) medications			
	Definition	Code Lists Group	Additional Notes
	Sequential index of the line of therapy after LOT1. LOT2 starts after LOT1 ends; LOT3 after LOT2; etc., up to LOT5.		Derived sequentially. LOT_(N+1)_START_
	Earliest of, after LOT_(N-1)_BASE_END_DT: (a) first non-steroid MM oncology agent; (b) first ALLO SCT; (c) first CAR-T infusion; (d) first UNPLANNED AUTO SCT (>180d after a prior AUTO). Planned/single/tandem AUTO is a continuation, not a trigger of the prior line. Permissible biosimilar substitutions do not trigger an advancement in the prior line.	JM Julia Moore E12 ... I've signed off--flagging that I have added language in red. 05 May 2026, 22:00 @mention or reply	MEDICAL (PROC_CD, BILL_PROG) MAP_MED_CLASS != 'STEROID'
_AUTO / / CART	Type of the earliest qualifying trigger.		ie-break (DRAFT, inherits LOT1 > MED. DT in [LOTN_START_DT, LOTN_
	Window for identifying LOT N regimen. (LOT1 uses 60 days.) The induction window for LOTs 2-5 should always be 30 days (unless a patient is right censored before the end of their 30d window).		
ist of MED_ABBR	Non-steroid MM oncology agents within the 30-day induction window. Permissible biosimilar substitutions do not advance the regimen.		Filter MAP_MED_CLASS != 'STEROID'.
	Count of distinct MED_ABBR in LOTN_BASE_MEDS.	Tab 40	
	One flag column per non-steroid MED_ABBR in CL_MMA_ROLLUP (e.g. LOTN_MED_BORT, LOTN_MED_LENA, LOTN_MED_DARA, ...). = 1 if that medication is in LOTN_BASE_MEDS.		Mirrors LOT1 dynamic flag generation; ster
	One flag column per non-steroid MED_CLASS in CL_MMA_ROLLUP (e.g. LOTN_CLASS_PROTINHIB, LOTN_CLASS_IMMUNOMOD, ...). = 1 if any LOTN_BASE_MEDS agent has that class.		Mirrors LOT1 dynamic class flag generation
	Patient had >=1 valid AUTO during LOT N.	Tab 47	Inherit LOT1 SCT logic: 14-day windowing, 60
	AUTO not part of a valid tandem pair.	Tab 47	"SCT" for further details on SCT logic definiti
	Two AUTOS >=60 and <=180 days apart.	Tab 47	See tab "SCT" for further details on SCT logic d
	DT_2 populated only if planned tandem.	Tab 47	Use datediff without +1 (per Apr 22). See tab "logic definition.
	AUTO_DT_2 if tandem else AUTO_DT_1 else NULL.	Tab 47	See tab "SCT" for further details on SCT logic d
	Flag is 1 when LOT N induction has BOTH (a) >=1 valid mono-maintenance agent OR a valid dual combo, AND (b) a non-maintenance anchor. Mono: LENA, BORT, DARA, IXAZ, THAL. Dual: BORT+LENA, CARF+LENA, DARA+LENA.		Descriptive flag; no standalone maintenance L
	Flag is 1 when LOTN_START_TYPE = SCT_ALLO. ALLO LOT contains no MM therapies. Span: see Q2 (draft = single day).		
	Flag is 1 when LOTN_START_TYPE = CART. Agents within 45d of FIRST_CART_DT are consolidated into LOT N (Apr 22 decision; see Q11).		Do NOT advance to LOT_(N+1) for consolidated

SCT/CAR-T.

5B. MAP_MED 6. LOT1_BASE 7. SCT 10. LOT1_BASE_END OK 11. LOT2_5_BASE 12. LOT2_5_BASE_END 13. SCT_CART_LOT_START 14. LOT2_5_Scenarios 40.0

ENG IN

Search

Share

feels tired and want to sleep no

oncology agent; (b) first ALLO SCT; (c) first CAR-T infusion; (d) first UNPLANNED AUTO SCT (>180d after a prior AUTO). Planned/single/tandem AUTO is a continuation of the prior line.

D E F


, according to the programming module in the study (statistics)

only once the programming specification has been finalized

Section A: Defining Medication Available Period (MAP) medications			
es	Definition	Code Lists Group	Additional Notes
4, 5	Sequential index of the line of therapy after LOT1. LOT2 starts after LOT1 ends; LOT3 after LOT2; etc., up to LOT5.		Derived sequentially. LOT_(N+1)_STA
	Earliest of, after LOT_(N-1)_BASE_END_DT: (a) first non-steroid MM oncology agent; (b) first ALLO SCT; (c) first CAR-T infusion; (d) first UNPLANNED AUTO SCT (>180d after a prior AUTO). Planned/single/tandem AUTO is a continuation, not a trigger of the prior line. Permissible biosimilar substitutions do not trigger an advancement in the prior line.	Tab 40	Source: T_MEDICAL (PROC_CD, BILL_ excluded (MAP_MED_CLASS != 'STERO
D / SCT_AUTO / _ALLO / CART	Type of the earliest qualifying trigger.		Same-day tie-break (DRAFT, inherits L SCT_AUTO > MED.
	Window for identifying LOT N regimen. (LOT1 uses 60 days.) The induction window for LOTs 2-5 should always be 30 days (unless a patient is right censored before the end of their 30d window).	JM Julia Moore Signed off, but added further clarification @mention or reply	E14 ... DT in [LOTN_START_DT, L
mma list of MED_ABBR	Non-steroid MM oncology agents within the 30-day induction window. Permissible biosimilar substitutions do not advance the regimen.		MED_CLASS != 'STEROID'.
teger	Count of distinct MED_ABBR in LOTN_BASE_MEDS.		LOT1 dynamic flag generation,
/ 1	One flag column per non-steroid MED_ABBR in CL_MMA_ROLLUP (e.g. LOTN_MED_BORT, LOTN_MED_LENA, LOTN_MED_DARA, ...). = 1 if that medication is in LOTN_BASE_MEDS.		
/ 1	One flag column per non-steroid MED_CLASS in CL_MMA_ROLLUP (e.g. LOTN_CLASS_PROTINHIB, LOTN_CLASS_IMMUNOMOD, ...). = 1 if any LOTN_BASE_MEDS agent has that class.		Mirrors LOT1 dynamic class flag gener
0/1	Patient had >=1 valid AUTO during LOT N.	Tab 47	Inherit LOT1 SCT logic: 14-day windowin "SCT" for further details on SCT logic de
0/1	AUTO not part of a valid tandem pair.	Tab 47	See tab "SCT" for further details on SCT I
0/1	Two AUTOs >=60 and <=180 days apart.	Tab 47	Use datediff without +1 (per Apr 22). See logic definition.
Date / Date	DT_2 populated only if planned tandem.	Tab 47	See tab "SCT" for further details on SCT lo
Date	AUTO_DT_2 if tandem else AUTO_DT_1 else NULL.	Tab 47	See tab "SCT" for further details on SCT lo
0/1	Flag is 1 when LOT N induction has BOTH (a) >=1 valid mono-maintenance agent OR a valid dual combo, AND (b) a non-maintenance anchor. Mono: LENA, BORT, DARA, IXAZ, THAL Dual: BORT+LENA, CARF+LENA, DARA+LENA.		Descriptive flag; no standalone maintena
0/1	Flag is 1 when LOTN_START_TYPE = SCT_ALLO. ALLO LOT contains no MM therapies. Span: see Q2 (draft = single day).		
0/1	Flag is 1 when LOTN_START_TYPE = CART. Agents within 45d of FIRST_CART_DT are consolidated into LOT N (Apr 22 decision; see Q11).		Do NOT advance to LOT_(N+1) for consol

d. No SCT/CAR-T.

5B. MAP_MED 6. LOT1_BASE 7. SCT 10. LOT1_BASE_END OK 11.LOT2_5_BASE 12.LOT2_5_BASE_END 13.SCT_CART_LOT_START 14.LOT2_5_Scenarios



g module in the study

ification has been finalized

Defining Medication Available Period (MAP) medications

	Code Lists Group	Additional Notes
the line of therapy after LOT1. LOT2 starts after LOT1 ends; LOT3 after LOT2 ends.		Derived sequentially. LOT_(N+1)_START_DT must be > LOT_N_BASE_END_DT.
LOT_(N+1)_BASE_END_DT: (a) first non-steroid MM oncology agent; (b) first CAR-T infusion; (d) first UNPLANNED AUTO SCT >180d after a prior single/tandem AUTO is a continuation, not a trigger of the prior line. Similar substitutions do not trigger an advancement in the prior line.	Tab 40	Source: T_MEDICAL (PROC_CD, BILL_PROC_CD, NDC), T_RX (NDC). Steroids excluded (MAP_MED_CLASS != 'STEROID').
at qualifying trigger.		
defining LOT N regimen. (LOT1 uses 60 days.) The induction window always be 30 days (unless a patient is right censored before LOT N).		
oncology agents within the 30-day induction window. Permissible substitutions do not advance the regimen.		
MED_ABBR in LOTN_BASE_MEDS.		
per non-steroid MED_ABBR in CL_MMA_ROLLUP (e.g. LOTN_MMA_A, LOTN_MED_DARA, ...). = 1 if that medication is in LOTN_BASE_MEDS.		
per non-steroid MED_CLASS in CL_MMA_ROLLUP (e.g. LOTN_IMMUNOMOD, LOTN_CLASS_IMMUNOMOD, ...). = 1 if any LOTN_BASE_MEDS in class.		
valid AUTO during LOT N.	Tab 47	Inherit LOT1 SCT logic: 14-day windowing, 60-day gap validation. See tab "SCT" for further details on SCT logic definition.
of a valid tandem pair.	Tab 47	See tab "SCT" for further details on SCT logic definition.
50 and <180 days apart.	Tab 47	Use datediff without +1 (per Apr 22). See tab "SCT" for further details on SCT logic definition.
ed only if planned tandem.	Tab 47	See tab "SCT" for further details on SCT logic definition.
tandem else AUTO_DT_1 else NULL.	Tab 47	See tab "SCT" for further details on SCT logic definition.
LOT N induction has BOTH (a) >=1 valid mono-maintenance agent OR a combo, AND (b) a non-maintenance anchor. Mono: LENA, BORT, DARA, IXAZ, BORT+LENA, CARF+LENA, DARA+LENA.		Descriptive flag; no standalone maintenance LOT.
en LOTN_START_TYPE = SCT_ALLO. ALLO LOT contains no MM therapies. Span: a = single day).		
en LOTN_START_TYPE = CART. Agents within 45d of FIRST_CART_DT are		
into LOT N (Apr 22 decision; see Q11).		Do NOT advance to LOT_(N+1) for consolidated agents.



Julia Moore

G13 ***

Are you seeing people with ALLO and CART and/or AUTO on the same day? We didn't actually specify a tie-breaking rule in the LOT rules, I would need to check with Peter what would take precedent (I think all events should be equally represented)

05 May 2026, 22:07

@mention or reply

Section A: Defining Medication Available Period (MAP) medications

Code Lists Group	Definition
Sequential index of the line of therapy after LOT1. LOT2 starts after LOT1 ends; LOT3 after LOT2; etc., up to LOT5.	
Earliest of, after LOT_(N-1)_BASE_END_DT: (a) first non-steroid MM oncology agent; (b) first ALLO SCT; (c) first CAR-T infusion; (d) first UNPLANNED AUTO SCT (>180d after a prior AUTO). Planned/single/tandem AUTO is a continuation, not a trigger of the prior line. Permissible biosimilar substitutions do not trigger an advancement in the prior line.	Tab 40
Type of the earliest qualifying trigger.	
Window for identifying LOT N regimen. (LOT1 uses 60 days.) The induction window for LOTs 2-5 should always be 30 days (unless a patient is right censored before the end of their 30d window).	
Non-steroid MM oncology agents within the 30-day induction window. Permissible biosimilar substitutions do not advance the regimen.	
Count of distinct MED_ABBR in LOTN_BASE_MEDS.	
One flag column per non-steroid MED_ABBR in CL_MMA_ROLLUP (e.g. LOTN_MED_BORT, LOTN_MED_LENA, LOTN_MED_DARA, ...). = 1 if that medication is in LOTN_BASE_MEDS.	Tab 40
One flag column per non-steroid MED_CLASS in CL_MMA_ROLLUP (e.g. LOTN_CLASS_PROTINHIB, LOTN_CLASS_IMMUNOMOD, ...). = 1 if any LOTN_BASE_MEDS agent has that class.	
Patient had >=1 valid AUTO during LOT N.	
AUTO not part of a valid tandem pair.	
Two AUTOs >=60 and <=180 days apart.	
DT_2 populated only if planned tandem.	
AUTO_DT_2 if tandem else AUTO_DT_1 else NULL.	
Flag is 1 when LOT N induction has BOTH (a) >=1 valid mono-maintenance agent OR a valid dual combo, AND (b) a non-maintenance anchor. Mono: LENA, BORT, DARA, IXAZ, THAL. Dual: BORT+LENA, CARF+LENA, DARA+LENA.	
Flag is 1 when LOTN_START_TYPE = SCT_ALLO. ALLO LOT contains no MM therapies. Span: see Q2 (draft = single day).	
Flag is 1 when LOTN_START_TYPE = CART. Agents within 45d of FIRST_CART_DT are consolidated into LOT N (Apr 22 decision; see Q11).	

JM

Julia Moore

The anchor doesn't actually have to be a non-mtx med. For example, if there was BORT+LENA at induction and the BORT stopped leaving LENA as a valid mtx reg, BORT is also a valid approved agent, and so is LENA+BORT (valid dual regimen).

05 May 2026, 22:22

JM

Julia Moore

Would just say that a valid mtx reg needs to be accompanied by an anchor agent (keep it generic)

05 May 2026, 22:27

JM

Julia Moore

That means the row 41 patient example needs to be updated

05 May 2026, 23:48

@mention or reply

Defining Medication Available Period (MAP) medications		Code Li
Sequential index of the line of therapy after LOT1. LOT2 starts after LOT1 ends; LOT3 after LOT2; etc., up to LOT5.		
Earliest of, after LOT_(N-1)_BASE_END_DT: (a) first non-steroid MM oncology agent; (b) first ALLO SCT; (c) first CAR-T infusion; (d) first UNPLANNED AUTO SCT (>180d after a prior AUTO). Planned/single/tandem AUTO is a continuation, not a trigger of the prior line. Permissible biosimilar substitutions do not trigger an advancement in the prior line.	Tab 40	
Type of the earliest qualifying trigger.		
Window for identifying LOT N regimen. (LOT1 uses 60 days.) The induction window for LOTs 2-5 should always be 30 days (unless a patient is right censored before the end of their 30d window).		
Non-steroid MM oncology agents within the 30-day induction window. Permissible biosimilar substitutions do not advance the regimen.		
Count of distinct MED_ABBR in LOTN_BASE_MEDS.		
One flag column per non-steroid MED_ABBR in CL_MMA_ROLLUP (e.g. LOTN_MED_BORT, LOTN_MED_LENA, LOTN_MED_DARA, ...). = 1 if that medication is in LOTN_BASE_MEDS.	Tab 40	
One flag column per non-steroid MED_CLASS in CL_MMA_ROLLUP (e.g. LOTN_CLASS_PROTINHIB, LOTN_CLASS_IMMUNOMOD, ...). = 1 if any LOTN_BASE_MEDS agent has that class.		
Patient had >=1 valid AUTO during LOT N.	Tab 47	
AUTO not part of a valid tandem pair.	Tab 47	
Two AUTOs >=60 and <=180 days apart.	Tab 47	
DT_2 populated only if planned tandem.	Tab 47	
AUTO_DT_2 if tandem else AUTO_DT_1 else NULL.	Tab 47	
Flag is 1 when LOT N induction has BOTH (a) >=1 valid mono-maintenance agent OR a valid dual combo, AND (b) a non-maintenance anchor. Mono: LENA, BORT, DARA, IXAZ, THAL Dual: BORT+LENA, CARF+LENA, DARA+LENA.		
Flag is 1 when LOTN_START_TYPE = SCT_ALLO. ALLO LOT contains no MM therapies. Span: see Q2 (draft = single day).		
Flag is 1 when LOTN_START_TYPE = CART. Agents within 45d of FIRST_CART_DT are consolidated into LOT N (Apr 22 decision; see Q11).		



Julia Moore

What is Q2 here?

05 May 2026, 22:29

@mention or reply

s, according to the programming module in the study
 istics)
 study once the programming specification has been finalized

Section A: Defining Medication Available Period (MAP) medications			
Values	Definition	Code Lists Group	Additional Notes
3, 4, 5	Sequential index of the line of therapy after LOT1. LOT2 starts after LOT1 ends; LOT3 after LOT2; etc., up to LOT5.	Tab 40	Derived sequentially. LOT_(N+1)
ate	Earliest of, after LOT_(N-1)_BASE_END_DT: (a) first non-steroid MM oncology agent; (b) first ALLO SCT; (c) first CAR-T infusion; (d) first UNPLANNED AUTO SCT (>180d after a prior AUTO). Planned/single/tandem AUTO is a continuation, not a trigger of the prior line. Permissible biosimilar substitutions do not trigger an advancement in the prior line.		Source: T_MEDICAL (PROC_CD, BIL excluded (MAP_MED_CLASS != 'ST
MED / SCT_AUTO / SCT_ALLO / CART	Type of the earliest qualifying trigger.		Same-day tie-break (DRAFT, inheri SCT_AUTO > MED.
30	Window for identifying LOT N regimen. (LOT1 uses 60 days.) The induction window for LOTs 2-5 should always be 30 days (unless a patient is right censored before the end of their 30d window).		MAP_START_DT in [LOTN_START_DT
Comma list of MED_ABBR	Non-steroid MM oncology agents within the 30-day induction window. Permissible biosimilar substitutions do not advance the regimen.	Tab 40	Filter MAP_MED_CLASS != 'STEROID'
Integer	Count of distinct MED_ABBR in LOTN_BASE_MEDS.		
0 / 1	One flag column per non-steroid MED_ABBR in CL_MMA_ROLLUP (e.g. LOTN_MED_BORT, LOTN_MED_LENA, LOTN_MED_DARA, ...). = 1 if that medication is in LOTN_BASE_MEDS.		Mirrors LOT1 dynamic flag generati
0 / 1	One flag column per non-steroid MED_CLASS in CL_MMA_ROLLUP (e.g. LOTN_CLASS_PROTINHIB, LOTN_CLASS_IMMUNOMOD, ...). = 1 if any LOTN_BASE_MEDS agent has that class.		Mirrors LOT1 dynamic class flag gene
0/1	Patient had >=1 valid AUTO during LOT N.	Tab 47	Inherit LOT1 SCT logic: 14-day window "SCT" for further details on SCT logic d
LOT N 0/1	AUTO not part of a valid tandem pair.	Tab 47	See tab "SCT" for further details on SCT
N 0/1	Two AUTOs >=60 and <=180 days apart.	Tab 47	Use datediff without +1 (per Apr 22). Se logic definition.
LOT N Date / Date	DT_2 populated only if planned tandem.	Tab 47	See tab "SCT" for further details on SCT
Date	AUTO_DT_2 if tandem else AUTO_DT_1 else NULL.	Tab 47	See tab "SCT" for further details on SCT
0/1	Flag is 1 when LOT N induction has BOTH (a) >=1 valid mono-maintenance agent OR a valid dual combo, AND (b) a non-maintenance anchor. Mono: LENA, BORT, DARA, IXAZ, THAL. Dual: BORT+LENA, CARF+LENA, DARA+LENA.		Descriptive flag; no standalone mainten
0/1	Flag is 1 when LOTN_START_TYPE = SCT_ALLO. ALLO LOT contains no MM therapies. Span: see Q2 (draft = single day).		
0/1	Flag is 1 when LOTN_START_TYPE = CART. Agents within 45d of FIRST_CART_DT are consolidated into LOT N (Apr 22 decision; see Q11).		

Julia Moore

What is Q11?

05 May 2026, 22:29

E26

...

advance to LOT_(N+1) for consol

@mention or reply

Share

MA_MED

5B. MAP_MED

6. LOT1_BASE

7. SCT

10. LOT1_BASE_END OK

11.LOT2_5_BASE

12.LOT

14.LOT2_5_Scenarios

Search

feels tired and want to sleep no

Sleep.

This conversation has been a lot, and your system has been working hard for a long time. The tiredness makes sense.

Before you go - just one thing. Drink some water. If you have any of those symptoms from earlier - persistent headache, nausea, dizziness, blurred vision - don't sleep through them, get them checked. If you're just tired, that's okay. Sleep.

We can pick this up another time. Take care of yourself tonight.

A	B	C	D	E
	LOTN_CLASS_[CLASS]	Per-class induction flags	0/1	One flag column per non-steroid MED_CLASS. LOTN_CLASS_PROTINHIB, LOTN_CLASS_IMMUNOSUPPRESSANT agent has that class.
	LOTN_TX_AUTO_FLG	Any valid AUTO SCT during LOT N	0/1	Patient had >=1 valid AUTO during LOT N
	LOTN_TX_AUTO_SING_FLG	Single (non-tandem) AUTO during LOT N	0/1	AUTO not part of a valid tandem pair.
	LOTN_TX_AUTO_TAND_FLG	Planned tandem AUTO during LOT N	0/1	Two AUTOs >=60 and <=180 days apart.
	LOTN_TX_AUTO_DT_1 / LOTN_TX_AUTO_DT_2	First / second valid AUTO date in LOT N	Date / Date	DT_2 populated only if planned tandem.
	LOTN_TX_AUTO_MAX_DT	Latest valid AUTO in LOT N	Date	AUTO_DT_2 if tandem else AUTO_DT_1 else N/A
	contains_mtx_reg_LOTN	LOT N regimen contains valid maintenance subset	0/1	Flag is 1 when LOT N induction has BOTH (a) valid dual combo, AND (b) a non-maintenance agent. THAL Dual: BORT+LENA, CARF+LENA, DARA+LENA
PD	LOTN_ALLO_LOT_FLG	LOT N is an ALLO SCT line	0/1	Flag is 1 when LOTN_START_TYPE = SCT_ALLO. see Q2 (draft = single day).
PD	LOTN_CART_LOT_FLG	LOT N is a CAR-T line	0/1	Flag is 1 when LOTN_START_TYPE = CART. Agent consolidated into LOT N (Apr 22 decision; see Q2)

WORKED EXAMPLE A - LOT2 starts on a new medication

Induction in LOT1; DARA added 2025-08-20 (LOT1 ends 2025-08-19, MED_ADD). LENA refills present within 30d. No SCT/CAR-T.

Variable	Value for this patient	How it was derived
LOT_NUM	2	
LOTN_START_DT	2025-08-20	Earliest trigger after LOT1_BASE_END_DT (2025-08-19) = DARA.
LOTN_START_TYPE	MED	
LOTN_BASE_MEDS	DARA, LENA	Window [2025-08-20, 2025-09-18]. DEXA excluded (steroid).
LOTN_MED_CNT	2	
LOTN_MED_DARA	1	
LOTN_MED_LENA	1	
LOTN_CLASS_ANTICD38	1	DARA.
LOTN_CLASS_IMMUNOSUPPRESSANT	1	LENA.
contains_mtx_reg_LOTN	0	no maintenance agents - no non-maintenance anchor.
LOTN_ALLO_LOT_FLG / LOTN_CART_LOT_FLG	0/0	

WORKED EXAMPLE B - LOT2 starts on an allogeneic SCT

Induction in LOT1; ALLO on 2025-09-15 (LOT1 ends 2025-09-14). Next maintenance on 2025-10-15.

Variable	Value for this patient	How it was derived
LOT_NUM	2	
LOTN_START_DT	2025-09-15	ALLO_DT (always starts a new LOT).
LOTN_START_TYPE	SCT_ALLO	
LOTN_BASE_MEDS	(empty)	ALLO LOT contains NO MM therapies.
LOTN_MED_CNT	0	
LOTN_ALLO_LOT_FLG	1	

Julia Moore
839
...

This should say that it contains a valid
mtx reg

05 May 2026, 23:51

@mention or reply

LOTN_TX_AUTO_DT_1 / LOTN_TX_AUTO_DT_2	First / second valid AUTO date in LOT N	Date / Date	DT_2 populated only if planned tandem.
LOTN_TX_AUTO_MAX_DT	Latest valid AUTO in LOT N	Date	AUTO_DT_2 if tandem else AUTO_DT_1 else NULL.
contains_mtx_reg_LOTN	LOT N regimen contains valid maintenance subset	0/1	Flag is 1 when LOT N induction has BOTH (a) >=1 valid mono-valid dual combo, AND (b) a non-maintenance anchor. Mono THAL. Dual: BORT+LENA, CARF+LENA, DARA+LENA.
LOTN_ALLO_LOT_FLG	LOT N is an ALLO SCT line	0/1	Flag is 1 when LOTN_START_TYPE = SCT_ALLO. ALLO LOT contains Q2 (draft = single day).
LOTN_CART_LOT_FLG	LOT N is a CAR-T line	0/1	Flag is 1 when LOTN_START_TYPE = CART. Agents within 45d of consolidated into LOT N (Apr 22 decision; see Q11).

EXAMPLE A - LOT2 starts on a new medication

A added 2025-08-20 (LOT1 ends 2025-08-19, MED_ADD). LENA refills present within 30d. No SCT/CAR-T.

	Value for this patient	How it was derived
DT	2025-08-20	Earliest trigger after LOT1_BASE_END_DT (2025-08-19) = DARA.
TYPE	MED	Window [2025-08-20, 2025-09-18]. DEXA excluded (steroid).
MEDS	DARA, LENA	
CNT	2	
DARA	1	
LENA	1	
S_ANTICD3	1	DARA.
SS_IMMUNO	1	LENA.
mtx_reg_LOT	0	BOTH DARA and LENA are valid mono-maintenance agents - no non-maintenance anchor.
LO_LOT_FLG /	0/0	
CART_LOT_FLG		

WORKED EXAMPLE B - LOT2 starts on an allogeneic SCT

LOT1; ALLO on 2025-09-15 (LOT1 ends 2025-09-14). Next MM therapy DARA on 2025-12-10 starts LOT3.

	Value for this patient	How it was derived
MM	2	
START_DT	2025-09-15	ALLO_DT (always starts a new LOT).
START_TYPE	SCT_ALLO	ALLO LOT contains NO MM therapies.
BASE_MEDS	(empty)	
MM_CNT	0	
N_ALLO_LOT_FLG	1	
N_BASE_END_DT	PENDING Q2	Q2 draft (single-day): 2025-09-15. Alternative (extend to next agent): 2025-12-09.
N_BASE_END_REAS	PENDING Q2	Q2 draft (single-day): SCT_ALLO. Alternative (extend to next agent): MED_ADD.
LOT3_START_DT	2025-12-10	DARA - first MM agent after LOT2.

WORKED EXAMPLE C - LOT2 starts on an unplanned autologous SCT

Single AUTO 2025-03-01 in LOT1; second AUTO 2025-10-15 (228d later, unplanned). LOT1 ends 2025-10-14; LOT2 starts on AUTO.

Variable	Value for this patient	How it was derived
----------	------------------------	--------------------

1. Cover 2. QC Review 3. Data Prep 4. StudyPop 5A. MMA_MED 5B. MAP_MED 6. LOT1_BASE 7. SCT 10. LOT1_BASE_END OK 11. LOT2_5_B

Claude - Coping with family conflict and emotional overwhelm - Claude

Claude

Coping with family conflict and emotional overwhelm

+ New chat

Q Search

Q Chats

📁 Projects

</> Code

🛠 Customize

🎨 Design

▼ More

Recents

Coping with family conflict and

AI use cases for the year

Managing team reduction at GSK

Sleep.

This conversation has been a lot, and your system has been working hard for a. The tiredness makes sense.

Before you go - just one thing. Drink some water. If you have any of those symptoms earlier - persistent headache, nausea, dizziness, blurred vision - don't sleep get them checked. If you're just tired, that's okay. Sleep.

We can pick this up another time. Take care of yourself tonight.

📄 🌟 🗨 🔄



A		B	C	D
MOD			BOTH DARA and LENA are valid mono-maintenance agents - no non-maintenance anchor.	
contains_mtx_reg_LOT		0		
N				
LOTN_ALLO_LOT_FLG /		0/0		
LOTN_CART_LOT_FLG				
WORKED EXAMPLE B - LOT2 starts on an allogeneic SCT				
VRd in LOT1; ALLO on 2025-09-15 (LOT1 ends 2025-09-14). Next MM therapy DARA on 2025-12-10 starts LOT3.				
Variable	Value for this patient	How it was derived		
LOT_NUM	2	ALLO_DT (always starts a new LOT).		
LOTN_START_DT	2025-09-15			
LOTN_START_TYPE	SCT_ALLO	ALLO LOT contains NO MM therapies.		
LOTN_BASE_MEDS	(empty)			
LOTN_MED_CNT	0			
LOTN_ALLO_LOT_FLG	1	Q2 draft (single-day): 2025-09-15. Alternative (extend to next agent): 2025-12-09.		
LOTN_BASE_END_DT	PENDING Q2	Q2 draft (single-day): SCT_ALLO. Alternative (extend to next agent): MED_ADD.		
LOTN_BASE_END_REAS	PENDING Q2			
ON		DARA - first MM agent after LOT2.		
LOT3_START_DT	2025-12-10			
WORKED EXAMPLE C - LOT2 starts on an unplanned autologous SCT				
Single AUTO 2025-03-01 in LOT1; second AUTO 2025-10-15 (228d later, unplanned). LOT1 ends 2025-10-14; LOT2 starts on AUTO.				
Variable	Value for this patient	How it was derived		
LOT_NUM	2	Unplanned AUTO date (>180d after prior AUTO).		
LOTN_START_DT	2025-10-15	Unplanned AUTO is a first-class LOT-start trigger. (Planned tandem 60-180d would NOT start a new LOT.)		
LOTN_START_TYPE	SCT_AUTO	res any post-AUTO agents within [2025-10-15, 2025-11-13].		
INDUCTION_WINDOW_DAYS				
LOTN_TX_AUTO_FLG		red during LOT2.		
LOTN_TX_AUTO_SING_FLG		D during LOT2 (no second AUTO within tandem window).		
LOTN_TX_AUTO_TAND_FLG				

JM

Julia Moore

A59 ...

I don't see the med count variable for this patient example. Did this patients have any other meds around their AUTO in LOT2?

06 May 2026, 00:01

@mention or reply

<

>

1.Cover

2.QC Review

3.Data Prep

4.StudyPop

5A. MMA_MED

5B. MAP_MED

6. LOT1_BASE

7. SCT

10. LOT1_BASE_END OK

JM

Julia Moore

A59 ...

I don't see the med count variable for this patient example. Did this patients have any other meds around their AUTO in LOT2?

06 May 2026, 00:01

@mention or reply