

members in NDMM cohort (1L cohort). Additional eligibility criteria will be applied to the 1L cohort to create the subset cohorts of patients who received 2+ (2L cohort) and 3+ (3L cohort) lines of therapy.

6.2.1.1. Inclusion Criteria

Yunji Hwang, 11-06-2026 21:00:00

commented:

Is this definition used in the publication that we can cite, just in case? It would be nice if we have a reference for this method.

Over *s defined for 1L NDMM cohort*

- **MM diagnosis:** At least one inpatient medical claim with a diagnosis code for MM in any position (any ICD-9-CM=203.0x or ICD-10-CM code=C90.0x) or ≥ 2 outpatient medical claims for MM in any position on the claim, on separate days within 90 days, during the study period
 - As data in Optum CDM is collected primarily for health insurance and not research purposes, all MM diagnosis codes are to be considered. Inclusion will focus on defining NDMM according to patients who are newly treated, defined as the receipt of a 1L treatment, and with no prior MM oncology therapy in the preceding 12 months (as described below)

- **Adult age:** Aged ≥ 18 years at the time of MM diagnosis according to calendar year
- **Eligible 1L treatment:** Received an eligible or expected treatment for MM on or after MM diagnosis (other than belantamab), occurring on or after 01 Jan 2017 (eligible treatment period).
 - The 1L cohort index date is the date of the first claim for MM treatment within the identification period
 - Eligible/expected treatments are defined as any treatment for MM except for treatments reserved strictly for later LOTs (see exclusion criteria).
 - For a full list of eligible/expected MM therapies, see Annex 2
- **Continuous enrollment (CE):** CE of at least 12-months with medical and pharmacy benefits before the 1L cohort index date. Patients with gaps in enrolment of ≤ 30 days are considered to be continuously enrolled
- **CE during follow-up:** CE of at least 3-months during follow-up or death with no gaps in enrollment

Additional eligibility for 2L and 3L RRMM Cohorts

The subset of patients with evidence of each subsequent LOT will be included in the 2L or 3L cohort at the initiation of 2L/3L.

To be eligible for each 2L or 3L cohorts, patients must meet the following criteria:

- **Received a subsequent LOT required to qualify for a specific cohort** (i.e., received a 2L treatment for 2L, 3L for 3L cohort)
- **Continuous enrollment (CE) for each cohort:** CE of at least 12-months with medical and pharmacy benefits before the cohort index date (2L or 3L). Patients with gaps in enrolment of ≤ 30 days are considered to be continuously enrolled
- **CE during follow-up for each cohort:** CE of at least 3-months during follow-up or death with no gaps in enrollment

6.2.1.2. Exclusion Criteria

- **Evidence of an MM oncology therapy during the 12-month 1L baseline period:** ≥ 1 medical or pharmacy claim for any MM oncology therapy

- This is to ensure that treatment exposure and LOT assignments reflect incident therapy starts at the index date and are not confounded by ongoing or recent prior MM treatments

• **Evidence of another cancer in the 1L baseline period:** Patients with either ≥ 1 inpatient or ≥ 2 outpatient ICD-9-CM or ICD-10-CM codes on separate days, within 30 days, for the same primary tumor type and/or metastatic cancer will be excluded

• **Evidence of pregnancy:** ≥ 1 of medical claim with a diagnosis, procedure, or revenue code indicating pregnancy or childbirth during the study period

• **Received belantamab mafodotin (i.e., an ADC) in any LOT**

- Note: at the time of study belantamab mafodotin was the only ADC in use for MM

6.2.2. SOC Categorization and Eligible 1L MM Regimens

To assess treatment patterns and time-to-event treatment outcomes according to LOT cohorts and SOC regimen, regimens will potentially be grouped according to commonly utilized quadruplets, triplets, doublets, anti-CD38 backbone, and class, allowing for potential off-label use.

purposes of inclusion criteria, eligible 1L treatments will include treatments commonly used or expected in the 1L setting, inclusive of approved, off-label combination regimens.

Annex 2 contains an exemplary list of potential treatment combinations that may be included in each group. Final treatment groupings may depend on data availability and distribution of observed treatments and may be re-categorized or summarized as 'other', where applicable.

Tentatively, the following SOC categories are proposed:

- **1L (NDMM):**

I

- Quadruplets with anti-CD38 backbone
- Triplets with anti-CD38 backbone
- Other triplets (non-anti-CD38)
- Doublets
- Monotherapies
- Other (if applicable)

- **Later lines (2L, 3L, 4L):**

- Quadruplets with anti-CD38 backbone (off-label; some use expected)