


Section A: Defining Medication Available Period (MAP) Medications									
Time Period	Variable	Label	Values	Definition	Concomitant Lists	Additional Notes		Date Modified (Initials + Date)	QC Reviewed (Initials + Date)
FU_PD	ENDDATE	The end date of a patient's observation period due to study end date, death, disenrollment, etc.	date	from the ELIG_COH dataset, merge ENDDATE onto the dataset via PAT_ID.				OK13APR2026	
FU_PD	LOT1_START_DT	LOT1 Start Date	date	min(MAP_START_DT) where MAP_MED_CLASS != STEROID		Patient level for LOT1_BASE dataset. Using the MAP_STACKED dataset, determine each patient's LOT1_START_DT by identifying the earliest date that a patient received an MMA medication. Note that LOT1_START_DT will be equal to at least one MAP_START_DT or potentially several if a patient started multiple [MED]s on the same day. In Figure 1 in Section B, the LOT1_START_DT is set by MAP [MED A]'s MAP_START_DT.		OK13APR2026	
FU_PD	LOT1_MED_[MED]	[MED] is an induction medication during LOT1 BASE.	0,1	For all [MED]s: if MAP_MED_TYPE=[MED] and LOT1_START_DT <= MAP_START_DT <= (LOT1_START_DT+59) then = 1 Else 0.		Identify induction regimen medications by using MAP_STACKED to identify medications occurring (MAP_START_DT) during the 60 day window of LOT1_START_DT + 59. A patient's LOT1 induction regimen is broadly defined as those MMA medications found during the 60 day window following the patient's earliest MMA medication claim. Once the induction regimen medications are identified, the induction regimen period is defined as the time period beginning with the earliest MMA medication claim and continuing until a valid medication add is introduced (see list of permissible substitutions in LOT1_BASE_MEDS), all induction regimen medications discontinue, censoring, HSCT occurs. The induction regimen and its corresponding period is referred to as LOT1 Base (see Figure 1 in Section B).		OK13APR2026	
FU_PD	LOT1_MED_CNT	Count of induction medications during LOT1 BASE	continuous	Set to the sum(LOT1_MED_[MED]) for all [MED]s. For example, if a patient has LOT1_MED_BORT=1 and LOT1_MED_LEN=1 and all other				OK13APR2026	

A	B	C	D	E	F	G	H	I
FU_PD	LOT1_MED_CNT	Count of induction medications during LOT1_BASE	continuous	Set to the sum(LOT1_MED_JMED) for all [MED]s. For example, if a patient has LOT1_MED_BORT=1 and LOT1_MED_LEN=1 and all other LOT1_MED_JMEDs = 0, then the value for this variable should be set to 2				OK13APR2026
FU_PD	LOT1_CLASS_CLASS	Presence of [CLASS] during LOT1_BASE	0,1	For each [MED], if LOT1_MED_JMED=1 then find the [MED]'s class [CLASS] and then set LOT1_CLASS_CLASS to 1. Else = 0 Example: LOT1_MED_BORT = 1 would indicate that LOT1_CLASS_PROTINHB should be set to 1.				OK13APR2026
FU_PD	LOT1_BASE_DISCON_DT	LOT1 Base Discontinue Date	date	A patient's temporary LOT1_BASE discontinuation date (LOT1_BASE_DISCON_DT) may be calculated using their induction regimen's MAPs and other MMA medications' MAPs that are permissible substitutions (see the additional notes column). Note that the calculated discontinuation date may be not mark the actual end date of LOT1_BASE (e.g., censoring). Obtain all induction regimen medication [MED] MAPs and all related permissible substitution [MED] MAPs that start on or after the established LOT1_Start_Dt and order them chronologically. Set the maximum MAP_END_DT observed as the LOT1_BASE_DISCON_DT. Obtain all MAPs from MAP_STACKED where (LOT1_MED_JMED=1 and MAP_MED_TYPE = [MED]) and MAP_START_DT >= LOT1_START_DT LOT1_BASE_DISCON_DT = max_map_end_dt. If the obtained discontinuation date is within 90 days of the patient's ENDDATE, then set the discontinuation date to missing as there was not enough follow-up time to determine if a patient's LOT1 Base period truly discontinued.				OK13APR2026

2

QC
(Unit



A B C D E F G H I

