

Section A: Defining Medication Available Period (MAP) medications									
Time Period	Variable	Label	Values	Definition	Code Lists Group	Additional Notes	Date Modified (Initials + Date)	Q	
FU_PD	ENDDATE	The end date of a patient's observation period due to study end date, death, disenrollment, etc.	date	from the ELUG_COH dataset, merge ENDDATE onto the dataset via PAT_ID.					
FU_PD	LOT1_START_DT	LOT1 Start Date	date	min(MAP_START_DT) where MAP_MED_CLASS != STEROID		Patient level for LOT1_BASE dataset. Using the MAP_STACKED dataset, determine each patient's LOT1_START_DT by identifying the earliest date that a patient received an MMA medication. Note that LOT1_START_DT will be equal to at least one MAP_START_DT or potentially several if a patient started multiple [MED]s on the same day. In Figure 1 in Section B , the LOT1_START_DT is set by MAP [MED A]'s MAP_START_DT.			
						Patient level for LOT1_BASE dataset. Identify induction regimen medications by using MAP_STACKED to identify medications occurring (MAP_START_DT) during the 60 day window of LOT1_START_DT + 59. A patient's <u>LOT1</u> induction regimen is broadly defined as those MMA medications found during the 60 day window following the patient's earliest MMA medication claim. Once the induction regimen medications are identified, the induction regimen period is defined as the time period beginning with the earliest MMA medication claim and continuing until a valid medication add is introduced (see list of permissible substitutions in LOT1_BASE_MEDS), all induction regimen medications discontinue, censoring, HSCT, or maintenance begins. The induction regimen and its corresponding period is referred to as LOT1 Base (see Figure 1 in Section B).			
FU_PD	LOT1_MED_[MED]	[MED] is an induction medication during LOT1 BASE.	0, 1	For all [MED]s: if MAP_MED_TYPE=[MED] and LOT1_START_DT <= MAP_START_DT <= (LOT1_START_DT+59) then = 1 Else 0					
13									

				A patient's temporary LOT1 BASE discontinuation date (LOT1_BASE_DISCON_DT) may be calculated using their induction regimen's MAPs and other MMA medications' MAPs that are permissible substitutions (see the additional notes column). Note that the calculated discontinuation date may be not mark the actual end date of LOT1 BASE (e.g., censoring).			
				Obtain all induction regimen medication [MED] MAPs and all related permissible substitution [MED] MAPs that start on or after the established LOT1_Start_Dt and order them chronologically. Set the maximum MAP_END_DT observed as the LOT1_BASE_DISCON_DT.		If the obtained discontinuation date is within 90 days of the patient's ENDDATE, then set the discontinuation date to missing as there was not enough follow-up time to determine if a patient's LOT1 Base period truly discontinued.	
				Obtain all MAPs from MAP STACKED where (LOT1_MED_MED)=1 and MAP_MED_TYPE = [MED] and MAP_START_DT >= LOT1_START_DT			
				LOT1_BASE_DISCON_DT = max_map_end_dt.			
				If ENDDATE - LOT1_BASE_DISCON_DT < 90 then LOT1 BASE DISCON DT =			

				For each LOT1_MED [MED]=1, create a string of the medication abbreviations separated by a space.		For example, if LOT1_MED_BORT=1 and LOT1_MED_LENA=1 and LOT1_MED_CARF=1 then this variable should equal "BORT LENA CARF".	
						Consider the list of permissible substitutions within a line, and that the line will ultimately be named according to the drug given for the longest duration within the line:	
						• Substitution of rituximab for rituximab/hyaluronidase and vice versa does not advance the LOT	
						• Substitution of daratumumab for daratumumab/hyaluronidase and vice versa does not advance the LOT	
						• Substitution of the reference product for a biosimilar or vice versa does not advance the LOT (e.g, trastuzumab and trastuzumab-anns)	
						• Substitution of bortezomib induction therapy for ixazomib as a maintenance therapy	
				If not missing(LOT1_BASE_DISCON_DT) then set to LOT1_BASE_DISCON_DT - LOT1_START_DT + 1			
				Otherwise if missing(LOT1_BASE_DISCON_DT) then set to ENDDATE - LOT1_START_DT + 1			

